


# BREAKING WATER WAVES

## A numerical study with viscosity

 Alan Riquier<sup>†,\*</sup> & Emmanuel Dormy<sup>†</sup>

### Model description



**Figure 1.** Geometry of the initial domain.

We solve (numerically) the non-dimensional incompressible Navier-Stokes equation in the moving domain  $\Omega_t$ . **Navier** conditions are used at the bottom  $\Gamma_0$  while **Stress-free** conditions are used on the water-air interface,


$$\begin{cases} \frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla p - \frac{1}{Re} \Delta \mathbf{u} = -\hat{\mathbf{y}} & \text{in } (0, T) \times \Omega_t \\ \nabla \cdot \mathbf{u} = 0 & \text{in } (0, T) \times \Omega_t \\ \mathbf{u} \cdot \hat{\mathbf{n}} = 0 & \text{on } (0, T) \times \Gamma_0 \\ \hat{\mathbf{t}} \cdot (\nabla \mathbf{u} + (\nabla \mathbf{u})^t) \cdot \hat{\mathbf{n}} = 0 & \text{on } (0, T) \times \Gamma_0 \\ -p \hat{\mathbf{n}} + \frac{1}{Re} (\nabla \mathbf{u} + (\nabla \mathbf{u})^t) \cdot \hat{\mathbf{n}} = 0 & \text{on } (0, T) \times \Gamma_{i,t} \\ \mathbf{u}(0, \cdot) = \nabla \phi_0 & \text{in } \Omega_0 \end{cases}.$$

The **initial potential**  $\phi_0$  is a harmonic function whose value on the interface is given by a finite amplitude extension of the linear wave solution,

$$\frac{\partial \phi_0}{\partial \hat{\mathbf{n}}}(x) = \mathbf{u}(0, x) \cdot \hat{\mathbf{n}} = a \sqrt{gk} \tanh(kh_0) \cdot \left[ \begin{array}{c} (\tanh kh_0)^{-1} \cos kx \\ \sin kx \end{array} \right].$$


The interface is advected with the fluid.

### Where does the viscous dissipation happens?




**Figure 4.** (a-e) The vorticity  $\omega = \partial_x u_y - \partial_y u_x$  at time  $t = 2.9$ . We see that the viscous dissipation is happening in a boundary layer below the surface of size  $\delta \approx Re^{-\frac{1}{2}}$ . Solid black lines correspond to vorticity cuts (not shown here). (f) A zoom on a positive vorticity region in the  $Re = 10^5$  case.

### Evolution of a Water Wave at $Re = 10^6$



**Figure 2.** Evolution of a wave of initial amplitude  $a = 0.5$  at  $Re = 10^6$ .

### Convergence to the inviscid solution



**Figure 3.** Influence of the viscosity on the shape of the wave as time increases. The  $Re = +\infty$  simulation has been achieved using an Euler-based code (see [1] for details).

<sup>†</sup> Département de Mathématiques et Applications, CNRS UMR-8553, ENS-PSL, 45 rue d'Ulm, 75005 Paris, France.

\* Corresp. email: alan.riquier@ens.psl.eu

This work makes use of the FreeFEM finite-element library [2].

[1] Dormy, E. & Lacave, C., **Inviscid Water-Waves and interface modeling**, accepted in Quarterly of Applied Mathematics, 2023.

[2] Hecht, F., **New development in freefem++**, J. Numer. Math. 20(3-4), 2012.