

Name:

NetId:

## Honors Ordinary Differential Equations

Midterm Exam, Fall 2021

**DO NOT OPEN YET**

**...and wait until the proctor announces that it is time to start.**

In the mean time, please write your name and NetID legibly,  
and **read the instructions below carefully.**

- \* Please do not fold or damage the exam papers. After you finish your exam, please put the pages in correct order back into the sleeve.
- \* There are 5 questions in this exam, the sleeve should contain 8 pieces of paper.
- \* The scratch paper is included: three last papers are blank. If your solution continues on scratch paper, please clearly indicate it.
- \* For questions asking to prove a result, the clarity of the mathematical argument will be taken into account in the score.
- \* Questions formulated in terms of real functions should be answered with real functions.
- \* Question marked with (†) is challenge.

**Good luck!**



1. Answer whether the statement is **TRUE** or **FALSE**.

(a) **TRUE / FALSE:**  $ty'' + t^2y' = t^3$  is a second order differential equation.

(b) **TRUE / FALSE:**  $y' + \sin(t)y = e^t$  is a non-linear differential equation.

(c) **TRUE / FALSE:** If  $y_1$  and  $y_2$  are solutions to the differential equation

$$y'' - t^2y' + 5y = e^t \sin(t),$$

then  $y_1 + y_2$  is also a solution of this equation.

(d) **TRUE / FALSE:** The initial value problem  $y'' + y = \sin(10t)$  has a unique solution on  $\mathbb{R}$  for  $y(0) = 0, y'(0) = 1$ .

(e) **TRUE / FALSE:** The initial value problem  $y' - \frac{1}{1+|t|}y = t^{\frac{1}{3}}, y(0) = 0, t \geq 0$  has more than one solution.

(f) **TRUE / FALSE:** Assume that  $p$  and  $q$  are continuous and that the functions  $y_1$  and  $y_2$  are solutions of the differential equation  $y'' + p(t)y' + q(t)y = 0$  on an open interval  $I$ . If  $y_1$  and  $y_2$  are zero at the same point in  $I$ , then  $y_1 = \lambda y_2$  for some  $\lambda \in \mathbb{R}$ .

(g) **TRUE / FALSE:** If the power series  $\sum_{k=0}^{\infty} a_k(x - x_0)^k$  has radius of convergence  $\rho$ , then  $\sum_{k=0}^{\infty} k a_k(x - x_0)^k$  also has a radius of convergence  $\rho$ .

(h) **TRUE / FALSE:** If the power series  $\sum_{k=0}^{\infty} a_k(x - x_0)^k$  has radius of convergence  $\rho$ , then  $\{\rho^n a_n\}_{n \in \mathbb{N}}$  is bounded.

2. Consider the differential equation

$$y' = p(t)y + \cos t, \tag{1}$$

where  $p$  is a continuous function on  $\mathbb{R}$ .

- (a) Show carefully that there exists a unique solution for that equation with initial value  $y(0) = y_0$ , and determine it explicitly.
- (b) ((††), could be considered later in the exam) Prove that, if  $y_0 > 1$ ,  $p(t) \geq 1$  on  $\mathbb{R}$ , then  $\lim_{t \rightarrow +\infty} y(t) = +\infty$ .
- (c) Now assume that  $p$  is the constant function  $p = 1$ . Show that there exists  $y_0 \in \mathbb{R}$  such that  $y(0) = y_0$  and  $y$  is a periodic solution, and prove carefully such  $y_0$  is unique.



3. Given continuous functions  $p$  and  $q$  on an open interval  $I$  containing  $t_0$ , let  $y_1$  and  $y_2$  be two solutions of the differential equation

$$y'' + p(t)y' + q(t)y = 0 \tag{2}$$

on that open interval  $I$ .

- (a) Define the Wronskian of the two solutions  $y_1$  and  $y_2$ .
- (b) State the theorem relating Wronskian to Fundamental Set of Solutions (FSS).
- (c) Write down (without proof) the expression of  $W_{[y_1, y_2]}(t)$  as a function of  $W_{[y_1, y_2]}(t_0)$ , for  $t_0, t \in I$ .
- (d) If  $y_1(t_0) = 1, y_1'(t_0) = 1, y_2(t_0) = 1, y_2'(t_0) = -1$ , write down the expression of the solution  $y$  of the initial value problem (2) with  $y(t_0) = a, y'(t_0) = b$ . Justify your answer.
- (e) Assume that, given  $\alpha \in \mathbb{R}$ , the functions  $p$  and  $q$  are constant equal to

$$\begin{aligned} p(t) &= -4, \\ q(t) &= -\alpha^2 + 2\alpha + 3. \end{aligned}$$

Compute the general solution. Determine the set of  $\alpha \in \mathbb{R}$  such that there exists a bounded non-zero solution to the ordinary differential equation (2).



4. The aim of this question is to prove Grönwall's inequality, which can be stated as follows.

Let  $a, y : \mathbb{R} \rightarrow [0, \infty)$  be two continuous nonnegative functions and suppose that, for any  $t \geq 0$ ,

$$y(t) \leq 1 + \int_0^t a(s)y(s) ds. \quad (3)$$

Then

$$y(t) \leq \exp\left(\int_0^t a(s) ds\right). \quad (4)$$

In the rest of the exercise, we assume (3), with the aim to prove (4).

- (a) Define  $w(t) := 1 + \int_0^t a(s)y(s) ds$ . Prove that  $w$  is a differentiable function, and compute  $w'(t)$ .
- (b) Plugging (3) in the expression for  $w'(t)$ , write down a differential inequality using only  $w'(t), w(t), a(t)$ .
- (c) Deduce from (b) an upper bound estimate for  $w(t)$ .
- (d) Conclude (4).



5. (a) Let  $\sum_{n=0}^{\infty} a_n x^n$  and  $\sum_{n=0}^{\infty} b_n x^n$  be two series with respectively radii of convergence  $\rho_1$  and  $\rho_2$ . Prove that the radius of convergence  $R$  of  $\sum_{n=0}^{\infty} a_n b_n x^n$  satisfies  $R \geq \rho_1 \rho_2$ . Do we always have equality ?

- (b) Consider the series

$$f(x) \rightarrow \sum_{n=1}^{\infty} \sin\left(\frac{1}{\sqrt{n}}\right) x^n. \quad (5)$$

- i. Determine the radius of convergence  $\rho$  of this series.
- ii. Study if this series converges at  $\rho$  and  $-\rho$ .
- iii. ((†), could be considered later in the exam) Prove that, for any  $M > 0$ , there exists  $\delta > 0$  and  $N \in \mathbb{N}$  such that for any  $x \in (1 - \delta, 1)$ , we have

$$\sum_{n=1}^N \sin\left(\frac{1}{\sqrt{n}}\right) x^n \geq M. \quad (6)$$

- iv. Compute the limit  $\lim_{x \rightarrow 1^-} f(x)$ .

