

UN PRINCIPE D'AX-KOCHEN-ERSHOV IMAGINAIRE

MARTIN HILS AND SILVAIN RIDEAU-KIKUCHI

ABSTRACT. We study interpretable sets in henselian and σ -henselian valued fields with value group elementarily equivalent to \mathbb{Q} or \mathbb{Z} . Our first result is an Ax-Kochen-Ershov type principle for weak elimination of imaginaries in finitely ramified characteristic zero henselian fields — relative to value group imaginaries and residual linear imaginaries. We extend this result to the valued difference context and show, in particular, that existentially closed equicharacteristic zero multiplicative difference valued fields eliminate imaginaries in the geometric sorts; the ω -increasing case corresponds to the theory of the non-standard Frobenius automorphism acting on an algebraically closed valued field.

On the way, we establish some auxiliary results on separated pairs of characteristic zero henselian fields and on imaginaries in linear structures which are also of independent interest.

1. INTRODUCTION

In his seminal work “Une théorie de Galois imaginaire” [Poi83], Poizat introduced the idea that the classification of certain abstract constructions of model theory — namely interpretable sets or Shelah’s imaginaries — could play an important role in our comprehension of specific structures. The classification of definable sets, in the guise of quantifier elimination results, has historically been used as a central ingredient in many applications of model theory. But the development of more sophisticated model theoretic

Date: April 25, 2023.

1991 *Mathematics Subject Classification.* Primary: 03C45; Secondary: 03C10, 03C60, 12J10, 12L12.

Key words and phrases. Model Theory, Valued Fields, Classification of Imaginaries, Non-standard Frobenius Automorphism, Separated pairs, Linear structures.

MH was partially supported by the German Research Foundation (DFG) via CRC 878, HI 2004/1-1 (part of the French-German ANR-DFG project GeoMod) and under Germany’s Excellence Strategy EXC 2044-390685587, ‘Mathematics Münster: Dynamics-Geometry-Structure’.

SRK was partially supported by GeoMod AAPG2019 (ANR-DFG), Geometric and Combinatorial Configurations in Model Theory.

Both authors were partially supported by ValCoMo (ANR-13-BS01-0006). A substantial part of this work was carried out when both authors participated in the thematic trimester “Model theory, combinatorics and valued fields” at the “Institut Henri Poincaré” in Paris. We are grateful to the IHP and its staff for hosting this event and providing us with this opportunity.

tools, in particular stability theory, naturally took place in the larger category of quotients of definable sets by definable equivalence relations, *i.e.*, interpretable sets. Shelah concretised this idea with his eq construction that formally makes every interpretable set definable.

However, these interpretable sets immediately escape the realm of well understood and classified objects, complicating the possibility of applying new tools from stability theory in specific examples, in particular coming from algebra. Poizat's idea was that these interpretable sets should also be classified, and he did so in algebraically closed fields and in differentially closed fields. In both cases, he showed that they are all definably isomorphic to definable sets, *i.e.*, the categories of definable and interpretable sets are equivalent — we say that these structures *eliminate imaginaries*. This property later became an essential feature in model-theoretic applications, *e.g.*, to diophantine geometry and algebraic dynamics.

The question of elimination of imaginaries also has a very geometric flavour: given a definable family of sets $X \subseteq Y \times Z$, one wishes to find a definable function $f : Z \rightarrow W$ such that for all $z_1, z_2 \in Z$, $X_{z_1} := \{y \in Y : (y, z_1) \in X\} = X_{z_2}$ if and only if $f(z_1) = f(z_2)$ — in other words, one wishes to find a canonical parametrisation of this family where each set appears exactly once. We refer the reader to Section 2.1 and [TZ12, Section 8.4] for further details on these notions and constructions.

Elimination of imaginaries results were then established for numerous structures, but it was not until work of Haskell, Hrushovski and Macpherson [HHM06] that the first complete classification of interpretable sets in a valued field was proved. However, in this case the situation is more complex. Indeed, the field itself does not eliminate imaginaries, as both the value group and the residue field are interpretable but not isomorphic to a definable set. Nevertheless, one can add certain well understood interpretable sets, the *geometric sorts*. These sorts consist of the field \mathbf{K} and, for all $n \in \mathbb{Z}_{>0}$, of the space $\mathbf{S}_n := \mathrm{GL}_n(\mathbf{K})/\mathrm{GL}_n(\mathcal{O})$ of free rank n \mathcal{O} -submodules of \mathbf{K}^n , where \mathcal{O} denotes the valuation ring, and of the space $\mathbf{T}_n := \bigcup_{s \in \mathbf{S}_n} s/\mathfrak{m}_s$ where $\mathfrak{m} \subseteq \mathcal{O}$ is the unique maximal ideal — *cf.* Section 2.2 for a precise definition of the geometric language. The main result of [HHM06] states that the theory ACVF of algebraically closed non-trivially valued fields eliminates imaginaries in the geometric sorts: given a definable family of sets $X \subseteq Y \times Z$, there exists a definable function $f : Z \rightarrow W$, with W a product of geometric sorts, such that for all $z_1, z_2 \in Z$, $X_{z_1} = X_{z_2}$ if and only if $f(z_1) = f(z_2)$ — equivalently the category of sets interpretable in an algebraically closed valued field is equivalent to the category of sets definable in its geometric sorts. One cannot overstate the impact of this result, as it opened the way for the development of geometric model theory in the context of valued fields. A beautiful illustration of the power of these new methods is the work by Hrushovski and Loeser on topological tameness in non-archimedean geometry [HL16].

In this paper we consider imaginaries in more general classes of henselian valued fields of characteristic 0, and also in certain valued difference fields, *i.e.*, valued fields endowed with a distinguished automorphism compatible with the valuation.

In the last 25 years, the model theory of existentially closed difference fields, largely developed by Chatzidakis and Hrushovski (see [CH99]), has led to several spectacular applications — among others in algebraic dynamics. Note that the corresponding theory ACFA does eliminate imaginaries and this fact plays an essential role in later developments. A very deep result of Hrushovski [Hru04], which takes the form of a Frobenius-twisted version of the Lang-Weil estimates, implies that ACFA is in fact the asymptotic theory of Frobenius automorphisms ϕ_q : any non-principal ultraproduct of (\mathbb{F}_q^a, ϕ_q) is a model of ACFA. Key properties of algebraic difference varieties may thus be read off from specializations to the Frobenius automorphisms.

It is also natural to consider the non-standard Frobenius acting on an algebraically closed valued field, *i.e.*, the limit theory of the valued difference fields $(\mathbb{F}_p(t)^a, v_t, \phi_p)$, as the prime p grows, where v_t is an extension of the t -adic valuation. By results of Hrushovski [Hru02] and Durhan [Azg10], this limit theory corresponds to the theory of existentially closed valued difference fields of equicharacteristic zero with ω -increasing automorphism — *cf.* Section 2.4 for a detailed discussion. In fact, these structures naturally arise, as early as in Hrushovski's proof of the twisted Lang-Weil estimates, in the study of algebraic difference varieties, by way of transformal specializations. One may thus expect that the development of a geometric model theory of valued difference fields will turn out useful in the future in geometric applications — as it did in the case of ACVF.

Main results. The classification of imaginaries in ACVF by the geometric sorts was later extended to other valued fields: real closed valued fields [Mel06], separably closed valued fields of finite imperfection degree [HKR18], and p -adic fields and their ultraproducts [HMR18] — which allowed to uniformise and extend Denef's result on the rationality of certain zeta functions to interpretable sets. The question remained whether a general principle underlined all these results. Such a principle was conjectured in the early 2000's by Hrushovski. The present paper establishes it for a large class of henselian fields, which covers most of the examples considered in applications, and extends it to valued fields with operators.

At this level of generality, one cannot expect elimination in the geometric sorts. Indeed, the residue field and the value group can be arbitrary and might not themselves eliminate imaginaries as is the case in all the results cited above. However, a fundamental idea of the model theory of valued fields, the so-called Ax-Kochen-Ershov principle, is that the model theory of a henselian equicharacteristic zero field should be controlled by its value group and residue field. This principle takes its name from the result of Ax and Kochen [AK65] and independently Ershov [Ers65] that this is indeed the

case for elementary equivalence, but this phenomenon has also been observed with respect to numerous other aspects of valued fields, from model theoretic tameness (starting with [Del81]) to motivic integration [HK06].

It is thus tempting to conjecture that, beyond the geometric sorts, imaginaries in equicharacteristic zero henselian fields only arise from the value group and the residue field. However, non trivial torsors of the residue field give rise to serious obstructions to this conjecture. One is thus naturally led to define the \mathbf{k} -linear imaginaries. Consider the two sorted language \mathcal{L}_{mod} of \mathbf{A} -modules \mathbf{V} with the ring structure on \mathbf{A} , the group structure on \mathbf{V} and scalar multiplication. Given an interpretable set X (more precisely a definable quotient of the vector space sort \mathbf{V}) in the \mathcal{L}_{mod} -theory of dimension n vector spaces over a field and given some \mathcal{O} -lattice $s \in \mathbf{S}_n$, we can consider the interpretation $X^{(\mathbf{k}, s/\mathfrak{m}s)}$ of X in the structure $(\mathbf{k}, s/\mathfrak{m}s)$. We then define:

$$\mathbf{T}_{n,X} := \bigsqcup_{s \in \mathbf{S}_n} X^{(\mathbf{k}, s/\mathfrak{m}s)}.$$

Note that if $X = \mathbf{V}$, $\mathbf{T}_{n,X} = \mathbf{T}_n$ and if X is the one element quotient of \mathbf{V} , $\mathbf{T}_{n,X} \cong \mathbf{S}_n$. In general, the $\mathbf{T}_{n,X}$ are essentially those interpretable sets that admit definable surjections $\mathbf{T}_n \rightarrow \mathbf{T}_{n,X} \rightarrow \mathbf{S}_n$. We write $\mathbf{k}^{\text{leq}} := \bigsqcup_{n,X} \mathbf{T}_{n,X}$. Before we state our main results, let us address some technical points. The first is that we prove a result not only in in equicharacteristic 0, but also in finitely ramified mixed characteristic. In the latter case one needs to also consider the higher residue rings $\mathbf{R}_\ell := \mathcal{O}/\ell\mathfrak{m}$, for $\ell \in \mathbb{Z}_{>0}$, where $\ell\mathfrak{m} := \{\ell \cdot x : x \in \mathfrak{m}\}$. These rings often play a crucial role in this situation, and they also come with their linear imaginaries and hence we define $\mathbf{T}_{n,\ell,X} := \bigsqcup_{s \in \mathbf{S}_n} X^{(\mathbf{R}_\ell, s/\ell\mathfrak{m}s)}$ and $\mathbf{R}^{\text{leq}} := \bigsqcup_{n,\ell,X} \mathbf{T}_{n,\ell,X}$, where X is now interpretable in the \mathcal{L}_{mod} -theory of free rank n modules. Moreover, if the residue field \mathbf{k} comes with additional structure, in the definition of \mathbf{k}^{leq} and \mathbf{R}^{leq} , we need to consider all X interpretable in the corresponding enrichment of the theory of free rank n modules.

The second point is that eliminating imaginaries often splits in two distinct problems: describing quotients under the action of finite symmetric groups (in other words finding canonical parameters for finite sets) and classifying interpretable sets up to one-to-finite correspondences: given a definable family of sets $X \subseteq Y \times Z$, one wishes to find a one-to-finite definable correspondence $F : Z \rightarrow W$ such that for all $z_1, z_2 \in Z$, $X_{z_1} = X_{z_2}$ if and only if $F(z_1) = F(z_2)$. This latter property is usually referred to as weak elimination of imaginaries and will be the main focus of this paper.

Our first main result is the following Ax-Kochen-Ershov principle for weak elimination of imaginaries, where $\mathbf{\Gamma}^{\text{eq}}$ refers to the collection all sets interpretable in the (enriched) ordered abelian group $\mathbf{\Gamma}$:

Theorem A (Theorem 6.1.1). *Let (K, v) be a characteristic zero henselian valued field, possibly with angular components and added structure on the value group $\mathbf{\Gamma}$ and, separately, the residue field \mathbf{k} . Assume that:*

- (\mathbf{C}_Γ) *The induced structure on Γ is definably complete;*
- (\mathbf{FR}) *For every $\ell \in \mathbb{Z}_{>0}$, the interval $[0, v(\ell)]$ is finite and \mathbf{k} is perfect;*
- ($\mathbf{I}_\mathbf{k}$) *The residue field \mathbf{k} is infinite;*
- ($\mathbf{E}_\mathbf{k}^\infty$) *The induced theory on \mathbf{k} eliminates \exists^∞ .*

Then K weakly eliminates imaginaries in the sorts $\mathbf{K} \cup \Gamma^{\text{eq}} \cup \mathbf{R}^{\text{leq}}$.

All the results with angular component, even in the algebraically closed (equicharacteristic zero) case, are new. Angular components are (compatible) multiplicative morphisms $\text{ac}_n : \mathbf{K}^\times \rightarrow \mathbf{R}_n^\times$ extending the residue map on \mathcal{O}^\times . They play a key role in the development of the model theoretic study of valued fields, in particular in the Cluckers-Loeser treatment of motivic integration [CL08], by providing uniform cell decomposition results.

Definable completeness of the value group is a necessary hypothesis for the conclusion to hold, since otherwise additional definable cuts appear, inducing more definable \mathcal{O} -submodules and hence more complex imaginaries. It is worth noting that $\text{PRES} = \text{Th}(\mathbb{Z})$ and $\text{DOAG} = \text{Th}(\mathbb{Q})$ are the only complete theories of pure ordered abelian groups which are definably complete. As both PRES and DOAG eliminate imaginaries, we thus get $\Gamma = \Gamma^{\text{eq}}$ under the assumptions of Theorem A in case Γ is not enriched.

As a corollary, Theorem A yields that if F is a field of characteristic 0 which eliminates \exists^∞ , the theories of the valued fields $F((t))$ and $F((t^\mathbb{Q}))$ (with or without angular components) weakly eliminate imaginaries in the sorts $(\mathbf{K} \cup \mathbf{k}^{\text{leq}})$ — noting that $\Gamma \cong \mathbf{S}_1$ may be identified with a sort in \mathbf{k}^{leq} . In the particular case that F is (of characteristic 0 and) algebraically closed, real closed or pseudofinite, using results of Hrushovski on linear imaginaries, we deduce that $F((t))$ and $F((t^\mathbb{Q}))$ (with or without angular components) eliminate imaginaries in the geometric sorts, after naming some constants in the pseudofinite and in the real closed case (see Corollary 6.1.7 for the precise statement), thus obtaining an absolute elimination result in these cases.

Without angular components, this provides alternate proofs of Mellor's result [Mel06] for $\mathbb{R}((t^\mathbb{Q}))$ and Hrushovski-Martin-Rideau's result [HMR18] for $F((t))$ where F is pseudofinite of characteristic 0. Independent work of Vicaria [Vic21] also yields the case of $\mathbb{C}((t))$, although her work also applies to more general value groups.

In mixed characteristic, the main example covered by Theorem A is $W(\mathbb{F}_p^a)$, the fraction field of the ring of Witt vectors with coefficients in \mathbb{F}_p^a , and more generally finite extensions of $W(F)$ for any perfect infinite field F of characteristic p which eliminates \exists^∞ . Thus, Theorem A provides an important step towards proving that the imaginaries of $W(\mathbb{F}_p^a)$ are classified by the geometric sorts as well.

The second main result of this paper concerns valued difference fields. Quantifier elimination (and hence an Ax-Kochen-Ershov principle for elementary equivalence) has been proved for various classes. First for isometries in [Sca03; BMS07; AvdD10], then for ω -increasing automorphisms — for every

$x \in \mathfrak{m}$, $v(\sigma(x)) > \mathbb{Z} \cdot v(x)$ — in [Azg10; Hru02]. Both of these contexts were subsumed in later work of Kushik [Pal12] on *multiplicative* automorphisms where the automorphism acts as multiplication by some element of an ordered field (*cf.* Definition 2.4.5 for precise definitions). Finally Durhan and Onay [DO15] proved that these results hold without any hypothesis on the automorphism.

Our second result focuses on the multiplicative setting where we prove an absolute elimination of imaginaries result for the respective model-companions:

Theorem B (Theorem 6.2.1). *The theory $\text{VFA}_{0,0}^{\text{mult}}$ eliminates imaginaries in the geometric sorts.*

Since the isometric case and the ω -increasing case correspond, respectively, to the asymptotic theory of \mathbb{C}_p with an isometric lifting of the Frobenius and to $\mathbb{F}_p(t)^a$ with the Frobenius, an immediate corollary of these results is a uniform elimination of imaginaries for large p in these structures. By elimination of imaginaries in ACVF, the result is even uniform for all p in the latter case.

Overview of the paper. The proofs of both theorems follow the same general strategy and many technical results are shared between the two. The proof consists of three largely independent steps (Sections 3 to 5).

In stable theories, every type p — a maximal consistent set of definable sets — is definable, *i.e.*, for every definable $X \subseteq Y \times Z$, the set $\{z \in Z : X_z \in p\}$ is definable. This was used in many proofs of weak elimination of imaginaries in the stable context to reduce the problem of finding canonical parameters for definable sets to finding canonical parameters for types; which, counter-intuitively maybe, is a simpler problem. In [Hru14], Hrushovski formalised the idea that even in an unstable context, this reduction could also prove useful, provided definable types were dense: over any algebraically closed imaginary set of parameters, any definable set contains a definable type.

The first step of the proof consists in proving such density results. But the above statement cannot hold in the full generality of henselian equicharacteristic zero valued fields since it might already fail in the residue field. We prove however that, under certain hypotheses, *quantifier free* definable types are dense, *cf.* Theorem 3.1.1. This result does not apply to discrete valued fields since the family of intervals contains arbitrarily large finite sets. In Theorem 3.1.3, we do however prove that the density of quantifier free invariant types holds in this context.

The proof improves on similar results in [Rid19] and the general idea is the same. In arity one, we look for a minimal finite set of balls covering the given definable set. The general case proceeds by fibration in relative arity one and by considering germs of functions into the space of (finite sets of) balls instead of actual balls. This fibration process is where most of the technical assumptions of Theorem A are used, in particular the elimination of \exists^∞ in the residue field.

In contexts with an absolute elimination of quantifiers (as, *e.g.*, in [Rid19; HKR18]) this first density result (and the implicit computation of canonical bases) suffices to conclude that weak elimination of imaginaries holds. In equicharacteristic zero henselian (and σ -henselian) fields, types come with more information than quantifier free types; an information that mostly lives in the short exact sequence $1 \rightarrow \mathbf{k}^\times \rightarrow \mathbf{RV}^\times := \mathbf{K}^\times/(1 + \mathfrak{m}) \rightarrow \mathbf{\Gamma}^\times \rightarrow 0$. The second step of our proof, Theorem 4.1.1, consists in showing that quantifier free invariant types have invariant completions over \mathbf{RV} (and \mathbf{k} -vector spaces) — this generalises to mixed characteristic by considering the higher residue rings.

By quantifier elimination relative to \mathbf{RV} , this step reduces to, given an invariant type, computing canonical generators of the structure generated by (realisations of) the type in \mathbf{RV} . Note that in the conclusion of Theorem 4.1.1 the types considered are invariant over definable sets which are of the same size as the model. We do however show various folklore results implying that this a well behaved notion when these sets are stably embedded.

The third step consists in studying imaginaries in \mathbf{RV} , which is left as a black box in the previous steps. We show, in the spirit of [HK06, Section 3.3], that the imaginaries in the short exact sequence $1 \rightarrow \mathbf{k}^\times \rightarrow \mathbf{RV}^\times \rightarrow \mathbf{\Gamma}^\times \rightarrow 0$ come essentially from \mathbf{k} and $\mathbf{\Gamma}$. To establish the results we need, as in [HK06, Section 3.3], we consider more generally structures given by (enriched) short exact sequences $0 \rightarrow \mathbf{A} \rightarrow \mathbf{B} \rightarrow \mathbf{C} \rightarrow 0$ of R -modules for some ring R . But our result is in a sense orthogonal to the one of Hrushovski and Kazhdan since we require \mathbf{B} to be a pure (in the sense of model theory) extension of \mathbf{A} and \mathbf{C} , which can be both arbitrarily enriched, whereas [HK06, Lemma 3.21] has strong hypotheses on \mathbf{A} and \mathbf{C} and no hypothesis on \mathbf{B} .

Theorem 5.1.4 is the first version of a series of such reductions of increasing complexity so as to cover the various cases that we require, the ultimate version, Variant 5.2.2 allowing controlled torsion in $\mathbf{\Gamma}$, auxiliary sorts on both the \mathbf{k} and $\mathbf{\Gamma}$ sides and considering not one but a projective system of short exact sequences.

These three steps put together allow us to prove relative results like Theorem A. However, absolute results like Theorem B or Corollary 6.1.7 require one last ingredient: the classification of imaginaries in collections of vector spaces — linear structures in the terminology of [Hru12]. Our contribution consists in a twisted version, *cf.* Lemma 2.5.9.(4), of Hrushovski's result on ACF_0 -linear structures with flags and roots endowed with an automorphism (the final step to prove Theorem B) and a version, Proposition 2.5.21, for real closed fields.

The plan of the paper is as follows. In Section 2, we provide some preliminary results on imaginaries, separated pairs of valued fields (in the sense of Baur), on valued difference fields and on linear structures. Section 3 is devoted to the proof of the two density results for definable (resp. invariant)

types mentioned above. The fact that invariant quantifier free types are invariant over \mathbf{RV} (and \mathbf{R}_n -modules), is established in Section 4. In Section 5, we prove the results about imaginaries in certain (enriched) short exact sequences of modules. Finally, in Section 6, we put everything together and prove our main results, in particular Theorem A and Theorem B.

Acknowledgement. The authors would like to thank the anonymous referee for their numerous comments and advice on an earlier version of this manuscript.

2. PRELIMINARIES

Convention 2.1. Throughout this paper, if M is an \mathcal{L} -structure, X is $\mathcal{L}(M)$ -definable and $A \subseteq M$, then $X(A)$ denotes $X \cap A$.

We adopt this convention, as there are too many structures at play to not be explicit as to which definable (or algebraic) closures we want to consider. For this reason, setting $X(A) := X \cap \text{dcl}(A)$ would lead to ambiguities.

2.1. Imaginaries. Let T be an \mathcal{L} -structure. The language \mathcal{L}^{eq} is the language containing \mathcal{L} with one additional sort S_X for every \mathcal{L} -definable set $X \subseteq Y \times Z$, where Y and Z are product of sorts, and one additional symbol $f_X : Z \rightarrow S_X$. The \mathcal{L}^{eq} -theory T^{eq} is then obtained as the union of T , the fact that the f_X are surjective and that their fibers are the classes of the equivalence relation defined by $X_{z_1} = \{y \in Y : (y, z_1) \in X\} = X_{z_2}$.

Any $M \models T$ has a unique expansion to a model of T^{eq} denoted M^{eq} — whose points are called the *imaginaries*. Throughout this paper, notations with exponent eq — like dcl^{eq} or acl^{eq} — will refer to the \mathcal{L}^{eq} -structure of some ambient M^{eq} .

Given $M \models T$ and an $\mathcal{L}(M)$ -definable set X , we denote by $\ulcorner X \urcorner \subseteq M^{\text{eq}}$ the intersection of all $A = \text{dcl}^{\text{eq}}(A) \subseteq M^{\text{eq}}$ such that X is $\mathcal{L}^{\text{eq}}(A)$ -definable. By construction of T^{eq} , X is $\mathcal{L}(\ulcorner X \urcorner)^{\text{eq}}$ -definable, so it is the smallest dcl^{eq} -closed set of definition for X . Any dcl^{eq} -generating subset of $\ulcorner X \urcorner$ is called a *code* of X .

If \mathcal{D} is a collection of sorts of \mathcal{L}^{eq} — equivalently, a collection of \mathcal{L} -interpretable sets — we say that X is *coded* in \mathcal{D} if it is $\mathcal{L}^{\text{eq}}(\mathcal{D}(\ulcorner X \urcorner))$ -definable — *i.e.*, it admits a code in \mathcal{D} . The theory T is said to *eliminate imaginaries* in \mathcal{D} if, for every $M \models T$, every $\mathcal{L}(M)$ -definable set X is coded in \mathcal{D} — equivalently, for every $e \in M^{\text{eq}}$, there is some $d \in \mathcal{D}(\text{dcl}^{\text{eq}}(e))$ such that $e \in \text{dcl}^{\text{eq}}(d)$. By compactness, this is equivalent to the definition in the introduction, provided $\text{dcl}(\emptyset)$ contains two elements. Finally, we say that the theory T *weakly eliminates imaginaries* in \mathcal{D} if for every $e \in M^{\text{eq}}$, there is some $d \in \mathcal{D}(\text{acl}^{\text{eq}}(e))$ such that $e \in \text{dcl}^{\text{eq}}(d)$.

We refer the reader to [TZ12, Section 8.4] for a detailed exposition of these notions.

2.2. The languages of valued fields. Any valued field (K, v) can be considered as a structure in the language \mathcal{L}_{div} with one sort \mathbf{K} for the valued field, the ring language and a binary relation $x|y$ interpreted as $v(x) \leq v(y)$. Note that in every language of valued fields that we will consider there is a sort \mathbf{K} for the valued field and hence, whenever M is a structure representing a valued field, $\mathbf{K}(M)$ will denote the underlying valued field.

The language \mathcal{L}_{div} owes its widespread use to the following result, essentially due to Robinson [Rob77]:

Fact 2.2.1. *The \mathcal{L}_{div} -theory ACVF of algebraically closed non trivially valued fields eliminates quantifiers.*

Notation. We will write $\mathcal{L}_0 := \mathcal{L}_{\text{div}}$ and throughout this paper, notations with an index 0 — like dcl_0 , acl_0 or tp_0 — will refer to the quantifier free \mathcal{L}_0 -structure; or equivalently the structure induced by any model of ACVF containing the valued field under consideration.

Given a valued field, seen as an \mathcal{L}_0 -structure, we will denote by $\mathcal{O} := \{x \in \mathbf{K} : v(x) \geq 0\}$ its valuation ring, $\mathfrak{m} := \{x \in \mathbf{K} : v(x) > 0\}$ its maximal ideal, $\mathbf{k} := \mathcal{O}/\mathfrak{m}$ its residue field and $\Gamma := \mathbf{K}/\mathcal{O}^\times$ its value *monoid*; it is the union of the value group $\Gamma^\times = \mathbf{K}^\times/\mathcal{O}^\times$ and the class of 0 that we usually denote ∞ . We also denote $\text{res} : \mathcal{O} \rightarrow \mathbf{k}$ and $v : \mathbf{K} \rightarrow \Gamma$ the canonical projections. More generally, for every $n \in \mathbb{Z}_{>0}$, we write $\mathbf{R}_n := \mathcal{O}/n\mathfrak{m}$. Let also $\text{res}_n : \mathcal{O} \rightarrow \mathbf{R}_n$ be the canonical projection, \mathbf{R}_∞ the (pro-definable) set $\varprojlim_n \mathbf{R}_n$ and $\text{res}_\infty : \mathcal{O} \rightarrow \mathbf{R}_\infty$ the natural map. Note that, working in a sufficiently saturated model, $\mathbf{R}_\infty \cong \mathcal{O}/\mathfrak{m}_\infty$, where $\mathfrak{m}_\infty := \{x \in \mathbf{K} : v_\infty(x) > \Delta_\infty\}$ and $\Delta_\infty \leq \Gamma$ is the convex subgroup generated by $v(\text{char}(\mathbf{k}))$, in mixed characteristic, and $\Delta_\infty = 0$, otherwise. It is a valuation ring whose fraction field is naturally identified with the residue field \mathbf{k}_∞ associated to the (equicharacteristic) valuation $v_\infty : \mathbf{K} \rightarrow \Gamma \rightarrow \Gamma/\Delta_\infty$. We also define $\mathbf{R} := \bigsqcup_{n>0} \mathbf{R}_n$.

Although most of the present paper is rather insensitive to the choice of language for valued fields — or, rather, we work in $\mathcal{L}_0^{\text{eq}}$ — we will at times need to work in certain languages tailored for specific elimination results. The first of them is the Haskell-Hrushovski-Macpherson geometric language. For every $n \in \mathbb{Z}_{>0}$, let $\mathbf{S}_n \cong \text{GL}_n(\mathbf{K})/\text{GL}_n(\mathcal{O})$ be the (interpretable) set of rank n free sub- \mathcal{O} -modules of \mathbf{K}^n , and $\mathbf{T}_n := \bigcup_{s \in \mathbf{S}_n} s/\mathfrak{m}s$. Let $\mathbf{S} := \bigcup_n \mathbf{S}_n$, $\mathbf{T} := \bigcup_n \mathbf{T}_n$ and $\mathcal{G} := \mathbf{K} \cup \mathbf{S} \cup \mathbf{T}$. We also denote by $s_n : \text{GL}_n(\mathbf{K}) \rightarrow \mathbf{S}_n$, $t_n : \text{GL}_n(\mathbf{K}) \rightarrow \mathbf{T}_n$ and $\tau_n : \mathbf{T}_n \rightarrow \mathbf{S}_n$ the canonical projections. Note that $\text{GL}_n(K)$ naturally acts transitively on \mathbf{T}_n and on \mathbf{S}_n , and that the map τ_n is compatible with these actions.

These interpretable sets (and the so called *geometric language* of which they are the sorts, which also contains the maps s_n , t_n and τ_n) were introduced to classify imaginaries in ACVF:

Fact 2.2.2 ([HHM06, Theorem 1.0.1]). *The theory ACVF eliminates imaginaries in the geometric sorts \mathcal{G} .*

The second language that we will use allows for a description of definable sets in certain henselian fields. The exact language that we use was introduced by Flenner in [Fle11]. For every $n \in \mathbb{Z}_{>0}$, let \mathbf{RV}_n be the multiplicative monoid $\mathbf{K}/(1 + n\mathfrak{m})$; it is the union of the group $\mathbf{RV}_n^* = \mathbf{K}^\times/(1 + n\mathfrak{m})$ and the class of 0, also denoted 0. Let $rv_n : \mathbf{K} \rightarrow \mathbf{RV}_n$ and $rv_{n,m} : \mathbf{RV}_n \rightarrow \mathbf{RV}_m$ denote the canonical projections. The valuation induces a map $\mathbf{RV}_n \rightarrow \Gamma$ that we also denote v . This map induces a short exact sequence:

$$1 \rightarrow \mathbf{R}_n^\times \rightarrow \mathbf{RV}_n^\times \rightarrow \Gamma^\times \rightarrow 0.$$

Remark 2.2.3. If $v(n) = v(m)$, then $rv_{n,m} : \mathbf{RV}_n \cong \mathbf{RV}_m$. In particular, in equicharacteristic zero all \mathbf{RV}_n are canonically isomorphic to \mathbf{RV}_1 . We allow this redundancy in order to have a uniform treatment of characteristic zero henselian valued fields.

Moreover, in positive characteristic p , if n is prime to p , $\mathbf{RV}_n \cong \mathbf{RV}_1$ and otherwise $\mathbf{RV}_n \cong \mathbf{K}$. In that case, it makes more sense to only consider \mathbf{RV}_1 — see below.

Moreover, \mathbf{RV}_n is endowed with the trace of addition which we denote, in Krasner's hyperfield manner: $\zeta \oplus \xi := \{rv_n(x + y) : rv_n(x) = \zeta \text{ and } rv_n(y) = \xi\} \subseteq \mathbf{RV}_n$. We say that $\zeta \oplus \xi$ is well-defined when $\zeta \oplus \xi = \{\chi\}$ is a singleton, and we often write $\zeta \oplus \xi = \chi$ in that case.

Note that for any two disjoint balls b_1 and b_2 , in some valued field (K, v) , and any $a_i, c_i \in b_i$, $rv_1(a_1 - a_2) = rv_1(c_1 - c_2)$. We will denote by $rv_1(b_1 - b_2)$ this common value. If $b_1 \cap b_2 \neq \emptyset$, by convention, $rv_1(b_1 - b_2) = 0$.

We denote by \mathbf{RV}_∞ the (pro-definable) set $\lim_{\leftarrow n} \mathbf{RV}_n$, and $rv_\infty : \mathbf{K} \rightarrow \mathbf{RV}_\infty$ denotes the natural map. Note that $\mathbf{RV}_\infty \cong \mathbf{K}/(1 + \mathfrak{m}_\infty)$, as pro-definable sets. We also denote $\mathbf{RV} := \bigsqcup_n \mathbf{RV}_n$.

Let $\mathcal{L}_{\mathbf{RV}}$ be the language with sorts \mathbf{K} , Γ and \mathbf{RV}_n , for all $n \in \mathbb{Z}_{>0}$, the ring structure on \mathbf{K} , ordered (abelian) monoid structure with a constant for ∞ on Γ , multiplication, constants 0, 1 and a ternary predicate \oplus on each \mathbf{RV}_n , the valuation map $v : \mathbf{K} \rightarrow \Gamma$ and the maps $rv_n : \mathbf{K} \rightarrow \mathbf{RV}_n$ and $rv_{n,m} : \mathbf{RV}_n \rightarrow \mathbf{RV}_m$. Let $\mathcal{L}_{\mathbf{RV}_1}$ be its restriction to the sorts \mathbf{K} , Γ and \mathbf{RV}_1 .

Remark 2.2.4. If the interval $[0, v(n)]$ is finite, then the predicate \oplus on \mathbf{RV}_n is definable (in general with parameters, and without parameters in case $v(n) = 0$) using addition on \mathbf{R}_n .

Proof. If $v(\zeta) \leq v(\xi)$, then $\zeta \oplus \xi = \zeta \cdot r_n^{-1}(1 + (\zeta^{-1}\xi))$, where r_n is the map sending $rv_n(x)$ to $\text{res}_n(x)$ whenever $x \in \mathcal{O}$. The remaining cases are dealt with by symmetry. Therefore, it suffices to show that the map r_n is definable. Let $\pi \in \mathbf{RV}_n$ be an element of minimal positive valuation. Then for every $\xi \in \mathbf{RV}_n$, if $v(\xi) = \ell v(\pi) \in [0, v(n)]$, then $r_n(\xi) = r_n(\pi)^\ell \cdot (rv_n(\pi))^{-\ell} \xi$; and if $v(\xi) > v(n)$, $r_n(\xi) = 0$. So r_n is indeed definable (with parameters π and $r_n(\pi)$, unless $v(n) = 0$). \square

Definition 2.2.5. We say that a valued field (K, v) is:

- *algebraically maximal* if it does not admit non trivial immediate algebraic extensions;
- *Kaplansky* if $\Gamma^\times(K)$ is p -divisible and any finite extension of $\mathbf{k}(K)$ has degree prime to p , where $p = \text{char}(\mathbf{k}(K))$ if it is positive and $p = 1$ otherwise;
- *finitely ramified* if for any $\ell \in \mathbb{Z}_{>0}$ the interval $[0, v(\ell)]$ in $\Gamma(K)$ is finite.

Note that a finitely ramified valued field is algebraically maximal if and only if it is henselian, *cf.* [EP05, Theorem 4.1.10].

The following quantifier elimination results are due, respectively, to Basarab [Bas91, Theorem A] in characteristic zero and Delon [Del82, Théorème 3.1] in positive characteristic (see also [Kuh94, Corollary 2.2 and Theorem 2.6]):

- Fact 2.2.6.**
- *Let \mathcal{L} be an \mathbf{RV} -enrichment of $\mathcal{L}_{\mathbf{RV}}$ and T an \mathcal{L} -theory containing the theory Hen_0 of henselian valued fields of characteristic zero. Then T eliminates field quantifiers.*
 - *Let \mathcal{L} be an \mathbf{RV}_1 -enrichment of $\mathcal{L}_{\mathbf{RV}_1}$ and T an \mathcal{L} -theory containing the theory of equicharacteristic p algebraically maximal Kaplansky valued fields, for some fixed $p > 0$. Then T eliminates field quantifiers.*

2.3. Separated pairs of valued fields. In this section, we will gather some results about separated pairs of valued fields, in particular concerning pure stable embeddedness of the residue field and value group pairs in specific contexts. In equicharacteristic zero, most of the results below follow from work of Leloup [Lel90]; and from work of Rioux [Rio17], in unramified mixed characteristic.

Recall that an extension L/K of valued fields is called *separated* if every finite-dimensional K -vector subspace of L admits a K -valuation basis, *i.e.*, a K -basis (b_1, \dots, b_n) which is *valuation independent* over K : for any $a_1, \dots, a_n \in K$ one has $v(\sum a_i b_i) = \min v(a_i b_i)$. Also, for field extensions $K \subseteq L \subseteq U$ and $K \subseteq K' \subseteq U$, we write $L \downarrow_K^{\text{ld}} K'$ if L and K' are linearly disjoint over K .

Definition 2.3.1. Let $K \subseteq L \subseteq U$ and $K \subseteq K' \subseteq U$ be valued field extensions.

- We say L and K' are $\mathbf{\Gamma k}$ -independent over K , denoted by $L \downarrow_K^{\mathbf{\Gamma k}} K'$, if $\mathbf{k}(L) \downarrow_{\mathbf{k}(K)}^{\text{ld}} \mathbf{k}(K')$ and $\Gamma(L) \cap \Gamma(K') = \Gamma(K)$.
- Assume that L/K is separated. Then L is said to be *valuatively disjoint* from K' over K , denoted by $L \downarrow_K^{\text{vd}} K'$, if whenever a tuple (b_1, \dots, b_n) from L is valuation independent over K , it is valuation independent over K' .

Fact 2.3.2. *Let $K \subseteq L \subseteq U$ and $K \subseteq K' \subseteq U$ be valued field extensions, with L/K separated and $L \downarrow_K^{\mathbf{\Gamma k}} K'$. Set $L' := LK'$. Then we have the following:*

- (1) $L \downarrow_K^{\text{vd}} K'$ — in particular, $L \downarrow_K^{\text{ld}} K'$;
- (2) L'/K' is separated;

- (3) $\mathbf{k}(L') = \mathbf{k}(L)\mathbf{k}(K')$ and $\mathbf{\Gamma}(L') = \mathbf{\Gamma}(L) + \mathbf{\Gamma}(K')$;
(4) If $L_1 \subseteq U$ and $f : L \cong L_1$ is an isomorphism over $K \cup \mathbf{k}(L) \cup \mathbf{\Gamma}(L)$, then f extends (uniquely) to an isomorphism $f' : L' \cong L_1 K'$ over K' .

Proof. This is shown by adapting the proof of the corresponding result for K maximally valued from [HHM08, Proposition 12.11]. \square

2.3.1. *Reduction to \mathbf{RV} .* Most of this paper will be concerned with characteristic zero finitely ramified fields, however, for future reference, we will state and prove certain results, mostly regarding pairs, in all characteristics, as the arguments are essentially identical.

Notation. Given a multisorted language \mathcal{L} , we let $\mathcal{L}_{\mathbf{P}}$ be the associated language of pairs, *i.e.*, for every sort \mathbf{S} from \mathcal{L} we add a unary predicate \mathbf{PS} of sort \mathbf{S} to the language. If N is an \mathcal{L} -substructure of M , we will consider the pair of \mathcal{L} -structures $\widetilde{M} = (M, N)$ as an $\mathcal{L}_{\mathbf{P}}$ -structure in the natural way, *i.e.*, $\mathbf{PS}(\widetilde{M}) = \mathbf{S}(N)$ for each sort \mathbf{S} . We denote $\mathbf{P}(\widetilde{M}) = N$ the whole \mathcal{L} -substructure singled out by the \mathbf{PS} 's. Instead of $\widetilde{M} = (M, \mathbf{P}(\widetilde{M}))$, we will often write $(M, \mathbf{P}(M))$. Given a quantifier free \mathcal{L} -definable set X , we extend the above notation and write $\mathbf{P}X$ for the $\mathcal{L}_{\mathbf{P}}$ -definable set whose points in $(M, \mathbf{P}(M))$ are the $\mathbf{P}(M)$ -points of X .

In $\mathcal{L}_{\mathbf{RV}_1, \mathbf{P}}$, the class of separated pairs of valued fields may be axiomatized. For technical reasons, we will consider such pairs in a hybrid language, adding higher \mathbf{RV} sorts for the small valued field. Formally, we let $\mathcal{L}_{\mathbf{RV}, \mathbf{P}}^{\text{hyb}}$ be the language consisting of $\mathcal{L}_{\mathbf{RV}_1, \mathbf{P}}$ together with additional sorts \mathbf{PRV}_n for all $n \geq 2$ and all symbols of $\mathcal{L}_{\mathbf{RV}}$, where for $n \geq 2$ we use \mathbf{PRV}_n instead of \mathbf{RV}_n . *E.g.*, in $\mathcal{L}_{\mathbf{RV}, \mathbf{P}}^{\text{hyb}}$, for $n \geq 2$ we have a function symbol $\text{rv}_n : \mathbf{K} \rightarrow \mathbf{PRV}_n$ and a ternary relation symbol \oplus on \mathbf{PRV}_n .

Let T^* be a theory of separated pairs $(M, \mathbf{P}(M))$ of henselian valued fields in the language $\mathcal{L}_{\mathbf{RV}, \mathbf{P}}^{\text{hyb}}$. Here, M is the $\mathcal{L}_{\mathbf{RV}_1}$ -structure associated to a valued field, $\mathbf{P}(M)$ the $\mathcal{L}_{\mathbf{RV}}$ -structure (interpreted on the respective \mathbf{PS} 's) of the corresponding valued subfield. (For $n \geq 2$, we extend rv_n from $\mathbf{PK}(M)$ to $\mathbf{K}(M)$ trivially, setting $\text{rv}_n(a) := 0 \in \mathbf{PRV}_n(M)$ for any $a \in \mathbf{K}(M) \setminus \mathbf{PK}(M)$.) We assume that M eliminates field quantifiers in $\mathcal{L}_{\mathbf{RV}_1}$ and $\mathbf{P}(M)$ eliminates fields quantifiers in $\mathcal{L}_{\mathbf{RV}}$. Note that we do not assume that \mathbf{PRV}_1 is stably embedded in \mathbf{RV}_1 .

Remark 2.3.3. In positive characteristic p , since $\mathbf{RV}_p \cong \mathbf{K}$, eliminating quantifiers from the sort \mathbf{K} in $\mathcal{L}_{\mathbf{RV}}$ is an empty assumption and it makes more sense to consider pairs of $\mathcal{L}_{\mathbf{RV}_1, \mathbf{P}}$ -structures instead, as in Remark 2.3.13.

By a hybrid \mathbf{RV} -structure, we mean a structure (elementarily equivalent to) $(\mathbf{RV}_1(M), \mathbf{PRV}(M))$, where $M \models T^*$ — with the restriction of the $\mathcal{L}_{\mathbf{RV}, \mathbf{P}}^{\text{hyb}}$ -structure. We also denote \mathbf{RV}^{hyb} the set of sorts $\{\mathbf{RV}_1, \mathbf{\Gamma}\} \cup \{\mathbf{PRV}_n : n \geq 2\}$.

Lemma 2.3.4. *Let $M_0 \preceq N_0$ be hybrid \mathbf{RV} -structures. Then $\mathbf{k}(M_0) \downarrow_{\mathbf{Pk}(M_0)}^{\text{ld}} \mathbf{Pk}(N_0)$ and $\mathbf{\Gamma}(M_0) \cap \mathbf{P}\mathbf{\Gamma}(N_0) = \mathbf{P}\mathbf{\Gamma}(M_0)$.*

Proof. Immediate from the elementarity of the extension. \square

Let M_0 be a hybrid \mathbf{RV} -structure, say (elementarily equivalent to) a structure of the form $(\mathbf{RV}_1(M), \mathbf{PRV}(M))$, where $M \models T^*$. We say that M_0 is finitely ramified if $\mathbf{P}(M)$ is — *i.e.*, $[0, v(\ell)] \cap \mathbf{P}\mathbf{\Gamma}$ is finite for every $\ell \in \mathbb{Z}_{>0}$. In that case, we also assume that $\mathbf{Pk}(M)$ is perfect. In mixed characteristic $(0, p)$, $R_\infty := \varprojlim_n \mathbf{PR}_n(M_0)$ is then a p -ring with perfect residue field $\mathbf{Pk}(M_0)$ — *cf.* [Ser68, Ch. II §5] — and p is not a divisor of zero. So it is a complete mixed characteristic discrete valuation ring and a finite extension of $W(\mathbf{Pk}(M_0))$ of degree $v(p)$, where $W(k)$ denotes the ring of Witt vectors over k . Let π be a uniformizer of R_∞ — *i.e.*, a generator of the maximal ideal — and P its minimal polynomial over $W(\mathbf{Pk}(M_0))$.

Definition 2.3.5. Ramification constants refer to the (infinite) tuple, in $\mathbf{Pk}(M_0)$, of Witt coordinates of the coefficients of a polynomial P as above.

Lemma 2.3.6. *Let M_0 be a finitely ramified hybrid \mathbf{RV} -structure. Assume that $\mathbf{k}^\times(M_0)$ is divisible, or that $\mathbf{P}\mathbf{\Gamma}(M_0)$ is a pure subgroup of $\mathbf{\Gamma}(M_0)$. Then, the following hold:*

- (1) \mathbf{k} and $\mathbf{\Gamma}$ are purely stably embedded and orthogonal.
- (2) The theory of M is determined by the theories of the \mathbf{k} -pair (with a choice of ramification constants), the $\mathbf{\Gamma}$ -pair and ramification data — *i.e.*, the theory stating that $\varprojlim_n \mathbf{PR}_n$ has a uniformizer which is a zero of the polynomial whose Witt coefficients are the ramification constants.

Moreover, the statements (1) and (2) hold in any \mathbf{k} - $\mathbf{\Gamma}$ -enrichment of M_0 , *i.e.*, a \mathbf{k} -enrichment of a $\mathbf{\Gamma}$ -enrichment of M_0 .

Proof. We may assume that M_0 is of the form $(\mathbf{RV}_1(M), \mathbf{PRV}(M))$ for some \aleph_1 -saturated $M \models T^*$. Then, as \aleph_1 -saturated modules are pure-injective, there is a section of the valuation map restricted to the small valued field $\mathbf{P}(M)$, inducing coherent splittings of the sequences

$$1 \rightarrow \mathbf{PR}_n^\times(M_0) \rightarrow \mathbf{PRV}_n^\times(M_0) \rightarrow \mathbf{P}\mathbf{\Gamma}^\times(M_0) \rightarrow 0$$

for all $n \geq 1$. In mixed characteristic, we may assume that the splitting is *normalized*: the chosen uniformizer π is in the image of the splitting, equivalently, $\text{ac}_n(\pi) = 1$ if ac_n is the angular component map induced by the splitting. Indeed, the group Δ generated by $v(\pi)$ is convex, so the quotient is also ordered and hence torsion free. Since $\mathbf{PK}^\times(M)/\mathcal{O}^\times(M) \cdot \pi^{\mathbb{Z}} \cong \mathbf{P}\mathbf{\Gamma}(M_0)/\Delta$, the extension $\mathbf{P}\mathcal{O}^\times(M) \cdot \pi^{\mathbb{Z}} \leq \mathbf{PK}^\times(M)$ is also pure. Pure-injectivity of $\mathbf{P}\mathcal{O}^\times(M)$ (which is an \aleph_1 -saturated abelian group) then allows to extend the retraction $\mathbf{P}\mathcal{O}^\times(M) \cdot \pi^{\mathbb{Z}} \rightarrow \mathbf{P}\mathcal{O}^\times(M)$ sending π to 1 to the whole of $\mathbf{PK}^\times(M)$.

It further follows from the assumptions that the splitting of $1 \rightarrow \mathbf{Pk}^\times(M_0) \rightarrow \mathbf{PRV}_1^\times(M_0) \rightarrow \mathbf{P}\mathbf{\Gamma}^\times(M_0) \rightarrow 0$ extends to a splitting of the sequence $1 \rightarrow$

$\mathbf{k}^\times(M_0) \rightarrow \mathbf{RV}_1^\times(M_0) \rightarrow \mathbf{\Gamma}^\times(M_0) \rightarrow 0$. To see this, note first that if $h : \mathbf{PRV}_1^\times(M_0) \rightarrow \mathbf{PK}^\times(M_0)$ is a retraction of the inclusion map, *i.e.*, $h|_{\mathbf{PK}^\times(M_0)} = \text{id}_{\mathbf{PK}^\times(M_0)}$, h extends (uniquely) to a homomorphism $\tilde{h} : \mathbf{k}^\times(M_0) \cdot \mathbf{PRV}_1^\times(M_0) \rightarrow \mathbf{k}^\times(M_0)$ which is the identity on $\mathbf{k}^\times(M_0)$, since $\mathbf{k}^\times(M_0) \cap \mathbf{PRV}_1^\times(M_0) = \mathbf{PK}^\times(M_0)$. It is enough to show that \tilde{h} may be extended to a homomorphism $h' : \mathbf{RV}_1^\times(M_0) \rightarrow \mathbf{k}^\times(M_0)$. In case $\mathbf{k}^\times(M_0)$ is divisible, this is clear, since divisible abelian groups are injective. In case $\mathbf{PT}(M_0)$ is a pure subgroup of $\mathbf{\Gamma}(M_0)$, we conclude as above.

Note that the additional structure on \mathbf{RV}^{hyb} , beyond the abelian structure, is given by \oplus and some $\mathbf{k}\text{-}\mathbf{\Gamma}$ -enrichment. As explained in Remark 2.2.4, \oplus can be defined using the ring structure on \mathbf{k} and the \mathbf{PR}_n (using the splitting, no further constants are required. Moreover, \mathbf{PR}_n is a finite extension, generated by the zero of a polynomial with coefficients the ramification constants, and, as such, is \emptyset -interpretable in $W_n(\mathbf{PK})$, which is itself \emptyset -interpretable in \mathbf{PK} . So, if we add the splittings, \mathbf{RV}^{hyb} is (identified to) a $\mathbf{k}\text{-}\mathbf{\Gamma}$ -enrichment of the product of \mathbf{k} and $\mathbf{\Gamma}$. In the product structure, (1) and (2) are clear, even for $\mathbf{k}\text{-}\mathbf{\Gamma}$ -enrichments. The result follows, as (1) and (2) are preserved in any reduct of the product structure that carries the whole structure on \mathbf{k} and $\mathbf{\Gamma}$. \square

We now get back to the $\mathcal{L}_{\mathbf{RV}, \mathbf{P}}^{\text{hyb}}$ -theory T^* of separated pairs of valued fields. Let $M, N \models T^*$, where we suppose that N is $|M|^+$ -saturated, and let $A \leq M$ and $f : A \rightarrow N$ be some embedding.

Definition 2.3.7. We say that:

- (1) A is *good* if $\mathbf{PK}(A) \leq \mathbf{K}(A)$ is a separated extension of valued fields with

$$\mathbf{K}(A) \downarrow_{\mathbf{PK}(A)}^{\mathbf{\Gamma}_{\mathbf{k}}} \mathbf{PK}(M),$$

- (2) f is *good* if $A \leq M$ and $f(A) \leq N$ are good and $f_{\mathbf{RV}^{\text{hyb}}}$ is elementary — for the $\mathcal{L}_{\mathbf{RV}, \mathbf{P}}^{\text{hyb}}|_{\mathbf{RV}^{\text{hyb}}}$ -structure.

Proposition 2.3.8. *Assume f is a good embedding. Then f extends to a good embedding $g : M \rightarrow N$.*

Proof. We proceed step by step.

Step 1. *We may extend f to a good map defined on $A \cup \mathbf{RV}^{\text{hyb}}(M)$ — and thus assume that $\mathbf{RV}^{\text{hyb}}(A) = \mathbf{RV}^{\text{hyb}}(M)$.*

Indeed, this follows from saturation, the fact that $f_{\mathbf{RV}^{\text{hyb}}}$ is elementary and that the only symbols in the language involving both \mathbf{K} and \mathbf{RV}^{hyb} are maps from \mathbf{K} to \mathbf{RV}^{hyb} .

Step 2. *We may extend f to a good map defined on (the substructure generated by) $A \cup \mathbf{P}(M)$ — and thus assume that $\mathbf{PK}(A) = \mathbf{PK}(M)$.*

Indeed, by \mathbf{K} -quantifier elimination in the $\mathcal{L}_{\mathbf{RV}}$ -theory of the small valued field $\mathbf{P}(M)$, the map $f|_{\mathbf{P}(A)}$ extends to an (elementary) $\mathcal{L}_{\mathbf{RV}}$ -embedding

$g : \mathbf{P}(M) \rightarrow \mathbf{P}(N)$. As $\mathbf{K}(A) \downarrow_{\mathbf{PK}(A)}^{\Gamma_{\mathbf{K}(A)}} \mathbf{PK}(M)$ and $\mathbf{RV}^{\text{hyb}}(A) = \mathbf{RV}^{\text{hyb}}(M)$, by Fact 2.3.2, $f \cup g$ induces a good embedding of $A \cup \mathbf{P}(M)$ into N .

Step 3. *We may extend f to a good embedding of M into N .*

Indeed, by \mathbf{K} -quantifier elimination in the $\mathcal{L}_{\mathbf{RV}_1}$ -theory of the valued field M , the map f extends to an (elementary) $\mathcal{L}_{\mathbf{RV}_1}$ -embedding $\tilde{f} : M \rightarrow N$. By Lemma 2.3.4, we get $\tilde{f}(\mathbf{K}(M)) \downarrow_{f(\mathbf{PK}(M))}^{\Gamma_{\mathbf{K}(M)}} \mathbf{PK}(N)$, so in particular $\tilde{f}(\mathbf{K}(M)) \downarrow_{f(\mathbf{PK}(M))}^{\text{ld}} \mathbf{PK}(N)$ (and thus $\tilde{f}(\mathbf{K}(M)) \cap \mathbf{PK}(N) = f(\mathbf{PK}(M))$) by Fact 2.3.2(1), showing that \tilde{f} is an $\mathcal{L}_{\mathbf{RV}_1, \mathbf{P}}$ -embedding, with image a good substructure of N . Thus \tilde{f} is a good embedding, since f was already defined on the whole of $\mathbf{RV}^{\text{hyb}}(M)$. \square

Corollary 2.3.9. *The theory T^* is complete relative to \mathbf{RV}^{hyb} , and \mathbf{RV}^{hyb} is purely stably embedded in T^* , i.e., the induced structure is that of a hybrid \mathbf{RV} -structure.*

This even holds resplendently, for any \mathbf{RV}^{hyb} -enrichment of the pair of valued fields.

Proof. Assume that $M, N \models T^*$ are models with $\mathbf{RV}^{\text{hyb}}(M) \cong \mathbf{RV}^{\text{hyb}}(N)$. The isomorphism between the prime substructures, i.e., the substructures of M and N generated by \emptyset , is easily seen to be a good embedding. It follows by Proposition 2.3.8, and a back and forth argument, that it is, in fact, elementary — i.e., $M \equiv N$.

Similarly, if $M \leq N$ — in particular a good substructure — and $f : M \rightarrow N$ is an elementary embedding — in particular a good embedding — inducing the identity on $\mathbf{RV}^{\text{hyb}}(M)$, then it remains a good embedding — and hence an elementary one — when extended by the identity on $\mathbf{RV}^{\text{hyb}}(N)$. Thus, $\text{tp}(M/\mathbf{RV}^{\text{hyb}}(M)) \vdash \text{tp}(M/\mathbf{RV}^{\text{hyb}}(N))$; in other words, \mathbf{RV}^{hyb} is stably embedded. Finally any $\mathcal{L}_{\mathbf{RV}, \mathbf{P}}^{\text{hyb}}|_{\mathbf{RV}^{\text{hyb}}}$ -elementary map on \mathbf{RV}^{hyb} is good and hence $\mathcal{L}_{\mathbf{RV}, \mathbf{P}}^{\text{hyb}}$ -elementary, so \mathbf{RV}^{hyb} is pure.

Resplendence of the result follows formally. (This is folklore. See, e.g., [Con+22, Proposition 2.7] for a proof.) \square

The proof of the corollary, together with the fact that \mathbf{PRV} is purely stably embedded in \mathbf{PK} , yields the following observation. (We will not use this in our paper.)

Remark 2.3.10. If \mathbf{PRV} is (purely) stably embedded in \mathbf{RV}^{hyb} , then \mathbf{PK} is (purely) stably embedded.

Corollary 2.3.11. *Let $M \models T^*$ be such that $\mathbf{P}(M)$ is finitely ramified with perfect residue field. Assume that $\mathbf{k}^\times(M)$ is divisible (which is the case for example if $M \models \text{ACVF}$) or that $\mathbf{PT}(M)$ is a pure subgroup of $\mathbf{T}(M)$. Then in any \mathbf{k} - \mathbf{T} -enrichment of M , the theory of M is determined by ramification data and the theories of \mathbf{k} (with ramification constants) and \mathbf{T} . Moreover \mathbf{k} and \mathbf{T} are purely stably embedded and orthogonal.*

Proof. By Corollary 2.3.9 together with Lemma 2.3.6. \square

Similarly to Remark 2.3.10, we get the following.

Remark 2.3.12. If \mathbf{Pk} is (purely) stably embedded in \mathbf{k} and \mathbf{PT} is (purely) stably embedded in $\mathbf{\Gamma}$, then \mathbf{PK} is (purely) stably embedded.

Remark 2.3.13. If we further assume that $\mathbf{P}(M)$ eliminates field quantifiers in $\mathcal{L}_{\mathbf{RV}_1}$ — *e.g.*, if it is algebraically closed or algebraically maximal Kaplansky of equicharacteristic — then all the above results can easily be adapted to pairs of $\mathcal{L}_{\mathbf{RV}_1}$ -structures (with no need for the rather exotic hybrid \mathbf{RV} -structures).

2.3.2. Characteristic 0 Laurent series fields. Let F be a field of characteristic 0, and let $K := F((t))$. In what follows, we are interested in the pair of valued fields (K^a, K) . Let us first deal with the pair of value groups. Let \mathcal{L}_{og} be the language of ordered groups and DOAG be the theory of non-trivial divisible ordered abelian groups. Let also $\mathcal{L}_{\text{Pres}}$ be the language \mathcal{L}_{og} enriched with a constant 1 and unary predicates for divisibility by integers. Let PRES be the $\mathcal{L}_{\text{Pres}}$ -theory of \mathbb{Z} .

Notation. Let $T_{\mathbb{Q}, \mathbb{Z}}$ be the theory of all structures (Γ, Δ) with $\Gamma \models \text{DOAG}$, $\Delta \models \text{PRES}$ and such for any $\gamma \in \Gamma$ there is a largest $\delta =: \lfloor \gamma \rfloor \in \Delta$ with $\delta \leq \gamma$, considered in the language $\mathcal{L}_{\mathbb{Q}, \mathbb{Z}}$ given by $\mathcal{L}_{\text{og}, \mathbf{P}}$ together with $\mathcal{L}_{\text{Pres}}$ on the predicate \mathbf{P} and the function $\lfloor \cdot \rfloor$.

The quantifier elimination result we state in part (3) of the following lemma has already been obtained by Weispfenning ([Wei99]). (We thank Matthias Aschenbrenner for having brought this to our attention.) We decided to include our proof for convenience of the reader.

Lemma 2.3.14.

- (1) Let $M = (\Gamma, \Delta) \models T_{\mathbb{Q}, \mathbb{Z}}$. Then the map $\gamma \mapsto (\lfloor \gamma \rfloor, \gamma - \lfloor \gamma \rfloor)$ is an \emptyset -definable bijection between Γ and $\Delta \times [0, 1)$, which identifies the \emptyset -definable sets in M with the \emptyset -definable sets in the product structure $(\Delta, 0, +, \leq) \times ([0, 1), 0, \tilde{+}, <)$, where $a \tilde{+} b := a + b - \lfloor a + b \rfloor$ is the group law on $[0, 1)$ induced by the natural bijection between $[0, 1)$ and Γ/Δ .
- (2) In $T_{\mathbb{Q}, \mathbb{Z}}$, the predicate \mathbf{P} is stably embedded with induced structure a pure model of PRES, and $[0, 1)$ is stably embedded, with induced structure given by \mathcal{L}_{og} , so in particular *o-minimal*.
- (3) $T_{\mathbb{Q}, \mathbb{Z}}$ eliminates quantifiers and is complete.

Proof. Let $f : \Gamma \rightarrow \Delta \times [0, 1)$ be the bijection given in (1). Clearly, f is \emptyset -definable, and the product structure $(\Delta, 0, +, <) \times ([0, 1), \tilde{+}, <)$ is \emptyset -definable in M . Conversely, under this identification \mathbf{P} corresponds to $f^{-1}(0)$, $<$ on Γ corresponds to the lexicographic ordering on $\Delta \times [0, 1)$, and the addition on Γ may also be recovered, since if $f(\gamma) = (z, a)$ and $f(\gamma') = (z', a')$, then

$$f(\gamma + \delta) = \begin{cases} (z + z', a \tilde{+} b) & \text{if } a \leq a \tilde{+} b \\ (z + z' + 1, a \tilde{+} b) & \text{otherwise.} \end{cases}$$

This proves (1). Part (2) follows directly from (1).

Let us now show (3). Completeness follows from quantifier elimination, since (\mathbb{Z}, \mathbb{Z}) embeds into every model of $T_{\mathbb{Q}, \mathbb{Z}}$ as a substructure. To prove quantifier elimination, we first note that the \mathcal{L}_{og} -theory of $([0, 1], 0, \tilde{+}, <)$ has quantifier elimination. (This is well known, and we leave the easy proof to the reader.) Moreover, PRES has quantifier elimination in $\mathcal{L}_{\text{Pres}}$. It is thus enough to establish the following claim:

Claim 2.3.15. *If $D \subseteq (\Delta \times [0, 1])^n$ is defined by an atomic formula in the product structure $\Delta \times [0, 1]$, with $\mathcal{L}_{\text{Pres}}$ on Δ and \mathcal{L}_{og} on $[0, 1]$, then $f^{-1}(D) \subseteq M^n$ is defined by a quantifier free formula (without parameters).*

To prove the claim, let us denote the projection onto Δ by π_1 , that onto $[0, 1]$ by π_2 . If φ is of the form $\psi(\pi_1(x_1), \dots, \pi_n(x_1))$ for some atomic $\mathcal{L}_{\text{Pres}}$ -formula $\psi(\bar{y})$, the statement is clear, as then $f^{-1}(D)$ is defined by the quantifier free formula $\psi(\lfloor x_1 \rfloor, \dots, \lfloor x_n \rfloor)$. Else, φ is (equivalent to) a formula of the form $z_1 \pi_2(x_1) \tilde{+} \dots \tilde{+} z_n \pi_2(x_n) = 0$, with $z_1, \dots, z_n \in \mathbb{Z}$, in which case $f^{-1}(D)$ is defined by $\mathbf{P}(\sum_{i=1}^n z_i x_i)$; or φ is (equivalent to) a formula of the form

$$z_1 \pi_2(x_1) \tilde{+} \dots \tilde{+} z_n \pi_2(x_n) < z'_1 \pi_2(x_1) \tilde{+} \dots \tilde{+} z'_n \pi_2(x_n),$$

with $z_1, z'_1, \dots, z_n, z'_n \in \mathbb{Z}$, in which case $f^{-1}(D)$ is defined by the quantifier free formula $\sum_{i=1}^n z_i x_i - \lfloor \sum_{i=1}^n z_i x_i \rfloor < \sum_{i=1}^n z'_i x_i - \lfloor \sum_{i=1}^n z'_i x_i \rfloor$. \square

In fact, the proof of Lemma 2.3.14 yields the following more general result.

Remark 2.3.16. Let $\mathcal{L}^+ \supseteq \mathcal{L}_{\text{Pres}}$ and $T^+ \supseteq \text{PRES}$ be a complete \mathcal{L}^+ -theory with quantifier elimination. Then the corresponding expansion $T_{\mathbb{Q}, \mathbb{Z}}^+$ of $T_{\mathbb{Q}, \mathbb{Z}}$ is complete, eliminates quantifiers, and \mathbf{P} is purely stably embedded with induced structure given by \mathcal{L}^+ .

Since $T_{\mathbb{Q}, \mathbb{Z}}$ admits the complete model (\mathbb{R}, \mathbb{Z}) , it is definably complete. Actually, this even holds resplendently, as shows the following corollary.

Corollary 2.3.17. *Assume that the expansion $T^+ \supseteq \text{PRES}$ is definably complete. Then $T_{\mathbb{Q}, \mathbb{Z}}^+$ is definably complete.*

Proof. Let $(\Gamma, \Delta) \models T_{\mathbb{Q}, \mathbb{Z}}^+$ and let $D \subseteq \Gamma$ be a definable subset which is bounded and non-empty. Then $\lfloor D \rfloor$ is a definable subset of Δ which is non-empty and bounded. By assumption, it admits a supremum s in Δ , which is then the maximum of $\lfloor D \rfloor$ as the order on Δ is discrete.

As the induced structure on $[0, 1]$ is o -minimal by Lemma 2.3.14, the induced structure on $[s, s+1]$ is o -minimal as well, and so $\sup(D) = \sup(D \cap [s, s+1])$ exists in $[s, s+1]$, proving definable completeness. \square

Let us now consider the residue field. By a classical result of Keisler [Kei64], if F and F' are fields such that $F \equiv F'$, then $(F^a, F) \equiv (F'^a, F')$. If $F = F^a$ or F^a is real closed, then the axiomatization, in $\mathcal{L}_{\text{ring}, \mathbf{P}}$, of (F^a, F) is clear, and \mathbf{P} is stably embedded with induced structure that of a ring. In case T_f is a complete theory of fields whose models are neither algebraically

nor real closed, and $F \models T_f$, then the models of the $\mathcal{L}_{ring, \mathbf{P}}$ -theory $T_{f,a}$ of (F^a, F) are precisely the pairs $(M, \mathbf{P}(M))$ of fields such that $M = M^a$ and $\mathbf{P}(M) \models T_f$. By [HKR18, Theorem 4.7], if one definably expands the theory, adding relation symbols \mathbf{ld}_n and function symbols $\ell_{n,i}$, the theory $T_{f,a}$ eliminates quantifiers relative to \mathbf{P} , even resplendently on \mathbf{P} . This yields in particular the following.

Fact 2.3.18. *For $T_f = \text{Th}(F)$ a complete theory of fields (in arbitrary characteristic), the predicate \mathbf{P} is stably embedded in the $\mathcal{L}_{ring, \mathbf{P}}$ -theory $T_{f,a} = \text{Th}(F^a, F)$, with induced structure given by T_f .*

This even holds resplendently, for any $\mathcal{L} \supseteq \mathcal{L}_{ring}$ and any \mathcal{L} -expansion of F .

Lemma 2.3.19. *Let F be some (enriched) field which eliminates \exists^∞ . Then the pair (F^a, F) also eliminates \exists^∞ .*

Proof. We may suppose that F is neither algebraically nor real closed, as otherwise the result is clear. By the relative quantifier elimination result [HKR18, Theorem 4.7] already mentioned above, if $(M, \mathbf{P}(M)) \equiv (F^a, F)$, and $(M, \mathbf{P}(M)) \preceq (\mathcal{U}, \mathbf{P}(\mathcal{U}))$ then for any $a, b \in \mathcal{U} \setminus (M\mathbf{P}(\mathcal{U}))^a$ we have $tp(a/M) = tp(b/M) =: p_{gen}(x)$, so, assuming that $(\mathcal{U}, \mathbf{P}(\mathcal{U}))$ is sufficiently saturated, any element of \mathcal{U} is the sum of two realizations of p_{gen} . By compactness, for any M -definable set $D = \psi(M) \subseteq M$ we then have

$$(2.1) \quad \begin{aligned} D + D = M &\Leftrightarrow \psi(\mathcal{U}) + \psi(\mathcal{U}) = \mathcal{U} \\ &\Leftrightarrow \psi(\mathcal{U}) \not\subseteq (M\mathbf{P}(\mathcal{U}))^a \Leftrightarrow p_{gen}(x) \vdash \psi(x). \end{aligned}$$

We will now assume that M is \aleph_0 -saturated. Let $\varphi(x, y)$ be a formula with x a single variable. Assume that c is a tuple from M such that $\varphi(M, c)$ is infinite. By compactness, it is enough to find a formula $\chi(y) \in tp(c)$ such that for any c' fro M satisfying χ the set $\varphi(M, c')$ is infinite.

If $p_{gen}(x) \vdash x \in \varphi(x, c)$, we may find such a $\chi(y)$ using Eq. (2.1). So assume now that $p_{gen}(x) \not\vdash x \in \varphi(x, c)$, and choose $a \notin M$ realizing $\varphi(x, c)$. Then $a \in (M\mathbf{P}(\mathcal{U}))^a \setminus M^a$, so in particular $M(a) \not\downarrow_{\mathbf{P}(M)}^{\text{ld}} \mathbf{P}(\mathcal{U})$. Set $A := \text{dcl}(Ma)$. If $\mathbf{P}(A) = \mathbf{P}(M)$, by [HKR18, Lemma 4.1], we have $A \downarrow_{\mathbf{P}(M)}^{\text{ld}} \mathbf{P}(\mathcal{U})$, contradicting that $M(a) \not\downarrow_{\mathbf{P}(M)}^{\text{ld}} \mathbf{P}(\mathcal{U})$. It follows that there is $a' \in \mathbf{P}(A) \setminus \mathbf{P}(M)$, so we find an M -definable function $f : D \rightarrow \mathbf{P}(M)$ with infinite image. Using Fact 2.3.18 and the assumption that the theory of F eliminates \exists^∞ , we may thus find a formula $\chi(y)$ as required, stipulating explicitly the existence of a definable function with infinite image in \mathbf{P} . \square

Fix some characteristic zero field F and let T_{Laur}^* be the theory of separated pairs of valued fields $(M, \mathbf{P}(M))$ with $(\mathbf{k}(M), \mathbf{Pk}(M)) \equiv (F^a, F)$ and $(\mathbf{\Gamma}(M), \mathbf{P}\mathbf{\Gamma}(M)) \models T_{\mathbb{Q}, \mathbb{Z}}$.

Proposition 2.3.20. *The theory T_{Laur}^* is complete, the definable sets \mathbf{k} , $\mathbf{\Gamma}$, \mathbf{RV} , \mathbf{PK} , \mathbf{Pk} , \mathbf{PT} and \mathbf{PRV} are all purely stably embedded, and \mathbf{k} and $\mathbf{\Gamma}$ are orthogonal. All these results hold resplendently for \mathbf{Pk} - \mathbf{PT} -enrichments, and also if one adds an angular component.*

Proof. By Corollary 2.3.9, \mathbf{RV} is stably embedded. Combining Fact 2.3.18 and Lemma 2.3.14 with Corollary 2.3.11, we see that T_{Laur}^* is complete, \mathbf{Pk} , $\mathbf{P}\Gamma$, \mathbf{k} and Γ are stably embedded and the latter are orthogonal. By Remark 2.3.12, \mathbf{PK} is thus stably embedded, and hence so is \mathbf{PRV} . \square

2.3.3. Finitely ramified fields. We will now prove analogous statements in mixed characteristic. Let K be a complete mixed characteristic \mathbb{Z} -valued field with perfect residue field F . We are interested in the pair of valued fields (K^a, K) . Let T_{Witt}^* be a theory of separated pairs of henselian valued fields with fixed ramification data such that $(\mathbf{k}(M), \mathbf{Pk}(M)) \equiv (F^a, F)$ (with ramification constants) and $(\Gamma(M), \mathbf{P}\Gamma(M)) \vDash T_{\mathbb{Q}, \mathbb{Z}}$.

Proposition 2.3.21. *The theory T_{Witt}^* is complete, the definable sets \mathbf{k} , Γ , \mathbf{RV}^{hyb} , \mathbf{PK} , \mathbf{PRV} , \mathbf{Pk} and $\mathbf{P}\Gamma$ are all purely stably embedded, and \mathbf{k} and Γ are orthogonal. All these results hold resplendently for \mathbf{Pk} - $\mathbf{P}\Gamma$ -enrichments, and also if one adds a coherent system of normalized angular components.*

Proof. One may argue as in the proof of Proposition 2.3.20. \square

2.3.4. Divisible value group. Let F be a field. If $\text{char}(F) = p > 0$, assume that F does not admit a finite extension of degree divisible by p (in particular F is perfect). Let $K := F((t^{\mathbb{Q}}))$. In what follows, we are interested in the pair of valued fields (K^a, K) .

Let T_{div}^* be the theory of separated pairs of equicharacteristic algebraically maximal valued fields $(M, \mathbf{P}(M))$ such that $(\mathbf{k}(M), \mathbf{Pk}(M)) \equiv (F^a, F)$ and $\mathbf{P}\Gamma(M) = \Gamma(M) \vDash \text{DOAG}$. Note that the Kaplansky conditions (*cf.* Definition 2.2.5) are satisfied in this case.

Proposition 2.3.22. *The theory T_{div}^* is complete. The definable sets \mathbf{k} , Γ , \mathbf{RV} , \mathbf{PK} , \mathbf{Pk} and \mathbf{PRV} are all purely stably embedded, and \mathbf{k} and Γ are orthogonal. All these results hold resplendently for \mathbf{Pk} - Γ -enrichments, and also if one adds angular components.*

Proof. One may argue as in the proof of Proposition 2.3.20. \square

2.4. Valued difference fields. Let (K, v, σ) be a valued field with an automorphism.

Definition 2.4.1. (1) For every $P \in K[x_0, \dots, x_n]$, $a \in K$, $d \in K^{n+1}$ and $\gamma \in \Gamma(K)^\times$, we say that (P, a, d, γ) is in σ -Hensel configuration if

$$v(P(\nabla a)) > \min_i v(d_i) + \sigma^i(\gamma)$$

and, for all $x, y \in K$ with $v(x - a), v(y - a) > \gamma$,

$$v(P(\nabla y) - P(\nabla x) - d \cdot \nabla(y - x)) > \min_i v(d_i \sigma^i(y - x)).$$

Here, $\nabla a := (\sigma^i(a))_{0 \leq i \leq n}$.

- (2) We say that (K, v, σ) is σ -henselian if for every (P, a, d, γ) in σ -Hensel configuration, there exists $c \in K(M)$ such that $P(\nabla c) = 0$ and

$$v(c - a) \geq \max_{i, d_i \neq 0} v(\sigma^{-i}(P(\nabla a))d_i^{-1}).$$

Fact 2.4.2. *Assume that for every linear non constant $L \in \mathbf{k}(K)[x_0, \dots, x_n]$ and $c \in \mathbf{k}(K)$, there exists $a \in \mathbf{k}(K)$ such that $L(\nabla(a)) = c$ and either:*

- *K is maximally complete;*
- *K is complete (rank one).*

Then (K, v, σ) is σ -henselian.

This follows from Newton approximation, *cf.* [Rid17, Proposition 4.14]. Note that at each step the approximation to the root of a σ -Hensel configuration (P, a, d, γ) improves by at least γ , *cf.* [Rid17, Lemma 4.16], and hence, in rank one, completeness suffices.

Remark 2.4.3. In [Azg10; Pal12; DO15] an *a priori* weaker notion of σ -henselian fields is considered. However, both notions hold in maximally complete fields (with linearly closed residue field) and both allow proving Fact 2.4.4. So any field satisfying either notion is elementarily equivalent to any maximal completion verifying both. It follows that the two notions are equivalent.

Let $\mathcal{L}_{\mathbf{RV}}^\sigma$ be the language $\mathcal{L}_{\mathbf{RV}}$ with three new unary functions $\sigma_{\mathbf{K}} : \mathbf{K} \rightarrow \mathbf{K}$, $\sigma_{\mathbf{RV}} : \mathbf{RV} \rightarrow \mathbf{RV}$ and $\sigma_{\mathbf{\Gamma}} : \mathbf{\Gamma} \rightarrow \mathbf{\Gamma}$. The expected quantifier elimination result also holds in characteristic zero σ -henselian fields, by [DO15, Theorem 7.3] in equicharacteristic 0, and by [Rid17, p. 41, Theorem A] in mixed characteristic:

Fact 2.4.4. *Let \mathcal{L} be an \mathbf{RV} -enrichment of $\mathcal{L}_{\mathbf{RV}}^\sigma$ and T an \mathcal{L} -theory containing the theory Hen_0^σ of characteristic zero σ -henselian valued fields. Then T eliminates field quantifiers.*

Note that it follows from field quantifier elimination that \mathbf{RV} is purely stably embedded. Its induced structure is the expansion of the short exact sequence of $\mathbb{Z}[\sigma]$ -modules $1 \rightarrow \mathbf{k}^\times \rightarrow \mathbf{RV}^\times \rightarrow \mathbf{\Gamma}^\times \rightarrow 0$ by \oplus and $<$ which, by Remark 2.2.4, can be defined (with parameters) in a $\mathbf{\Gamma}$ - \mathbf{k} -enrichment.

In order to obtain model complete theories, one often restricts the behaviour of the automorphism on the value group, *e.g.*, the class of (existentially closed) multiplicative difference valued fields introduced in [Pal12]:

Definition 2.4.5. Let $\text{VFA}_{0,0}^{\text{mult}}$ be the theory of σ -henselian non-trivially valued fields such that:

- (1) For every $P \in \mathbb{Z}[\sigma]$, either $P(\mathbf{\Gamma}_{>0}) = \mathbf{\Gamma}_{>0}$, $P(\mathbf{\Gamma}_{<0}) = \mathbf{\Gamma}_{>0}$ or $P(\mathbf{\Gamma}) = 0$;
- (2) We have $(\mathbf{k}, \sigma_{\mathbf{k}}) \models \text{ACFA}_0$
- (3) The embedding of $\mathbb{Z}[\sigma]$ -modules $\mathbf{k}^\times \rightarrow \mathbf{RV}$ is pure.

Remark 2.4.6. Two multiplicative behaviours of σ are of particular interest.

- (1) The ω -increasing case — *i.e.*, for all $x \in \mathcal{O}$ and $n \in \mathbb{Z}_{>0}$, $v(\sigma(x)) \geq nv(x)$ — studied in [Azg10]. One then gets the asymptotic theory of $(\mathbb{F}_p(t)^a, v_t, \phi_p)$, where ϕ_p is the Frobenius automorphism.
- (2) The isometric case, studied in [BMS07]. In that case, one gets the asymptotic theory of $(\mathbb{C}_p, v_p, \sigma_p)$, where σ_p is an isometric lift of the Frobenius automorphism on $\mathbf{k}(\mathbb{C}_p) = \mathbb{F}_p^a$.

Both characterisations follow (see, *e.g.*, [CH14]) from the Ax-Kochen-Ershov principle for σ -henselian valued fields and Hrushovski's deep result that ACFA_0 is the asymptotic theory of (\mathbb{F}_p^a, ϕ_p) , *cf.* [Hru04].

Fact 2.4.7. *In $\text{VFA}_{0,0}^{\text{mult}}$, \mathbf{k} is a stably embedded pure difference field and Γ is a stably embedded σ -minimal pure ordered $\mathbb{Z}[\sigma]$ -module, and \mathbf{k} and Γ are orthogonal. These results also hold if one adds a σ -equivariant angular component.*

Proof. Condition (3) in Definition 2.4.5 ensures that in any \aleph_1 -saturated model of $\text{VFA}_{0,0}^{\text{mult}}$ there is a σ -equivariant angular component map ac . [Pal12, Theorem 11.8] yields the results if we add such an ac map to the language, and they obviously go down to the reduct without ac . \square

2.5. Linear structures. Let us now recall the results of [Hru12] on linear structures. In the case of valued difference fields, we will need “twisted” versions of these results. As it took us a while to get the arguments clear, we decided to spell them out in detail.

2.5.1. Independent amalgamation. We will first recall some material from [Hru12, Section 4]. We fix a complete stable theory T in some language \mathcal{L} , and we assume that T eliminates quantifiers and imaginaries. Let \mathcal{C}_T be the category consisting of those \mathcal{L} -structures that are algebraically closed substructures of a model of T , with \mathcal{L} -embeddings (which are \mathcal{L} -elementary in T by assumption) as morphisms. Let $\mathbf{n} = \{0, 1, \dots, n-1\}$, and set $\mathcal{P}(\mathbf{n})^- := \mathcal{P}(\mathbf{n}) \setminus \{\mathbf{n}\}$. We consider $\mathcal{P}(\mathbf{n})^-$ and $\mathcal{P}(\mathbf{n})$ as categories, with inclusion maps as morphisms.

Given a functor $A : P \rightarrow \mathcal{C}_T$, where P equals $\mathcal{P}(\mathbf{n})^-$ or $\mathcal{P}(\mathbf{n})$, if $\iota : w_1 \rightarrow w_2$ denotes the inclusion map for two sets $w_1 \subseteq w_2 \in P$, in what follows we will write $A(w_1)$ for the subset $A(\iota)(A(w_1))$ of $A(w_2)$, thus omitting the map $A(\iota)$ in our notation. This slight abuse of notation should not lead to any confusion.

Definition 2.5.1. Let P equal $\mathcal{P}(\mathbf{n})^-$ or $\mathcal{P}(\mathbf{n})$.

- (1) A functor $A : P \rightarrow \mathcal{C}_T$ is called *independence preserving* if for any $w, w' \in P$ with $w \cup w' \in P$ one has $A(w) \downarrow_{A(w \cap w')} A(w')$ (inside $A(w \cup w')$).
- (2) A functor $A : P \rightarrow \mathcal{C}_T$ is called *bounded* if for any $\emptyset \neq w \in P$ one has $A(w) = \text{acl}(\cup_{i \in w} A(\{i\}))$.

- (3) An n -amalgamation problem in T is a bounded independence preserving functor $A^- : \mathcal{P}(\mathbf{n})^- \rightarrow \mathcal{C}_T$. A *solution* of A^- is a bounded independence preserving functor $A : \mathcal{P}(\mathbf{n}) \rightarrow \mathcal{C}_T$ extending A^- .

Definition 2.5.2. The theory T is said to have

- n -existence if every n amalgamation problem in T has a solution;
- n -uniqueness if whenever A and A' are solutions of the same n -amalgamation problem A^- in T , then A and A' are isomorphic over A^- , i.e., there is an \mathcal{L} -isomorphism $f : A(\mathbf{n}) \cong A'(\mathbf{n})$ fixing $A^-(w)$ pointwise for every $w \in \mathcal{P}(\mathbf{n})^-$.

Remark 2.5.3. In the terminology of [Hru12], these notions corresponds to n -existence / n -uniqueness of T over every parameter set.

Remark 2.5.4. It follows from stability and elimination of imaginaries in T (as then types over algebraically closed sets are stationary) that T has 2-existence, 2 uniqueness and 3-existence.

Let σ be a new unary function symbol, $\mathcal{L}_\sigma := \mathcal{L} \cup \{\sigma\}$. Consider the category $\tilde{\mathcal{C}}_T$ of \mathcal{L}_σ -structures of the form (A, σ) , where $A \in \mathcal{C}_T$ and $\sigma \in \text{Aut}_{\mathcal{L}}(A)$, with \mathcal{L}_σ -embeddings as morphisms.

Definition 2.5.5. Let P equal $\mathcal{P}(\mathbf{n})^-$ or $\mathcal{P}(\mathbf{n})$.

- (1) A functor $A : P \rightarrow \tilde{\mathcal{C}}_T$ is called *independence preserving* (*bounded*, respectively), if it is so when composed with the forgetful functor from $\tilde{\mathcal{C}}_T$ to \mathcal{C}_T .
- (2) We say that $\tilde{\mathcal{C}}_T$ has n -existence if every bounded independence preserving functor $A^- : \mathcal{P}(\mathbf{n})^- \rightarrow \tilde{\mathcal{C}}_T$ extends to a bounded independence preserving functor $A : \mathcal{P}(\mathbf{n}) \rightarrow \tilde{\mathcal{C}}_T$.

It follows from 2-uniqueness and 2-existence in T that $\tilde{\mathcal{C}}_T$ has 2-existence. Let T_σ be the \mathcal{L}_σ -theory of all $(M, \sigma) \in \tilde{\mathcal{C}}_T$ such that $M \models T$. Recall that if T_σ admits a model-companion, it is denoted by TA . If this is the case, we will say that “ TA exists”. (We refer to [CP98] for fundamental facts about TA .)

Fact 2.5.6 ([Hru12, Proposition 4.7 and Corollary 4.10]). *For T as above, the following are equivalent:*

- (1) T has 3-uniqueness.
- (2) $\tilde{\mathcal{C}}_T$ has 3-existence.

Moreover, assuming in addition that TA exists, the above conditions imply:

- (3) TA eliminates imaginaries. □

It is easy to see that if TA exists and it eliminates bounded hyperimaginaries (e.g., when T is superstable, by [BPW01] combined with [CP98, Corollary 3.8]), then (3) is actually equivalent to (1) and (2). We will not use this in our paper.

2.5.2. *Twisted independent amalgamation.* Let $\tau : \mathcal{L} \cong \mathcal{L}'$ be a bijection between two first order languages (sending sorts to sorts, function symbols to function symbols consistently with their arity, similarly for constants and relations). Then τ extends naturally to a bijection between the set of \mathcal{L} -formulas and the set of \mathcal{L}' -formulas. Given an \mathcal{L} -formula φ , we denote by φ^τ its image under this map. If T is an \mathcal{L} -theory, $T^\tau := \{\varphi^\tau : \varphi \in T\}$ is an \mathcal{L}' -theory. Of course, up to changing the names of the symbols using τ , T^τ is the “same” theory as T .

If M is an \mathcal{L} -structure, we denote by M^τ the \mathcal{L}' -structure with base set M and interpretations $(\Sigma^\tau)^{M^\tau} = \Sigma^M$, for any symbol $\Sigma \in \mathcal{L}$. If N' is an \mathcal{L}' -structure, we call an \mathcal{L}' -isomorphism $\sigma : M^\tau \cong N'$ a τ -twisted isomorphism between M and N' . Similarly, one defines the notion of a τ -twisted elementary map $\sigma : A \rightarrow A'$, where $A \subseteq M$ and $A' \subseteq N'$, i.e., one requires that for any \mathcal{L} -formula $\varphi(x)$ and any tuple a from A of the right length, one has $M \models \varphi(a)$ if and only if $N' \models \varphi^\tau(\sigma(a))$.

Lemma 2.5.7. *Let T be a complete stable \mathcal{L} -theory eliminating quantifiers and imaginaries. Assume that T has n -uniqueness. Let $A : \mathcal{P}(\mathbf{n}) \rightarrow \mathcal{C}_{\mathbf{T}}$ and $A' : \mathcal{P}(\mathbf{n}) \rightarrow \mathcal{C}_{\mathbf{T}^\tau}$ be bounded independence preserving functors.*

Then for any coherent system $(\sigma_w)_{w \in \mathcal{P}(\mathbf{n})^-}$ of τ -twisted elementary bijections $\sigma_w : A(w) \rightarrow A'(w)$ there exists a τ -twisted elementary bijection $\sigma_{\mathbf{n}} : A(\mathbf{n}) \rightarrow A'(\mathbf{n})$ extending σ_w for every $w \in \mathcal{P}(\mathbf{n})^-$.

Proof. The result follows from n -uniqueness of T^τ , since we may consider A as a functor to \mathcal{C}_{T^τ} , replacing $A(w)$ by $A(w)^\tau$. \square

We now consider the special case where $\mathcal{L}' = \mathcal{L}$, τ is a permutation of \mathcal{L} and T is a complete \mathcal{L} -theory such that $T = T^\tau$. Let $\tilde{\mathcal{C}}_T^{(\tau)}$ be the category of \mathcal{L}_σ -structures (B, σ) with $B \in \mathcal{C}_T$ and $\sigma : B \rightarrow B$ a τ -twisted elementary bijection. When T is stable, we use the same terminology as in Definition 2.5.5, for functors $A : \mathcal{P} \rightarrow \tilde{\mathcal{C}}_T^{(\tau)}$. Lemma 2.5.7 then yields:

Corollary 2.5.8. *Let T be a complete stable \mathcal{L} -theory eliminating quantifiers and imaginaries, and let $\tau : \mathcal{L} \rightarrow \mathcal{L}$ be a bijection such that $T^\tau = T$. Then $\tilde{\mathcal{C}}_T^{(\tau)}$ has 2-existence. If in addition we assume that T has 3-uniqueness, then $\tilde{\mathcal{C}}_T^{(\tau)}$ has 3-existence.*

Proof. Lemma 2.5.7 yields that if T has n -uniqueness and n -existence, then $\tilde{\mathcal{C}}_T^{(\tau)}$ has n -existence. We may thus conclude by Remark 2.5.4 \square

Given a complete \mathcal{L} -theory T and a permutation τ of \mathcal{L} such that $T^\tau = T$, we let $T_\sigma^{(\tau)}$ be the \mathcal{L}_σ -theory whose models are of the form (M, σ) , where $M \models T$ and where σ is a τ -twisted automorphism of M .

Assume now in addition that T is stable and eliminates quantifiers and imaginaries. It follows from quantifier elimination in T that $T_\sigma^{(\tau)}$ is then a $\forall\exists$ -theory, and so it has a model-companion if and only if the e.c. models of $T_\sigma^{(\tau)}$ form an elementary class. If this is the case, denote by $T^{(\tau)}A$ the

model-companion of $T_\sigma^{(\tau)}$. Then the models of $T^{(\tau)}A$ are precisely the e.c. models of $T_\sigma^{(\tau)}$. The basic results on TA , due to Chatzidakis and Pillay ([CP98]), generalize to this context in a straightforward manner. We will only state some facts which we will need.

Lemma 2.5.9. *Let T and τ be as above, and assume that $T^{(\tau)}A$ exists. Then the following holds:*

(1) *If $(M, \sigma) \models T^{(\tau)}A$ and $B \subseteq M$ then*

$$\text{acl}_{(M, \sigma)}(B) = \text{acl}_\sigma(B) := \text{acl}_M\left(\bigcup_{z \in \mathbb{Z}} \sigma^z(B)\right).$$

(2) *Quantifier reduction: If $(M_i, \sigma_i) \models T^{(\tau)}A$ and $B_i \subseteq M_i$ for $i = 1, 2$, then $B_1 \equiv_{\mathcal{L}_\sigma} B_2$ if and only if there is an \mathcal{L}_σ -isomorphism from $\text{acl}_\sigma(B_1)$ to $\text{acl}_\sigma(B_2)$ sending B_1 to B_2 .*

(3) *$T^{(\tau)}A$ is simple and*

$$A \downarrow_E^{T^{(\tau)}A} B \text{ if and only if } \text{acl}_\sigma(EA) \downarrow_{\text{acl}_\sigma(E)}^T \text{acl}_\sigma(EB).$$

If T is superstable, then $T^{(\tau)}A$ is supersimple.

(4) *Assume that $\tilde{\mathcal{C}}^{(\tau)}$ has 3-existence. (Equivalently, in $T^{(\tau)}A$, the independence theorem holds over acl_σ -closed sets.) Then $T^{(\tau)}A$ eliminates imaginaries.*

Proof. To prove (2), assume that $B = \text{acl}_\sigma(B)$ is a common substructure of two models (M, σ) and (N, σ) of $T^{(\tau)}A$, such that (N, σ) is $|M|^+$ -saturated. We need to show that $(M, \sigma) \mathcal{L}_\sigma(B)$ -embeds into (N, σ) . As $\tilde{\mathcal{C}}_T^{(\tau)}$ has 2-existence, there is an amalgam $(A, \sigma) \in \tilde{\mathcal{C}}_T^{(\tau)}$ of N and M over B . Enlarging (A, σ) if necessary, we may assume that $(A, \sigma) \models T^{(\tau)}A$, hence $(A, \sigma) \succcurlyeq (N, \sigma)$. In particular, $\text{tp}_{\mathcal{L}_\sigma}(M/B)$ is finitely satisfiable in (N, σ) , so this type is realized in (N, σ) by saturation, yielding an $\mathcal{L}_\sigma(B)$ -embedding of (M, σ) into (N, σ) .

We now prove (1). Let $(M, \sigma) \models T^{(\tau)}A$ and let $B \subseteq M$. Clearly, $\text{acl}_{(M, \sigma)}(B) \supseteq \text{acl}_\sigma(B)$. To prove the other inclusion, it suffices to show that if $B = \text{acl}_\sigma(B)$, then B is algebraically closed in (M, σ) . Let $a \in M \setminus B$, and set $A := \text{acl}_\sigma(Ba)$. Then $(B, \sigma) \subseteq (A, \sigma)$ is an extension in $\tilde{\mathcal{C}}_T^{(\tau)}$.

For $n \in \mathbb{Z}_{>0}$, using 2-existence in $\tilde{\mathcal{C}}_T^{(\tau)}$ and induction, we may construct an extension $(B, \sigma) \subseteq (C_n, \sigma) \in \tilde{\mathcal{C}}_T^{(\tau)}$ such that C_n contains n isomorphic copies $(A_1, \sigma), \dots, (A_n, \sigma)$ of (A, σ) over B , which are \mathcal{L} -independent over B . Replacing (M, σ) by an elementary extension if necessary, using (2) we may assume that $C_n \subseteq M$. Since $B = \text{acl}_\mathcal{L}(B)\mathcal{L}$, $A_i \cap A_j = B$ for any $i \neq j$, so (M, σ) contains n distinct realizations of $\text{tp}_{\mathcal{L}_\sigma}(a/B)$, by part (2). As n was arbitrary, $a \notin \text{acl}_{(M, \sigma)}(B)$.

To show (3), we proceed exactly as in the proof of the corresponding result for TA ([CP98, Corollary 3.8]). If A, B, E are subsets of a model of $T^{(\tau)}A$, we say that A and B are *independent* over E if $\text{acl}_\sigma(EA) \downarrow_{\text{acl}_\sigma(E)}^T \text{acl}_\sigma(EB)$,

where \perp^T denotes forking independence in T . This relation satisfies the all abstract properties of an independence notion that guarantee, by the Theorem of Kim-Pillay (see, *e.g.*, [Wag00, Theorem 2.6.1]), that $T^{(\tau)}A$ is simple and that non-forking is given by the independence notion in question. This is clear for all properties except the independence theorem (over a model). To establish the latter, one shows that every 3-amalgamation problem $A^- : \mathcal{P}(\mathbf{3})^- \rightarrow \tilde{\mathcal{C}}_T^{(\tau)}$ with $A^-(\emptyset) \models T^{(\tau)}A$ has a solution; equivalently, the independence theorem even holds over models of $T_\sigma^{(\tau)}$. The proof of this is identical to the proof of [CP98, Theorem 3.7].

The statement about supersimplicity follows as in the proof of [CP98, Corollary 3.8].

Part (4) is the analog of [Hru12, Proposition 4.7]. Weak elimination of imaginaries in $T^{(\tau)}A$ follows directly from a formalization of Hrushovski's argument by Montenegro and the second author (see [MR21, Proposition 1.17]). But finite sets are coded in models of T , as T eliminates imaginaries by assumption. so they are also coded in the expansion $T^{(\tau)}A$ of T . Hence $T^{(\tau)}A$ eliminates imaginaries. \square

2.5.3. *Linear imaginaries.* Let us now recall some notions from [Hru12, Section 5].

Definition 2.5.10. Let \mathfrak{t} be a theory of fields (possibly with additional structure). A \mathfrak{t} -linear structure M is an \mathcal{L} -structure with a sort \mathbf{k} for a model of \mathfrak{t} , and additional sorts V_i ($i \in I$) denoting finite-dimensional \mathbf{k} -vector spaces, such that the family $(V_i)_{i \in I}$ is closed under tensor products and duals. Each V_i has (at least) the \mathbf{k} -vector space structure. One assumes that \mathbf{k} is stably embedded in M with induced structure given by \mathfrak{t} .

We now fix such a \mathfrak{t} -linear structure M .

- (1) M is said to *have flags* if for any i with $\dim(V_i) > 1$, for some j, k with $\dim(V_j) = \dim(V_i) - 1$, there exists a \emptyset -definable exact sequence $0 \rightarrow V_k \rightarrow V_i \rightarrow V_j \rightarrow 0$. We will call such a short exact sequence a *flag*.
- (2) M is said to *have roots* if for any one-dimensional $V = V_i$, and any $m \geq 2$, there exists a (one-dimensional) $W = V_j$ and a \emptyset -definable \mathbf{k} -linear isomorphism $f : W^{\otimes m} \cong V$.

Let us now mention two results from [Hru12]. The proof of the first one is rather elementary, whereas that of the second one is quite involved.

Fact 2.5.11 ([Hru12, Lemma 5.6]). *The theory of an ACF-linear structure with flags (in any characteristic) eliminates imaginaries.*

The following fact follows from [Hru12, Proposition 5.7] in combination with [Hru12, Proposition 4.3 and Corollary 4.10].

Fact 2.5.12. *Let T be the theory of an ACF₀-linear structure with flags and roots. Then T has 3-uniqueness.*

Our main interest in linear structures stems from the fact that the \mathbf{k} -internal sets in a given model of ACVF give rise to such a structure. For every $M \models \text{ACVF}$ and $A \subseteq \mathcal{G}(M)$, we define

$$\mathbf{Lin}_A := \bigsqcup_{\substack{s \in \mathbf{S}(\text{dcl}_0(A)) \\ \ell \in \mathbb{Z}_{>0}}} s/\ell \mathbf{m} s.$$

In equicharacteristic zero, this is exactly the *stable part* $\bigsqcup_{s \in \mathbf{S}(\text{dcl}_0(A))} s/\mathbf{m} s$ of [HHM08]. In mixed characteristic, however, this is a more complicated structure since it also consists of (free) \mathbf{R}_ℓ -modules — and this more complicated structure is actually needed in Section 4. Note that, by Convention 2.1 and our choice of representation of the geometric sorts, $\mathbf{Lin}_A(M)$ is the set of cosets $c + \ell \mathbf{m} s$ where $s \in \mathbf{S}(\text{dcl}_0(A))$ has a basis in M and $c \in s(M)$.

Lemma 2.5.13. *Let $M \models \text{Hen}_{0,0}$ and $A \subseteq \mathcal{G}(M)$. Then $\mathbf{Lin}_A(M)$, with its $\mathcal{L}_0(A)$ -induced structure, is an $\text{Th}(\mathbf{k}(M))$ -linear structure with flags. Moreover, if $\Gamma(M)$ is divisible, then $\mathbf{Lin}_A(M)$ has roots.*

Proof. We may assume that $\text{dcl}_0(A) \cap \mathcal{G}(M) \subseteq A$. The fact that the residue field \mathbf{k} is stably embedded in $\text{Hen}_{0,0}$, with induced structure that of a pure field, is well known — and follows from the existence of splittings as in Lemma 2.3.6.

Now let V, W be two sorts from $\mathbf{Lin}_A(M)$, *i.e.*, vector spaces over \mathbf{k} of the form $V = a/\mathbf{m}a$, $W = b/\mathbf{m}b$ for some $a \in \mathbf{S}_m(A)$ and $b \in \mathbf{S}_n(A)$ — with bases in M . Then $a \otimes_{\mathcal{O}} b$ is canonically isomorphic to an element c from $\mathbf{S}_{m \cdot n}(A)$, so we may identify $V \otimes_{\mathbf{k}} W$ with $c/\mathbf{m}c$, which is a sort from $\mathbf{Lin}_A(M)$. Similarly, $\check{a} = \text{Hom}_{\mathcal{O}}(a, \mathcal{O})$ can be identified with $\{z \in \mathbf{K}^n : \forall v \in a, \sum z_i v_i \in \mathcal{O}\} \in \mathbf{S}_m(A)$, so $\check{V} = \text{Hom}_{\mathbf{k}}(V, \mathbf{k}) \cong \check{a}/\mathbf{m}\check{a}$ is a sort from \mathbf{Lin}_A as well.

Flags: For $a \in \mathbf{S}_n(A)$ define $a_1 := a \cap (\mathbf{K} \times \{0\}^{n-1})$. Then the projection onto the first coordinate identifies a_1 with an element of $\mathbf{S}_1(A)$. Let $\pi : a \rightarrow \mathbf{K}^{n-1}$ be induced from the projection on the last $n - 1$ coordinates. Then

$$0 \rightarrow \ker(\pi) = a_1 \rightarrow a \rightarrow \pi(a) \rightarrow 0$$

is an A -definable exact sequence of free \mathcal{O} -modules, and $\pi(a) \in \mathbf{S}_{n-1}(A)$ — this follows from the fact that $\pi(a)$ is a finitely generated torsion free \mathcal{O} -submodule of \mathbf{K}^{n-1} of rank $n - 1$. Reducing modulo \mathbf{m} , we conclude.

Roots: Assume $\Gamma(M)$ is divisible. Let $n \geq 1$, and let V be a one-dimensional sort from \mathbf{Lin}_A . Then $V = \gamma \mathcal{O}/\gamma \mathbf{m}$ for some $\gamma \in \Gamma(\text{dcl}_0(A))$. Consider $V_n := \delta \mathcal{O}/\delta \mathbf{m}$, for $\delta = \frac{\gamma}{n}$. The map

$$x/\gamma \mathbf{m} \mapsto y/\delta \mathbf{m} \otimes \dots \otimes y/\delta \mathbf{m} : V \rightarrow V_n^{\otimes n},$$

where $y^n = x$, is well-defined and an A -definable isomorphism of \mathbf{k} -vector spaces defined over A . In particular, \mathbf{Lin}_A has roots. \square

The result above actually holds of the *stable part* $\bigsqcup_{s \in \mathbf{S}(\text{dcl}_0(A))} s/\mathbf{m} s$ in all characteristic (provided \mathbf{k} is stably embedded).

Corollary 2.5.14. *Let $M \models \text{ACVF}_{0,0}$ and $A \subseteq \mathcal{G}(M)$. Then \mathbf{Lin}_A satisfies 3-uniqueness. \square*

2.5.4. Twisted linear imaginaries.

Lemma 2.5.15. *Let \mathfrak{t} be a stable theory of fields, and let T be the theory of a \mathfrak{t} -linear structure such that T eliminates quantifiers. Let τ be a permutation of the language with $T = T^\tau$ such that τ fixes all the formulas on the sort \mathbf{k} . Suppose $\mathfrak{t}A$ exists. Then $T_\sigma^{(\tau)} \cup \mathfrak{t}A$ is the model-companion of $T_\sigma^{(\tau)}$. In particular, this holds for $\mathfrak{t} = \text{ACF}$.*

Proof. Let $(M, \sigma) \models T_\sigma^{(\tau)}$. Then, as $M \models T$, for any sort V from \mathcal{L} there is an M -definable surjection $f : \mathbf{k}(M) \rightarrow V(M)$. For any $N \succeq_{\mathcal{L}} M$, f then also defines a surjection from $\mathbf{k}(N)$ onto $V(N)$, hence $N = \text{dcl}_{\mathcal{L}}(M\mathbf{k}(N))$. Thus, any extension of σ to a τ -twisted automorphism on N is uniquely determined by its restriction to $\mathbf{k}(N)$. It follows that (M, σ) is an e.c. model of $T_\sigma^{(\tau)}$ if and only if $(\mathbf{k}(M), \sigma|_{\mathbf{k}(M)})$ is an e.c. model of \mathfrak{t}_σ . This yields the statement of the lemma. \square

As a special case of Lemma 2.5.15, we get the following.

Remark 2.5.16. Let \mathfrak{t} be a stable theory of fields, and let T be the theory of a \mathfrak{t} -linear structure, such that T eliminates quantifiers. Suppose $\mathfrak{t}A$ exists. Then TA exists and is given by $T_\sigma \cup \mathfrak{t}A$. In particular, this holds for $\mathfrak{t} = \text{ACF}$.

Definition 2.5.17. Let \mathbf{k} be a stably embedded sort in a theory T . An A_0 -definable set D is said to be *internally \mathbf{k} -internal* (over A_0), if there is a tuple $d \in D$ and an A_0d -definable surjection $f : Y \rightarrow D$, where $Y \subseteq \mathbf{k}^n$ for some n .

Lemma 2.5.18. *Let \mathbf{k} be a stably embedded sort in a theory T , and let D be A_0 -definable and internally \mathbf{k} -internal (over A_0). Then $\mathbf{k} \cup D$ is stably embedded (over A_0).*

Proof. Set $D' := \mathbf{k} \cup D$. It follows from the assumptions that D' is A_0 -definable and internally \mathbf{k} -internal over A_0 . Let f be an A_0d -definable surjection as in the definition, with $d \in D'$. Taking the preimage under $f \times \cdots \times f$, one sees that any \mathcal{U} -definable subset X of D'^m is $\mathbf{k}(\mathcal{U})A_0d$ -definable, by stable embeddedness of \mathbf{k} . In particular, X is $A_0D'(\mathcal{U})$ -definable, proving stable embeddedness of D' (over A_0). \square

Proposition 2.5.19. *Let $M \models \text{VFA}_{0,0}^{\text{mult}}$ and $A \subseteq \mathcal{G}(M)$. Then \mathbf{Lin}_A is stably embedded in $\text{VFA}_{0,0}^{\text{mult}}$ and its A -induced structure eliminates imaginaries.*

Proof. Stable embeddedness follows from stable embeddedness of \mathbf{k} — cf. Fact 2.4.7 — and the fact that \mathbf{Lin}_A is internally \mathbf{k} -internal (by naming a basis for every sort).

Now, let T be the theory of $\mathbf{Lin}_A(M)$, with its $\mathcal{L}_0(A)$ -induced structure. By Fact 2.5.11 and Lemma 2.5.13, T eliminates imaginaries. Let τ be the

permutation of $\mathcal{L}_0(A)$ induced by σ . Then τ fixes all the formulas on the sort \mathbf{k} , and we have $T^\tau = T$. It follows from Corollary 2.5.14 and Lemma 2.5.9(4) that $T^{(\tau)}A$ eliminates imaginaries.

Also, by Fact 2.4.7 $(\mathbf{k}(M), \sigma_{\mathbf{k}})$ is a stably embedded pure model of ACFA, and hence $\mathbf{Lin}_A(M) \models T^{(\tau)}A$ by Lemma 2.5.15. Since the A -induced structure on \mathbf{Lin}_A is a definable expansion of its ACF-linear structure with a twisted automorphism, elimination of imaginaries follows — *e.g.*, by [Hru12, Lemma 5.4]. \square

2.5.5. Real linear imaginaries. We conclude these preliminaries with a study of RCF-linear structures.

Definition 2.5.20. An RCF-linear structure with flags is said to be *oriented* if for every sort V of dimension one, the two half lines are \emptyset -definable.

Proposition 2.5.21. *Any oriented RCF-linear structure with flags eliminates imaginaries.*

Proof. Let us first prove a few preliminary results. Let M be a sufficiently saturated and homogeneous oriented RCF-linear structure with flags. First, note that if $0 \rightarrow W \rightarrow V \rightarrow U \rightarrow 0$ is an \emptyset -definable flag, then any translate of W in V is ordered by $a < b$ if $a - b$ is in a fixed half line of W .

Claim 2.5.22. *M is rigid: for every $A \subseteq M$, $\text{acl}(A) = \text{dcl}(A)$.*

Proof. Let X be a non-empty finite A -definable set such that all elements of X have the same type over A . We need to show that X is a singleton. Using tensors, we may assume that X is contained in some sort V . We proceed by induction on $\dim(V)$. Let $0 \rightarrow W \rightarrow V \rightarrow U \rightarrow 0$ be an \emptyset -definable flag for V . By induction, we may assume that X projects to a singleton $b \in U$, *i.e.*, X is contained in a translate $a + W$ of W in V and we can conclude since $a + W$ inherits an \emptyset -definable total order from the ordered group structure on W . \square

Claim 2.5.23. *Let $X \subseteq c + W \subseteq V$ be definable for some \emptyset -definable flag $0 \rightarrow W \rightarrow V \rightarrow U \rightarrow 0$ and some $c \in V$. Then X is coded.*

Proof. Since \mathbf{k} is o -minimal and there is a definable order preserving bijection between $c + W$ and \mathbf{k} , X is a finite union of points and intervals and hence it is coded by its (finite) border. \square

Let $K = \mathbf{k}(M)^a$ and $K \otimes M$ be the structure whose sorts are the sorts V of M interpreted as $K \otimes_{\mathbf{k}(M)} V(M)$, with the field structure on \mathbf{k} , the \mathbf{k} -vector space structure on each V and the tensor, dual and flag structure. Then $K \otimes M$ is an ACF-structure with flags and for every $N \preceq M$, and tuple $c \in M$, since all of the vector spaces have bases in N , $\text{dcl}(Nc) \subseteq \text{acl}_0(Nc)$ where acl_0 (respectively dcl_0) denotes the algebraic (respectively definable) closure in $K \otimes M$. Note also that in $K \otimes M$, we have $\mathbf{k}(\text{dcl}_0(M)) = \mathbf{k}(M)$ and since each of the vector spaces has a basis in M , $\text{dcl}_0(M) \subseteq M$. Since $K \otimes M$

eliminates imaginaries by Fact 2.5.11, it follows that any M -definable set in $K \otimes M$ has a code consisting of elements in M .

By [HHM06, Remark 3.2.2], to prove elimination of imaginaries in M , it suffices to code every definable function $f : V \rightarrow S$, where S is a sort. We will proceed by induction on the dimension of V . Let $0 \rightarrow W \rightarrow V \rightarrow U \rightarrow 0$ be an \emptyset -definable flag for V . Let F be the Zariski closure of the graph of f in $K \otimes M$ — any choice of basis induces a Zariski topology on V , but this topology is independent of the choice of coordinates. For every $c \in V(M)$, since $\text{dcl}(Nc) \subseteq \text{acl}_0(Nc)$, for every $N \preccurlyeq M$, the fiber F_c of F above c is a finite set containing $f(c)$. As was noted above, F has a code in M . Note that any $\sigma \in \text{Aut}(M/\ulcorner f \urcorner)$ can be extended to an automorphism of $K \otimes M$ fixing F . It follows that $F(M)$ is defined over $\ulcorner f \urcorner$ and hence it has a code in $M \cap \ulcorner f \urcorner$. Moreover, by compactness and Claim 2.5.22, in M , we find $\ulcorner F \urcorner$ -definable maps $(h_i)_{i < n}$ such that for all $c \in V(M)$, $F_c(M) = \{h_i(a) : i < n\}$. Now, fix $a \in U(M)$ and $c \in V$ above a . Let f_a be the restriction of f to the fiber $c + W$ above a in V . Let $X_{a,i} = \{x \in c + W : f_a(x) = h_i(x)\}$. By Claim 2.5.23, this set is coded in M . Let $g(a)$ denote the tuple consisting of codes of the $X_{i,a}$. The function g is $\ulcorner f \urcorner$ -definable with domain U , so, by induction, g is coded in M . This concludes the proof since f is $(\ulcorner F \urcorner, \ulcorner g \urcorner)$ -definable. \square

Proposition 2.5.24. *Let $(K, <, v)$ be an ordered field with a non-trivial convex valuation and $A \subseteq \mathcal{G}(A)$. Then $\mathbf{Lin}_A(K)$ is oriented. In particular, if v is henselian and $\mathbf{k}(K) \models \text{RCF}$, then $\mathbf{Lin}_A(K)$ is stably embedded and its A -induced structure eliminates imaginaries.*

Proof. Dimension one sorts in \mathbf{Lin}_A are of the form $\gamma\mathcal{O}/\gamma\mathfrak{m}$ for some $\gamma \in \Gamma(K)$. But this quotient inherits the order on $\gamma\mathcal{O}$; so it is oriented. The rest of the proposition follows from Lemma 2.5.18, Proposition 2.5.21 and Lemma 2.5.13. \square

Remark 2.5.25.

- Any $K \equiv \mathbb{R}((\mathbb{Q}))$, being real closed, admits a unique field ordering which is definable (without parameters).
- Any $K \equiv \mathbb{R}((t))$, admits exactly two field orderings, depending on the sign of a choice of uniformizer π . Both orders are definable (using an imaginary parameter for a half line in $\mathbf{RV}_{1,v(\pi)} := \{\xi \in \mathbf{RV}_1 : v(\xi) = v(\pi)\}$, in particular any element of $\mathbf{RV}_{1,v(\pi)}$).

3. C -MINIMAL DEFINABLE GENERICS

We will now consider generalisations of [Rid19, Theorem 8.7].

Notation. Let $\mathcal{L}_0 = \mathcal{L}_{\text{div}}$ and T_0 be the \mathcal{L}_0 -theory ACVF. Let $\mathcal{L} \supseteq \mathcal{L}_0$ and T be a (complete) \mathcal{L} -theory of valued fields. Let $M \models T$ be sufficiently saturated and homogeneous and $M_0 = M^a \models T_0$. Note that since ACVF eliminates quantifiers, we will implicitly assume that every \mathcal{L}_0 -formula is quantifier free. We will denote by $S_x^0(M)$ the set of (quantifier free) $\mathcal{L}_0(M)$ -types (in M_0) in variables x and whenever $\Psi(x;t)$ is a set of \mathcal{L}_0 -formulas,

$S_x^\Psi(M)$ will denote the set of Ψ -types over M — that is, maximal consistent sets (in M_0) of formulas $\psi(x; a)$ and $\neg\psi(x, a)$ with $\psi \in \Psi$ and $a \in M^t$.

3.1. Main results. In this section we prove the following two density results:

Theorem 3.1.1. *Assume*

- (**C_B**) *T is definably spherically complete;*
- (**C_Γ**) *T has definably complete value group;*
- (**E_k[∞]**) *The full induced theory on \mathbf{k} eliminates \exists^∞ : for any $M \models T$ and $\mathcal{L}(M)$ -definable $(Y_z)_z \subseteq \mathbf{k}$, there exists $n \in \mathbb{Z}_{\geq 0}$ such that $|Y_z| < \infty$ implies $|Y_z| \leq n$;*
- (**E_Γ[∞]**) *The full induced theory on Γ eliminates \exists^∞ : for any $M \models T$ and $\mathcal{L}(M)$ -definable $(Y_z)_z \subseteq \Gamma$, there exists $n \in \mathbb{Z}_{\geq 0}$ such that $|Y_z| < \infty$ implies $|Y_z| \leq n$.*

Then, for every strict pro- $\mathcal{L}(A)$ -definable $X \subseteq \mathbf{K}^x$, with x countable and $A = \text{acl}^{\text{eq}}(A) \subseteq M^{\text{eq}} \models T^{\text{eq}}$, there exists an $\mathcal{L}_0(\mathcal{G}(A))$ -definable $p \in S_x^0(M)$ consistent with X .

In other terms, there exists $N \geq M$ and $a \in X(N)$ such that $\text{tp}_0(a/M)$ is $\mathcal{L}_0(\mathcal{G}(A))$ -definable. Recall (see, e.g., [HL16, Section 2.2]) that a set is strict pro-definable if it is the limit of a small directed system of definable sets with surjective transition maps. In other terms, it is a \star -definable set whose projection on any finite set of variables is definable.

Proof. This is a particular case of Proposition 3.5.1 □

- Remark 3.1.2.**
- Any (non zero) definably complete ordered abelian group Γ is elementarily equivalent to either \mathbb{Z} or \mathbb{Q} . Indeed, Γ is necessarily regular as it clearly cannot have a proper non trivial definable convex subgroup. We may thus assume it is a subgroup of $(\mathbb{R}, +, <)$. If it is not elementarily equivalent to \mathbb{Z} or \mathbb{Q} , it is a dense non divisible subgroup of \mathbb{R} . For any $\gamma \in \Gamma$ non divisible by $n \in \mathbb{Z}_{>0}$, the cut at γ/n yields a counter-example.
 - Hypotheses (**C_B**) and (**C_Γ**) are necessary for the conclusion to hold.
 - Hypothesis (**E_Γ[∞]**) does not allow for discrete value groups. Note however that the conclusion of the theorem fails in p -adic fields. So the hypotheses (**C_B**), (**C_Γ**) and (**E_k[∞]**) cannot be sufficient.
 - As Theorem 3.1.3 illustrates, by restricting to a mild class of enrichments of ACVF_v , one can trade hypothesis (**E_Γ[∞]**) for purely algebraic conditions and a weaker conclusion.

Let Hen_0 be the \mathcal{L}_0 -theory of characteristic zero henselian valued fields.

Theorem 3.1.3 (cf. Corollary 3.5.6). *Let T be a \mathbf{k} - Γ -enrichment of Hen_0 , such that:*

- (**C_Γ**) *T has definably complete value group;*
- (**FR**) *For every $n \in \mathbb{Z}_{>0}$, the interval $[0, v(n)]$ is finite and \mathbf{k} is perfect;*
- (**I_k**) *The residue field \mathbf{k} is infinite;*

(\mathbf{E}_k^∞) *The full induced theory on k eliminates \exists^∞ .*

Then, for every strict pro- $\mathcal{L}(A)$ -definable $X \subseteq \mathbf{K}^x$, with $A = \text{acl}^{\text{eq}}(A) \subseteq M^{\text{eq}} \models T^{\text{eq}}$, there exist an $\text{Aut}(M/\mathcal{G}(A))$ -invariant $p \in S_x^0(M)$ consistent with X .

Note that in this setting k is stably embedded so the full induced structure coincides with the \emptyset -induced structure.

Remark 3.1.4.

- Contrary to Theorem 3.1.1, Theorem 3.1.3 requires finite ramification in mixed characteristic. Even if Theorem 3.1.3 does not apply to characteristic zero non-Archimedean local fields either, *cf.* the stronger [HMR18, Remark 4.7].
- Under the hypotheses of Theorem 3.1.3, locally, we do find definable types: for any *finite* set $\Psi(x;t)$ of \mathcal{L}_0 -formulas, we can find an $\mathcal{L}_0(\mathcal{G}(A))$ -definable $p \in S_x^\Psi(M)$ consistent with X , *cf.* Proposition 3.5.4. This local statement does not hold in characteristic zero non-Archimedean local fields.
- In both theorems, hypothesis (\mathbf{E}_k^∞) is an artefact of our proof. It is necessary to prove certain intermediate results. However, we do not know if it is necessary to prove either theorems. Moreover, these theorems are the only reason hypothesis (\mathbf{E}_k^∞) appears in the imaginary Ax-Kochen-Ershov principle, *cf.* Theorem 6.1.1.

Given these observations, the following questions are quite natural:

Question 3.1.5.

- (1) *Can the density of either invariant or definable types — i.e., the conclusion of either theorem — be proved without assuming (\mathbf{E}_k^∞) ?*
- (2) *Under the hypotheses of Theorem 3.1.3, can we find an $\mathcal{L}_0(\mathcal{G}(A))$ -definable type p ?*
- (3) *Can the hypotheses of Theorem 3.1.3 be weakened to also encompass characteristic zero non-Archimedean local fields?*

3.2. The uniform arity one case. We start by giving a succinct (and slightly more general) presentation of terminology and results from [Rid19, Section 6 and 7].

We define \mathbf{B} to be the (\mathcal{L}_0 -definable) set of closed and open balls in models of T_0 — the field itself is the open ball of radius $-\infty$ and points are closed balls of radius $+\infty$. For every $r \in \mathbb{Z}_{>0}$, $\mathbf{B}^{[r]}$ is the set of finite (potentially empty) subsets of cardinality at most r of \mathbf{B} of the same radius and either all open or all closed. Let also $\mathbf{B}^{[<\infty]} := \bigcup_{r>0} \mathbf{B}^{[r]}$.

For every finite set of balls B , we define $B^\cup := \bigcup_{b \in B} b$ and for every finite sets of balls B_1 and B_2 , we write $B_1 \leq B_2$ if $B_1^\cup \subseteq B_2^\cup$. For every $b_1, b_2 \in \mathbf{B}$, we also define $d(b_1, b_2) := \inf\{v(x_1 - x_2) : x_i \in b_i\}$. Note that this is not a metric on the space of balls since $d(b_1, b_1) = \text{rad}(b_1)$, the radius of b_1 . Finally, for $B_1, B_2 \in \mathbf{B}^{[r]}$, we define $D(B_1, B_2) := \{d(b_1, b_2) : b_i \in B_i\}$. We fix an enumeration of $D(B_1, B_2)$ as follows: for every $i \geq 0$ below the cardinal

of $D(B_1, B_2)$, let $d_i(B_1, B_2)$ be the $i + 1$ -th element of $D(B_1, B_2)$, for every other $i < r^2$, let $d_i(B_1, B_2) = d_{r^2-1}(B_1, B_2)$ be the last element of $D(B_1, B_2)$. The actual enumeration does not actually matter as long as it is chosen uniformly.

Let us now fix a set $\Psi(x; t)$ of \mathcal{L}_0 -formulas, an \mathcal{L}_0 -definable set Λ , an integer r and an \mathcal{L}_0 -definable family $F = (F_\lambda)_{\lambda \in \Lambda}$ of functions $F_\lambda : \mathbf{K}^x \rightarrow \mathbf{B}^{[r]}$. We now wish to give sufficient conditions on Ψ and F that will allow us to proceed with certain classical unary constructions in valued fields, uniformly over realisations of Ψ -types. In particular, it will allow us to describe (local) types in $n + 1$ -variables as generics of balls parametrized by n -variables.

Definition 3.2.1. Let $p \in S_x^\Psi(M)$.

- (1) We say that p is adapted to F if each of the following statement, or its negation, is implied by p :
 - $F_\lambda(x) \sqcap \bigcup_{i < r} F_{\mu_i}(x)$ where $\lambda, \mu_i \in \Lambda(M)$ and $\square \in \{=, \subseteq, \subset, \leq, <\}$;
 - $F_\lambda(x) = F_{\mu_1}^\cup(x) \cap F_{\mu_2}^\cup(x)$, where $\lambda, \mu_i \in \Lambda(M)$;
 - Every ball in $F_\lambda(x)$ is closed;
 - $\text{rad}(F_\lambda(x)) \sqcap d_i(F_{\mu_1}(x), F_{\mu_2}(x))$, where $\lambda, \mu_i \in \Lambda(M)$, $\square \in \{=, \leq\}$ and $i < r^2$;
- (2) We say that F is closed under intersections over p if for every $\lambda, \mu \in \Lambda(M)$, there exists $\epsilon \in \Lambda(M)$ with $p(x) \vdash F_\lambda^\cup(x) \cap F_\mu^\cup(x) = F_\epsilon^\cup(x)$ — and we further assume that there exists $\eta \in \Lambda(M)$ such that $F_\lambda(x) = \{\mathbf{K}\}$.
- (3) We say that F is closed under complement over p if for every $\lambda, \mu \in \Lambda(M)$, with $p(x) \vdash F_\mu(x) \subseteq F_\lambda(x)$, there exists $\epsilon \in \Lambda(M)$ with $p(x) \vdash F_\epsilon(x) = F_\lambda(x) \setminus F_\mu(x)$.

Let now $(\Psi(x; t), F)$ and $p \in S_x^\Psi(M)$ be adapted to F , such that F is closed under intersections and complement over p .

Definition 3.2.2. Let $\lambda \in \Lambda(M)$. We say that F_λ is *irreducible over p* , if for every $\mu \in \Lambda(M)$, $p(x) \vdash F_\mu(x) \subseteq F_\lambda(x)$ implies $p(x) \vdash F_\mu(x) = \emptyset \vee F_\mu(x) = F_\lambda(x)$.

We define $\Lambda_p(M) := \{\lambda \in \Lambda(M) : F_\lambda \text{ irreducible over } p\}$.

Lemma 3.2.3. For every $\lambda \in \Lambda(M)$, there exists finitely many $(\mu_i \in \Lambda_p(M))$ with $p(x) \vdash F_\lambda(x) = \bigcup_i F_{\mu_i}(x)$.

Proof. Let $a \models p$. The lemma now follows, by induction on the cardinality of $F_\lambda(a)$, from closure under complement. \square

Lemma 3.2.4. For every $\lambda, \mu \in \Lambda_p(M)$, we have

$$p(x) \vdash F_\lambda^\cup(x) \cap F_\mu^\cup(x) = \emptyset \vee F_\lambda(x) \leq F_\mu(x) \vee F_\mu(x) \leq F_\lambda(x).$$

Proof. Let $a \models p$. We may assume that the balls in $F_\lambda(a)$ have smaller (or equal) radius than those in $F_\mu(a)$ and if the radii are equal and the balls in $F_\lambda(a)$ are open, so are the balls in $F_\mu(a)$. By closure under intersection,

we find $\epsilon \in \Lambda(M)$ such that $F_\lambda^\cup(a) \cap F_\mu^\cup(a) = F_\epsilon^\cup(a)$. By hypothesis on the radii, $F_\epsilon(a) \subseteq F_\lambda(a)$ and hence, by irreducibility, either $F_\epsilon(a) = \emptyset$ or $F_\epsilon(a) = F_\lambda(a)$. \square

We will later need some further hypotheses (*cf.* Lemma 3.3.1) on Ψ and F leading to the following definition:

Definition 3.2.5. Let $\Psi(x;t)$ be a set of \mathcal{L}_0 -formulas, and $F = (F_\lambda)_{\lambda \in \Lambda} : \mathbf{K}^x \rightarrow \mathbf{B}^{[r]}$ be a definable family of functions — for some \mathcal{L}_0 -definable Λ and integer r . We say that (Ψ, F) is a good presentation if, for any $p \in S_x^\Psi(M)$

- (1) p is adapted to F ;
- (2) F is closed under intersection and complement over p ;
- (3) F has large balls over p , *i.e.*, for every $\lambda, \mu \in \Lambda(M)$ and $i \in \mathbb{Z}_{\geq 0}$:
 - If $p(x) \vdash F_\lambda(x) \neq \mathbf{K}$, there is $\eta \in \Lambda(M)$ such that $p(x) \vdash \text{rad}(F_\eta(x)) = d_i(F_\lambda(x), F_\mu(x)) \wedge \text{“}F_\eta(x) \text{ is closed”} \wedge F_\lambda(x) \leq F_\eta(x)$;
 - If $p(x) \vdash \text{“}F_\lambda(x) \text{ is open”} \vee \text{rad}(F_\lambda(x)) < d_i(F_\lambda(x), F_\mu(x))$, there is $\eta \in \Lambda(M)$ such that, $p(x) \vdash \text{rad}(F_\eta(x)) = d_i(F_\lambda(x), F_\mu(x)) \wedge \text{“}F_\eta(x) \text{ is open”} \wedge F_\lambda(x) \leq F_\eta(x)$.

Let $\Delta(xy; s)$ with $|y| = 1$ be a set of \mathcal{L}_0 -formulas and $G := (G_\omega)_{\omega \in \Omega} : \mathbf{K}^x \rightarrow \mathbf{B}^{[\ell]}$ be an \mathcal{L}_0 -definable family of functions. We say that (Ψ, F) is a *good presentation for Δ* if (Ψ, F) is a good presentation and every M -instance of Δ is a Boolean combination of M -instances of Ψ and formulas $y \in F_\lambda(x)^\cup$ with $\lambda \in \Lambda(M)$. If, moreover, for every $\omega \in \Omega(M)$, there exists $\lambda \in \Lambda(M)$ such that $G_\omega = F_\lambda$. we say that (Ψ, F) is a *good presentation for (Δ, G)* .

An important point is that *finite* good presentations always exist:

Proposition 3.2.6. *Let $\Delta(xy; s)$ be a finite set of \mathcal{L}_0 -formulas with $|y| = 1$ and $(G_\omega)_{\omega \in \Omega} : \mathbf{K}^x \rightarrow \mathbf{B}^{[\ell]}$ be \mathcal{L}_0 -definable. Then there exists a finite set of \mathcal{L}_0 -formulas $\Psi(x;t)$ and an \mathcal{L}_0 -definable $F := (F_\lambda)_{\lambda \in \Lambda} : \mathbf{K}^x \rightarrow \mathbf{B}^{[r]}$ such that (Ψ, F) is a good presentation for (Δ, G) .*

We only sketch the proof — details of the precise encodings can be found in [Rid19, Proposition 6.14, 6.15, 6.18 and 7.12].

Proof. The existence of Ψ and F such that any instance of Δ is an Boolean combination of instances of Ψ and $y \in F_\lambda^\cup(x)$ follows by compactness from the Swiss cheese decomposition. Enlarging F , we may assume it contains G and that condition (3) hold. At any point, enlarging Ψ , we may assume that condition (1) holds. Since the intersection of two balls is either empty or one of these balls, F can be closed under intersection by considering the family of $r+1$ -fold intersections of F . Closure under complement can be obtained by considering the finite Boolean algebra generated by the subsets, appearing in F , of any given F_λ (over some realisation of p). They are generated in (uniformly) finitely many steps and hence can be considered as the elements of one single family. This concludes the proof since the previous two steps preserve condition (3). \square

Remark 3.2.7. A good presentation $(\Psi(x;t), F)$ remains a good presentation as Ψ grows. So, given a set $\Psi(xy;t)$ of \mathcal{L}_0 -formulas and an \mathcal{L}_0 -definable $F := (F_\lambda)_{\lambda \in \Lambda} : \mathbf{K}^x \rightarrow \mathbf{B}^{[r]}$, we say that $(\Psi(xy;t), F)$ is a good presentation if there exists $\Phi(x;t) \subseteq \Psi$ such that $(\Phi(x;t), F)$ is a good presentation.

We now fix a good presentation $(\Psi(xy;t), F)$, with $|y| = 1$, and $p \in S_{xy}^\Psi(M)$. Let (Ψ, F) be the set of \mathcal{L}_0 -formulas $\Psi \cup \{y \in F_\lambda^\cup(x)\}$ in variables xy and parameters $t\lambda$. Let $S_{xy}^{\Psi, F}(M)$ denote the space of (Ψ, F) -types over M (in M_0).

Definition 3.2.8. Let $E \subseteq \Lambda_p(M)$. We define

$$\begin{aligned} \eta_{E,p}(xy) &:= p(xy) \\ &\cup \{y \in F_\lambda^\cup(x) : \lambda \in \Lambda(M) \wedge \exists \mu \in E \ p(xy) \vdash F_\mu(x) \leq F_\lambda(x)\} \\ &\cup \{y \notin F_\lambda^\cup(x) : \lambda \in \Lambda(M) \wedge \forall \mu \in \Lambda(M) \ F_\mu^\cup(x) \cap F_\lambda^\cup(x) \subset F_\mu^\cup(x)\}. \end{aligned}$$

Note that for any $E \subseteq \Lambda_p(M)$, provided $\eta_{E,p}$ is consistent, we have $\eta_{E,p} \in S_{xy}^{\Psi, F}(M)$. Further assuming that p is $\mathcal{L}(M)$ -definable (as a Ψ -type), then $\Lambda_p(M)$ is a $\mathcal{L}(M)$ -definable set, and if $\mathcal{E} \subseteq \Lambda_p$ is an $\mathcal{L}(M)$ -definable then the type $\eta_{\mathcal{E}(M),p}$ is an $\mathcal{L}(M)$ -definable (Ψ, F) -type, provided it is consistent. We denote it $\eta_{\mathcal{E},p}$.

Definition 3.2.9. For every $\mathcal{L}(M)$ -definable maps $f, g : X \rightarrow Y$ and every partial $\mathcal{L}(M)$ -type q concentrating on X , we say that f and g have the same q -germ, and we write $[f]_q = [g]_q$, if $q(x) \vdash f(x) = g(x)$.

We write $\lambda \leq_p \mu$ whenever $p(xy) \vdash F_\lambda(x) \leq F_\mu(x)$. Note that \leq_p is an $\mathcal{L}(M)$ -definable pre-order whose associated equivalence relation is equality of p -germs. Moreover, restricted to Λ_p , by Lemma 3.2.4, there is a largest element, \mathbf{K} , and the \leq_p -upwards closure of any $\lambda \in \Lambda_p \setminus [\emptyset]_p$ is totally ordered: $(\Lambda_p \setminus [\emptyset]_p, \leq_p)$ is a tree.

Definition 3.2.10. Let $\pi(x)$ be a partial $\mathcal{L}(M)$ -type and $A \subseteq M^{\text{eq}}$. We say that π is $\mathcal{L}(A)$ -quantifiable over \mathcal{L} if, for every \mathcal{L} -formula $\varphi(x;t)$, there exists an $\mathcal{L}(A)$ -formula $\theta(t)$ such that $\{b \in M^t : \pi(x) \vdash \varphi(x;b)\} = \theta(M)$. When it exists, we write $\forall_\pi x \varphi(x;t) := \theta(t)$ and $\exists_\pi x \varphi(x;t) = \neg(\forall_\pi x \neg\varphi(x;t))$.

Remark 3.2.11. If, for some set of \mathcal{L}_0 -formulas Δ , p is a complete $\mathcal{L}(A)$ -quantifiable Δ -type over M , then it is $\mathcal{L}(A)$ -definable, as a Δ -type. As we will see in Lemma 3.3.1, under certain hypotheses on T and Δ , the converse also holds.

We can now prove the crucial step in proving Theorems 3.1.1 and 3.1.3: the relative arity one case. Let us now assume that p is $\mathcal{L}(A)$ -quantifiable over \mathcal{L} , where $A = \text{acl}^{\text{eq}}(A) \subseteq M^{\text{eq}}$, and consistent with some $\mathcal{L}(A)$ -definable $X \subseteq \mathbf{K}^{xy}$.

Definition 3.2.12. For every $\lambda, \mu \in \Lambda_p$, let $\lambda \trianglelefteq \mu$ hold whenever $\forall_p xy \ y \in (X_x \cap F_\lambda^\cup(x)) \rightarrow y \in F_\mu^\cup(x)$.

The relation \trianglelefteq is an $\mathcal{L}(A)$ -definable preorder on Λ_p and we denote \equiv the associated equivalence relation. Since \leq_p refines \trianglelefteq on $\Theta_p := \Lambda_p \setminus (\emptyset/\equiv)$, this is also a tree with root \mathbf{K}/\equiv and \equiv -classes are \leq_p -convex.

Lemma 3.2.13. *For every $\lambda \in \Theta_p$, if $\eta_{\lambda/\equiv, p}$ is not consistent with X , then λ/\equiv has finitely many \trianglelefteq -daughters $(\mu_i/\equiv)_{0 \leq i < n} \in \text{acl}^{\text{eq}}(A \upharpoonright \lambda/\equiv)$ in Θ_p/\equiv . Moreover, $n \geq 2$ and $p(xy) \vdash y \in X_x \cap F_\lambda^\cup(x) \rightarrow \bigvee_{i < n} y \in F_{\mu_i}^\cup(x)$.*

Proof. By compactness, there exist $(\nu_i)_{0 \leq i < m} \in \Theta_p(M)$ such that $\nu_i \triangleleft \lambda$ and $p(xy) \vdash y \in X_x \cap F_\lambda^\cup(x) \rightarrow \bigvee_{i < m} y \in F_{\nu_i}^\cup(x)$. The existence of the μ_i now follows from the facts that any $\mu \trianglelefteq \lambda$ is \trianglelefteq -comparable to one of the ν_i and that since the F_{ν_i} are irreducible, the subtree with root λ and leaves $(\nu_i)_{i < m}$ embeds in the lattice of subsets of $\{0, \dots, m-1\}$, which is finite — we refer the reader to [Rid19, Claim 8.4] for details. Finally, if $n = 1$, we would have $\lambda \trianglelefteq \mu_0$, contradicting that μ_0 is a daughter of λ . \square

Let $\overline{\Theta}_p := \{\lambda \in \Theta_p : \forall_p xy \text{ “} F_\mu(x) \text{ is closed”}\}$ and, for every $\lambda \in \Lambda$, let $Y_\lambda := \{F_\mu \in \overline{\Theta}_p : \forall_p xy \text{ rad}(F_\mu(x)) = \text{rad}(F_\lambda(x))\}$.

Lemma 3.2.14. *One of the following holds:*

- *There exists a $\lambda \in \Theta_p$ such that $\lambda/\equiv \in A$ and $\eta_{\lambda/\equiv, p}$ is consistent with X .*
- *For every $n \in \mathbb{Z}_{\geq 0}$, there exists $\lambda/\equiv \in A$ such that $[Y_\lambda]_p$ is finite of cardinality larger than n , where $[Y_\lambda]_p := \{[F_\mu]_p : \mu \in Y_\lambda\}$.*

Proof. Assume that X is consistent with no $\eta_{\lambda/\equiv, p}$, where $\lambda/\equiv \in A$. Then, by Lemma 3.2.13, Θ_p/\equiv admits an initial finitely strictly branching discrete tree — that is every, element has at least two daughters — with every branch infinite. Note that, for every $\lambda \in \Theta_p$ with $\lambda <_p \mathbf{K}$, by the large ball property, there is $\mu \in \Lambda$ with $\lambda \leq_p \mu$ and $\forall_p xy \text{ “} F_\mu(x) \text{ is closed”} \wedge \text{rad}(F_\mu(x)) = \text{rad}(F_\lambda(x))$. We may assume that F_μ is irreducible over p . Then $\lambda = \mu$ or λ is the unique \leq_p -daughter of μ . Note also that, by the large ball property, $\overline{\Theta}_p \cap \mathbf{K}/\equiv \neq \emptyset$. It follows that $\overline{\Theta}_p/\equiv$ also admits an initial finitely branching discrete tree, denoted Ξ_p , with every branch infinite.

Note that, for any two $\mu, \nu \in Y_\lambda$, since $\forall_p xy \text{ rad}(F_\mu(x)) = \text{rad}(F_\nu(x))$, we have that $[F_\mu]_p = [F_\nu]_p$ implies $\mu \equiv \nu$, which implies that $\forall_p xy F_\mu^\cup(x) \cap F_\nu^\cup(x) \neq \emptyset$, which, by irreducibility, implies that $[F_\mu]_p = [F_\nu]_p$, so these three statements are equivalent. In particular, the identity induces a bijection between $[Y_\lambda]_p$ and Y_λ/\equiv .

We now build, by induction, $\lambda_i \in \Lambda$ such that $Y_{\lambda_i/\equiv} \subseteq \Xi_p$ and $|Y_{\lambda_i/\equiv}| = |[Y_{\lambda_i}]_p|$ is finite and strictly increasing. Start with any $\lambda_0 \in \overline{\Theta}_p \cap \mathbf{K}/\equiv$. Then $Y_{\lambda_0} = [F_\lambda]_p$ and $Y_{\lambda_0/\equiv} = \lambda_0/\equiv = \mathbf{K}/\equiv$. If λ_i is built, let $(\mu_j)_{j < m}$ enumerate all the \trianglelefteq -daughters — in Ξ_p — of the elements in $Y_{\lambda_i/\equiv}$. Let j_0 be such that, for all j , $\forall_p xy \text{ rad}(F_{\mu_{j_0}}(x)) \leq \text{rad}(F_{\mu_j}(x))$. For every $\nu \in Y_{\mu_{j_0}}$, by the large ball property, we find $\lambda \in Y_{\lambda_i}$ such that $\nu \leq_p \lambda$. Since $\forall_p xy \text{ rad}(F_\nu(x)) = \text{rad}(F_{\mu_{j_0}}(x)) \leq \text{rad}(F_{\mu_j}(x))$, we cannot have $\nu \triangleleft \mu_j$ and hence ν/\equiv is either in $Y_{\lambda_i/\equiv}$ or it is one of the μ_j/\equiv . So $Y_{\mu_{j_0}/\equiv} \subseteq \Xi_p$ is finite.

Furthermore, for every μ_j , by the large ball property, there exists $\nu \in Y_{\mu_{j_0}}$ such that $\mu_j \leq_p \nu$. It follows that, for every element of Y_{λ_i}/\equiv , either it or all of its (more than one) daughters appear in $Y_{\mu_{j_0}}$. In particular, all the sisters of μ_{j_0}/\equiv appear, and hence $|Y_{\mu_{j_0}}/\equiv| > |Y_{\lambda_i}/\equiv|$. Thus, we can choose $\lambda_{i+1} = \mu_{j_0}$. \square

We can now eliminate the second option in Lemma 3.2.14 by imposing a uniform bound on the size of finite instances of $(Y_\lambda)_{\lambda \in \Lambda}$:

Corollary 3.2.15. *Assume:*

$(\mathbf{E}_{p,\bar{F}}^\infty)$ For every $\mathcal{L}(M)$ -definable family $(Y_z)_z$ of subsets of $[F_{\Lambda_p}]_p := \{[F_\lambda]_p : \lambda \in \Lambda_p\}$ such that for all z and $[F_\lambda]_p, [F_\mu]_p \in Y_z$, $p(xy) \vdash "F_\mu(x) \text{ is closed}" \wedge \text{rad}(F_\lambda(x)) = \text{rad}(F_\mu(x))$, there exists $n \in \mathbb{Z}_{>0}$ such that, for all z , $|Y_z| < \infty$ implies $|Y_z| \leq n$.

Then there exists an $\mathcal{L}(A)$ -definable $\mathcal{E} \subseteq \Lambda_p$ such that $\eta_{\mathcal{E},p}$ is consistent with X . \square

However the family $[Y_\Lambda]_p$ is not any definable family in $[F_{\Lambda_p}]_p$. It has certain geometric properties that reflect that of X . In particular, with further hypotheses on X , we can dispense with $(\mathbf{E}_{p,\bar{F}}^\infty)$ altogether.

We now wish to apply the construction above in the pair (M_0, M) which is naturally an $\mathcal{L}_{0,P}$ structure enriched with the \mathcal{L} -structure on M . To be precise and avoid an unnecessary conflict of notation:

Notation. Let \mathcal{L}_1 be some expansion of \mathcal{L}_0 and T_1 some \mathcal{L}_1 -theory of valued fields. In the following lemma, we apply the above with T the theory of the pair $M := (M_1^a, M_1)$, where $M_1 \models T_1$ is sufficiently saturated and homogeneous, in the language $\mathcal{L} := \mathcal{L}_P$ consisting of $\mathcal{L}_{0,P}$ structure enriched with the \mathcal{L}_1 -structure on \mathbf{P} — so $M_0 = M_1^a$.

Let us now introduce some useful terminology from [CHR19]:

Definition 3.2.16. Fix $n \in \mathbb{Z}_{>0} \cap \mathbf{K}^\times(M)$. For any ball b , we define $b[n] := \{a + n^{-1}(a - c) : a, c \in b\}$. It is a ball of radius $\text{rad}(b) - v(n)$ around b , open if b is, closed otherwise. For a set of balls B , we set $B[n] := \{b[n] : b \in B\}$.

- (1) An $\mathcal{L}(M_1)$ -definable set $X \subseteq \mathbf{K}(M_1)$ is *n-prepared* by some finite set $C \subseteq \mathbf{K}(M_0)$ if for every ball $b \in \mathbf{B}(M_0)$ with $b[n] \cap C = \emptyset$, either $b \cap X(M_1) = \emptyset$ or $b \cap X(M_1) = b(M_1)$.
- (2) We say that some $\mathcal{L}_0(M_1)$ -definable $G : \mathbf{K}^n \rightarrow \mathbf{K}^{[r]}$ *n-prepares* $X \subseteq \mathbf{K}^{n+1}(M_1)$ if, for every $x \in \mathbf{K}(M_1)^n$, $G(x)$ *n-prepares* X_x .
- (3) We say that $X \subseteq \mathbf{K}^{n+1}(M_1)$ is *n-prepared* by F if there exists $\lambda \in \Lambda(M_1)$ such that F_λ *n-prepares* X .

Remark 3.2.17. By field quantifier elimination (*cf.* Fact 2.2.6), if M_1 is a pure henselian field of characteristic zero, any $\mathcal{L}(M_1)$ -definable $X \subseteq \mathbf{K}$ is p^ℓ -prepared, for some ℓ , by the finite set of roots of polynomials that appear in the (field quantifier free) definition of X , where p is either 1 or the residue characteristic when it is positive.

Let also $A_1 = \text{acl}_1^{\text{eq}} \subseteq M_1^{\text{eq}}$, $X \subseteq \mathbf{K}^{xy}(M_1)$ be $\mathcal{L}_1(A_1)$ -definable, $A = A_P = \text{acl}^{\text{eq}}(A)$ and $p \in S_{xy}^\Psi(M_0)$ be $\mathcal{L}_P(A_P)$ -definable and consistent with X .

Lemma 3.2.18. *Let $\Phi(x; t) \subseteq \Psi$ be such that (Φ, F) is a good presentation for Ψ . Assume that there is some $n \in \mathbb{Z}_{>0} \cap \mathbf{K}^\times(M)$ such that:*

- ($\mathbf{P}_X^{F,n}$) *The set X is n -prepared by F ;*
- (\mathbf{FR}_n) *The interval $[0, v(n)] \subseteq \Gamma(M_1)$ is finite and $\mathbf{k}(M_1)$ is perfect;*
- ($\mathbf{I}_\mathbf{k}$) *The residue field $\mathbf{k}(M_1)$ is infinite.*

Then, there exists an $\mathcal{L}_P(A_P)$ -definable $\mathcal{E} \subseteq \Lambda_p(M_0)$ such that $\eta_{\mathcal{E},p}|_{M_0}$ is consistent with X .

Proof. Let ρ be such that $F_\rho(x)$ n -prepares X_x for all x . By Lemma 3.2.14, applied in $M = (M_1^a, M_1)$, either the conclusion of the Lemma holds, or we can find $\lambda/\equiv \in A$ with $||Y_\lambda||_p > r$. Then some element of Y_λ , say F_λ , does not contain any point of $F_\rho(x)$. Replacing λ with any $\mu \trianglelefteq \lambda$ in Ξ_p such that $||[\mu, \lambda]|| \geq ||[0, v(n)]||$, we may further assume that $F_\rho(x) \cap F_\lambda(x)[n]^\cup = \emptyset$ — where, for any $B \in \mathbf{B}^{<\infty}$, $B[n] := \{b[n] : b \in B\}$ — and that $\lambda \triangleleft \mathbf{K}$.

By Lemma 3.2.13, we have $p(xy) \vdash y \in X_x \cap F_\lambda^\cup(x) \rightarrow \forall_{i < n} y \in F_{\mu_i}^\cup(x)$, where the $(\mu_i/\equiv)_{i < n}$ are the daughters of λ/\equiv . By compactness, there exists some $\psi(xy) \in p$ such that $q := p|_\Phi \vdash \forall y \psi(xy) \wedge y \in X_x \cap F_\lambda^\cup(x) \rightarrow \forall_{i < n} y \in F_{\mu_i}^\cup(x)$. Since (Φ, F) is a good presentation for Ψ , there are κ and $(\mu_i)_{n \leq i < m} \in \Lambda_p$ with $\vDash \psi(xy) \leftrightarrow y \in F_\kappa^\cup(x) \setminus (\bigcup_{n \leq i < m} F_{\mu_i}^\cup(x))$. In particular, $p(xy) \vdash y \in F_\kappa^\cup(x)$. It follows that $\kappa \equiv \mathbf{K}$ and hence $\lambda \trianglelefteq \kappa$. So, we have

$$q \vdash F_\lambda^\cup(x) \cap X_x \subseteq \bigcup_i F_{\mu_i}^\cup(x).$$

Since $\lambda \neq \emptyset$, there exists $ac \vDash p$ in some $N_1 \geq M_1$ such that $c \in X_a \cap F_\lambda^\cup(a)$. Let $b_0 \in \mathbf{B}(N_1^a)$ be the ball of $F_\lambda(a)$ containing c . Since $F_\rho(a) \cap b_0[n] = \emptyset$, we have $b_0 \cap X_a(N_1) = b_0(N_1)$. It follows that $b_0(N_1) \subseteq \bigcup_i F_{\mu_i}^\cup(a)$. By construction, $b_0(N_1)$ is not covered by any single ball in $\bigcup_i F_{\mu_i}^\cup(a)$. So $\{v(x - y) : x, y \in b_0(N_1)\}$ has a minimal element (realised by some x, y in distinct balls of $\bigcup_i F_{\mu_i}^\cup(a)$). Let $b \in \mathbf{B}(N_1)$ be the smallest closed ball containing $b_0(N_1)$. Then $b(N_1) = b_0(N_1)$ is covered by finitely many of its maximal open subballs, contradicting hypothesis ($\mathbf{I}_\mathbf{k}$). \square

3.3. Quantifiable types. To use the above constructions in an induction, we need a number of results on quantifiable types. The first one is that $\eta_{\mathcal{E},p}$ is itself quantifiable when p is. For any finite set B of balls, let \mathbf{k}_B be the set of maximal open subballs of the balls $b \in B$ and $\text{res}_B : B^\cup \rightarrow \mathbf{k}_B$ be the projection.

Lemma 3.3.1 (cf. [Rid19, Corollary 6.9]). *Let $(\Psi(xy; t), (F_\lambda(x))_{\lambda \in \Lambda})$ be a good presentation. Let $p \in S_{xy}^\Psi(M)$ be $\mathcal{L}(A)$ -quantifiable over \mathcal{L} , where $A \subseteq M^{\text{eq}}$. Assume:*

- ($\mathbf{E}_{p,F}^\infty$) *For any $\mathcal{L}(M)$ -definable $(Y_z)_z \subseteq [F_\Lambda]_p$ and $\lambda \in \Lambda(M)$ such that, for all z and $[F_\mu]_p \in Y_z$, $p(xy) \vdash F_\mu(x) \subseteq \mathbf{k}_{F_\lambda(x)}$, there exists $n \in \mathbb{Z}_{\geq 0}$ such that, for all z , $|Y_z| < \infty$ implies $|Y_z| \leq n$.*

Then, any $\mathcal{L}(A)$ -definable $q \in S_{xy}^{\Psi, F}(M)$ containing p is $\mathcal{L}(A)$ -quantifiable over \mathcal{L} .

Proof. Let $\mathcal{E} := \{\lambda \in \Lambda_p : q(xy) \vdash y \in F_\lambda^\cup(x)\}$. Then \mathcal{E} is $\mathcal{L}(A)$ -definable and $q = \eta_{\mathcal{E}, p}$. If \mathcal{E} does not have an \leq_p -minimal element which is closed, for any \mathcal{L} -formula $\varphi(xy; s)$ and $e \in M^s$, $q(xy) \vdash \varphi(xy; e)$ if and only if there exists $\lambda \in \mathcal{E}(M)$ and $\mu \in \Lambda_p(M)$ with $\lambda <_p \mathcal{E}$ and $q(xy) \vdash y \in F_\lambda^\cup(x) \setminus F_\mu^\cup(x) \rightarrow \varphi(xy; e)$ — cf. [Rid19, Proposition 6.4]. So let us assume that \mathcal{E} has an \leq_p -minimal element λ_0 which consists of closed balls. If $p(xy) \vdash \text{rad}(F_{\lambda_0}(x)) = +\infty$, then $q(xy) \vdash \varphi(xy; e)$ if and only if $p(xy) \vdash F_{\lambda_0}^\cup(x) \rightarrow \varphi(xy; e)$ — cf. [Rid19, Proposition 6.6]. If $p(xy) \vdash \text{rad}(F_{\lambda_0}(x)) \neq +\infty$, let

$$Y_s := \{\mu \in \Lambda_p : \forall_p xy \ F_\mu(x) \subseteq \mathbf{k}_{F_\lambda(x)} \wedge \exists_p xy \ \varphi(xy; s) \wedge y \in F_\mu^\cup(x)\}.$$

Let n be a uniform bound on the cardinality of finite $[Y_s]_p$, as in $(\mathbf{E}_{p, F}^\infty)$.

Claim 3.3.2. *For every $e \in M^s$, q is consistent with $\varphi(xy; e)$ if and only if, for every $(\mu_i)_{i < n} \in \Lambda_p$, with $\forall_p xy F_{\mu_i}(x) \in \mathbf{k}_{F_{\lambda_0}(x)}$, $\exists_p xy \ \varphi(xy; e) \wedge y \in F_{\lambda_0}^\cup(x) \setminus \bigcup_{i < n} F_{\mu_i}^\cup(x)$.*

Proof. Assume q is not consistent with $\varphi(xy; e)$, then, by compactness, there exists $(\mu_i)_{i < m} \in \Lambda_p$ such that $\mu_i <_p \mathcal{E}$ and $\forall_p xy \ \varphi(xy; e) \wedge y \in F_{\lambda_0}^\cup(x) \rightarrow \bigvee_{i < m} y \in F_{\mu_i}^\cup(x)$. By the large ball property, we may assume $\forall_p xy \ F_{\mu_i}(x) \in \mathbf{k}_{F_{\lambda_0}(x)}$. Choosing a minimal m , we may also assume that, $\exists_p xy \ \varphi(xy; e) \wedge y \in F_{\mu_i}^\cup(x)$. In particular, $\mu_i \in Y_e$.

By definition of Y_s , for every $\mu \in Y_e(M)$, we find $ac \models p$ such that $\varphi(ac; e)$ and $c \in F_\mu^\cup(a) \subseteq F_{\lambda_0}^\cup(a)$. So there is an i such that $c \in F_{\mu_i}^\cup(a)$. By irreducibility, $F_\mu(a) = F_{\mu_i}(a)$. It follows that $[Y_e]_p$ is finite and thus $m \leq |[Y_e]_p| \leq n$. \square

Since $q \vdash \varphi(xy; e)$ if and only if q is not consistent with $\neg\varphi(xy; e)$, Claim 3.3.2 allows us to conclude. \square

We also need a better understanding of the interpretable set $[F_\Lambda]_p$. Note that it is, a priori, $\mathcal{L}(M)$ -interpretable which is exactly the kind of sets elimination of imaginaries aims at describing. However, if p happens to be the restriction to M of a global $\mathcal{L}_0(M)$ -definable type, then $[F_\Lambda]_p$ naturally embeds in an $\mathcal{L}_0(M)$ -interpretable set. The goal of the following lemmas is to give (necessary) hypotheses under which any definable p verifies that condition. Valued vector spaces will play an important role:

Definition 3.3.3. Let (K, v) be a valued field and V be a K -vector space. A *valuation on V* is a map $v : V \rightarrow X$ where X is an ordered set with a maximal element ∞ and an action $+$ of Γ , respecting the order, such that:

- $v(0) = \infty$;
- For all $x, y \in V$, $v(x + y) \geq \min\{v(x), v(y)\}$;
- For all $a \in K$ and $x \in V$, $v(a \cdot x) = v(a) + v(x)$.

We say that a family $(x_i)_i \in V$ is *separating* if for every almost everywhere zero $(a_i)_i \in K$, $v(\sum_i a_i x_i) = \min_i (v(a_i) + v(x_i))$.

The following lemma owes much to Johnson's computation of the canonical basis of definable types in ACVF, cf. [Joh20, Section 5.2]. Let $\varphi_d(x; yz) := v(\sum_{|I|<d} y_I x^I) \geq v(\sum_{|I|<d} z_I x^I)$.

Proposition 3.3.4. *Assume:*

(C_V) *Every $\mathcal{L}(M)$ -definable valuation v on \mathbf{K}^n has a separating basis;*

(C_T) *T has definably complete value group.*

Then, for every $A = \text{dcl}^{\text{eq}}(A) \subseteq M^{\text{eq}}$, $\mathcal{L}(A)$ -definable $p \in S_{\varphi_d}(M)$ and algebraic extension $K(M) \leq L$, any $q \in S_{\varphi_d}(L)$, extending p and finitely satisfiable in M , is $\mathcal{L}_0(\mathcal{G}(A))$ -definable.

Proof. For every field F with $K = \mathbf{K}(M) \leq F \leq L$, we define a valuation v on the F -vector space $V_d(F) := F[x]_{\leq d}$ of polynomials in variables x over F of degree at most d by $v(P(x)) \leq v(Q(x))$ if $v(P(x)) \leq v(Q(x)) \in q$. The valuation v on $V_d(K)$ is $\mathcal{L}(A)$ -definable. By hypothesis (C_V), it has a separating basis $(P_i)_i \in V_d(K)$.

Claim 3.3.5. *$(P_i)_i$ is a separating basis of $V_d(L)$ over L .*

Proof. We may assume that $K \leq L$ is finite. By [Joh21, Remark 2.7], the valuation on L (interpreted in K) is then $\mathcal{L}(M)$ -definable and hence, by hypothesis (C_V), also has a separating basis $(c_j)_j \in L$ over K . Let $b_i = \sum_j b_{i,j} c_j \in L$, where $b_{i,j} \in K$. If $v(\sum_j (\sum_i b_{i,j} P_i) c_j) > \min_j v(\sum_i b_{i,j} P_i) + v(c_j)$, since q is finitely satisfiable in M , we have $v(\sum_j (\sum_i b_{i,j} P_i(a)) c_j) > \min_j v(\sum_i b_{i,j} P_i(a)) + v(c_j)$ for some $a \in M$, contradicting that $(c_j)_j$ is separating over K . So

$$\begin{aligned} v(\sum_i b_i P_i) &= v(\sum_j (\sum_i b_{i,j} P_i) c_j) \\ &= \min_j v(\sum_i b_{i,j} P_i) + v(c_j) \\ &= \min_{i,j} v(b_{i,j}) + v(c_j) + v(P_i) \\ &\leq \min_i v(b_i) + v(P_i) \\ &\leq v(\sum_i b_i P_i) \end{aligned} \quad \square$$

We now define the L -Archimedean equivalence on $v(V_d(L))$: $v(P) \sim_L^\infty v(Q)$ holds if there exists $c \in L^\times$ such that $-v(c) + v(P) \leq v(Q) \leq v(c) + v(P)$. One can check that $|v(V_d(L))/\sim_L^\infty| \leq |v(V_d(L))/v(L)| \leq \dim_L(V_d(L)) + 1 < \infty$. We also define the L -infinitesimal equivalence on $v(V_d(L))$: $v(P) \sim_L^0 v(Q)$ holds if for every $\gamma \in v(L)_{>0}$ we have $-\gamma + v(P) < v(Q) < \gamma + v(P)$. Note that two elements of the same $v(L)$ -orbit cannot be \sim_L^0 -equivalent unless they are equal. It follows that \sim_L^0 -classes are finite.

Let C be any K -Archimedean class and \overline{C} denote its upwards closure. Then $V_C := v^{-1}(\overline{C}) \leq V_d(K)$ is a sub- K -vector space.

Claim 3.3.6 ([Joh20, Observation 4.3]). *V_C has a basis of elements in A .*

Proof. Some coordinatewise projection $V_C \subseteq K^l \rightarrow K^m$ restricts to an isomorphism on V_C . The preimage of the standard basis of K^m then has the required properties. \square

Let C_0 be the successor of C in $V_d(K)/\sim_K^\infty$. Since $V_{C_0} \subset V_C$, any basis of V_C has an element outside V_{C_0} . In particular $(V_C \setminus V_{C_0})(A) \neq \emptyset$ and we find $\gamma_C \in C(A) \neq \emptyset$. Then the whole (finite) K -infinitesimal class of γ_C is in A . Let i be such that $v(P_i) \in C$. By (\mathbf{C}_Γ) , the set $\{\gamma \in v(K) : \gamma + v(P_i) \leq \gamma_C\}$ has a supremum γ_i . Multiplying P_i by some constant $c \in K$ with $v(c) = -\gamma_i$, we may assume that $\gamma_i = 0$ in which case $v(P_i)$ is K -infinitesimally close to $\gamma_C \in A$ and hence is itself in A . Since every $v(K)$ -orbit is contained in some K -Archimedean class, we now have that for any i , $v(P_i) \in A$ and for any j , if $v(P_i) \sim_K^\infty v(P_j)$, then $v(P_i) \sim_K^0 v(P_j)$.

Note that, since $v(L)$ is in the convex hull of $v(K)$, \sim_L^∞ extends \sim_K^∞ . Also, if $v(K)$ is dense, then \sim_L^0 extends \sim_K^0 . However, if $v(K)$ is discrete then \sim_K^0 reduces to equality. In particular, we also have that, if $v(P_i) \sim_L^\infty v(P_j)$, then $v(P_i) \sim_L^0 v(P_j)$.

For every i, d , let $M_{i,d}(L) = \{P \in V_d(L) : v(P) \geq v(P_i)\}$. Note that, by Claim 3.3.5, $\sum \lambda_j P_j \in M_{i,d}(L)$ if and only if:

- $\lambda_j = 0$ for every j with $v(P_j) < v(P_i)$ and $v(P_j) \not\sim_K^\infty v(P_i)$;
- $\lambda_j \in \mathfrak{m}$ for j with $v(P_j) < v(P_i)$ and $v(P_j) \sim_K^0 v(P_i)$;
- $\lambda_j \in \mathcal{O}$ for j with $v(P_i) \leq v(P_j)$ and $v(P_j) \sim_K^0 v(P_i)$.

So $M_{i,d}$ is (quantifier free) $\mathcal{L}_0(\mathcal{G}(A))$ -definable. Since q is $\mathcal{L}_0(\bigcup_{i,d} \ulcorner M_{i,d} \urcorner)$ -definable, it is indeed $\mathcal{L}_0(\mathcal{G}(A))$ -definable. \square

Remark 3.3.7. Any definably complete ordered abelian group is elementarily equivalent to either \mathbb{Q} , \mathbb{Z} or 0 .

If $p \in S^0(M)$ the existence (and uniqueness) of such a q follows, on general grounds, from the finite satisfiability of p :

Lemma 3.3.8. *Let $p \in S^0(M)$ be finitely satisfiable in M . Then any two realisations of p have the same $\mathcal{L}_0(\text{acl}_0(M))$ -type. In particular, the unique extension of p to $\text{acl}_0(M)$ is finitely satisfiable in M .*

Proof. Fix any $c \in \text{acl}_0(M)$, $\varphi(xy)$ an \mathcal{L}_0 -formula and $\psi(y)$ an $\mathcal{L}_0(M)$ -formula that algebraises c . Then $\forall y [\psi(y) \rightarrow (\varphi(x_1y) \leftrightarrow \varphi(x_2y))]$ defines an $\mathcal{L}_0(M)$ -definable equivalence relation with finitely many classes. Then the E -class of any $a \in N \geq M$ realising p has an element $e \in M$. It follows that $p(x) \vdash xEe$. In particular $p(x) \vdash \varphi(xe)$ whenever $\varphi(ec)$ holds.

Let q be the unique extension of p to $\text{acl}_0(M)$. Then $p \vdash q$ and hence q is finitely satisfiable in M . \square

Following [Bau82], we can prove that (\mathbf{C}_V) follows from definable spherical completeness. Up to definability, this is a standard result. But we include its proof, on the one hand, for the sake of completeness, and, on the other, to show that the proof can indeed be done definably.

Lemma 3.3.9. *Assume:*

(C_B) T is definably spherically complete : any $\mathcal{L}(M)$ -definable chain of balls has a non-empty intersection.

Then any (finite dimensional) $\mathcal{L}(M)$ -interpretable valued \mathbf{K} -vector space (V, v) has a separating basis.

Proof. Let us proceed by induction on $n + 1 := \dim(V)$. In particular, we may assume that we have found a separating family $(y_i)_{0 \leq i < n} \in V$.

Claim 3.3.10. For every $x \in V$, $\{v(x - \lambda y) : \lambda \in \mathbf{K}^n\}$ has a maximal element.

Proof. For every $\lambda, \mu \in \mathbf{K}^n$, we have $v(x - \mu y) \geq v(x - \lambda y) =: \gamma$ if and only if $\min_i \{v(\mu_i - \lambda_i) + v(y_i)\} = v((\mu - \lambda)y) \geq \gamma$. For every $i < n$ and $\lambda \in \mathbf{K}^n$, let $B_{i,\lambda} := \{\mu \in \mathbf{K} : v(x - \lambda_{\neq i} y_{\neq i} - \mu y_i) \geq v(x - \lambda y)\} = \{\mu \in \mathbf{K} : v(\mu - \lambda_i) + v(y_i) \geq v(x - \lambda y)\}$. They form a chain for inclusion.

If there is a minimal B_{i,λ_0} , pick any $\lambda_i \in B_{i,\lambda_0}$. If there is no minimal $B_{i,\lambda}$, for every $\gamma \in v(\mathbf{K})$, let $b_{i,\gamma}$ be the closed ball of radius γ containing some $B_{i,\lambda}$, if it exists, or \mathbf{K} otherwise. Since the chain of $B_{i,\lambda}$ does not have a minimal element, any $B_{i,\lambda}$ contains a $b_{i,\gamma}$ that itself contains a $B_{i,\mu}$. By definable spherical completeness, we find $\lambda_i \in \bigcap_{\gamma} b_{i,\gamma} = \bigcap_{\lambda} B_{i,\lambda}$. Then λ_i has the property that, for any $\mu \in \mathbf{K}^n$, $v(x - \mu y) \leq v(x - \mu_{\neq i} y_{\neq i} - \lambda_i y_i)$. It follows that $v(x - \lambda y)$ is maximal. \square

Let x be linearly independent from the y_i . By the Claim 3.3.10, we may assume that $v(x) = \max\{v(x - \lambda y) : \lambda \in \mathbf{K}^n\}$. Then, for every $\mu \in \mathbf{K}$ and $\lambda \in \mathbf{K}^n$, we have $v(\mu x + \lambda y) \leq v(\mu x)$ and thus, $v(\mu x + \lambda y) = \min\{v(\mu x), v(\lambda y)\} = \min_i \{v(\mu x), v(\lambda_i y_i)\}$. \square

3.4. Counting germs. The last ingredient in this section is to reduce the (seemingly horrendous) hypotheses $(\mathbf{E}_{p,\tilde{F}}^\infty)$ and $(\mathbf{E}_{p,\tilde{F}}^\infty)$ to something more tractable. We start by generalizing [Rid19, Section 7]. Let $M \models T$, $M_0 := M^a \models T_0$ and Q be either Γ or \mathbf{k} .

Lemma 3.4.1. Let $(f_\lambda)_{\lambda \in \Lambda} : \mathbf{K}^n \rightarrow Q$ be an \mathcal{L}_0 -definable family and $c \in \mathbf{K}(N_0)^n$, where $N_0 \geq M_0$. There exists an $\mathcal{L}_0(M)$ -definable family $(g_\rho)_{\rho \in R} : \mathbf{K}^n \rightarrow Q^{[\leq \infty]}$, with $R \subseteq Q^m$, such that for all $\lambda \in \Lambda(M_0)$, there exists $\rho \in R(M_0)$ with $f_\lambda(c) \in g_\rho(c)$.

In particular, if $p \in S^0(M_0)$ is $\mathcal{L}_0(M)$ -definable, there is a $\mathcal{L}_0(M)$ -definable finite-to-one map $\{[f_\lambda]_p : \lambda \in \Lambda(M_0)\} \rightarrow Q^m$, for some $m \in \mathbb{Z}_{>0}$.

Proof. We start with the non-uniform version of the result:

Claim 3.4.2. For every $N_0 \models \text{ACVF}$, $A \leq \mathbf{K}(N_0)$ and finite tuple $c \in \mathbf{K}(N_0)$, there exists a finite tuple $a \in A$, such that $Q(\text{acl}_0(Ac)) \subseteq \text{acl}_0(Q(A)ac)$.

Proof. If $|c| = 1$, we find $a_0 \in \text{acl}_0(A)$ with $Q(\text{acl}_0(Ac)) = \text{acl}_0(Q(A(c))) \subseteq \text{acl}_0(Q(\text{acl}_0(A))ca_0) = \text{acl}_0(Q(A)ca_0)$, by the characterization of transcendental 1-types in ACVF. If $a \in A$ is such that $a_0 \in \text{acl}_0(a)$, we indeed have $Q(\text{acl}_0(Ac)) \subseteq \text{acl}_0(Q(A)ac)$. If $c = de$ with $|e| = 1$, we proceed by induction: $Q(\text{acl}_0(Ade)) \subseteq \text{acl}_0(Q(\text{acl}_0(Ad))be) \subseteq \text{acl}_0(Q(A)acbe)$, with $a, b \in A$. \square

By Claim 3.4.2, and compactness in a saturated model of the pair (N_0, M_0) , there exists an $\mathcal{L}_0(M_0)$ -definable g as above. The union of its conjugates over M has the same properties and is $\mathcal{L}_0(M)$ -definable.

Now, if $p \in S^0(M_0)$ is $\mathcal{L}_0(M)$ -definable, then for any $\lambda \in \Lambda(M_0)$, let $Y_\lambda := \{\rho : \forall_p x f_\lambda(x) \in g_\rho(x)\}$ and $h([f_\lambda]_p) := \ulcorner Y_\lambda \urcorner \in Q^m$. Note that $p(x) \models f_\mu(x) \in \bigcap_{\rho \in Y_\mu} g_\rho(x)$, so h is finite-to-one. \square

Lemma 3.4.3. *Assume:*

(\mathbf{E}_k^∞) For any $\mathcal{L}(M)$ -definable $(Y_z)_z \subseteq \mathbf{k}$, there exists $n \in \mathbb{Z}_{\geq 0}$ such that, for all z , $|Y_z| < \infty$ implies $|Y_z| \leq n$;

Then, for every $\mathcal{L}_0(M)$ -definable $p \in S_x^0(M_0)$ and \mathcal{L}_0 -definable $(F_\lambda)_{\lambda \in \Lambda} : \mathbf{K}^x \rightarrow \mathbf{B}^{[r]}$, $(\mathbf{E}_{p, \bar{F}}^\infty)$ holds, uniformly in λ .

Proof. The core of the proof is the following almost internality result:

Claim 3.4.4. For every $\lambda \in \Lambda(M)$, there exists an $\mathcal{L}_0(M)$ -definable finite-to-one map $g_\lambda : X_\lambda := \{[F_\mu]_p : \mu \in \Lambda(M_0) \text{ and } p(x) \models \text{``}F_\mu(x) \text{ are maximal open balls of } F_\lambda(x)\text{''}\} \rightarrow \mathbf{k}^m$, for some $m \in \mathbb{Z}_{>0}$.

Proof. Let $c \models p$. Working in $M(c)^a$ and then taking unions of Galois conjugates, we may find $G_i(c) \in \mathbf{B}^{[<\infty]}$, for $i := 1, 2$, two $\mathcal{L}_0(Mc)$ -definable sets picking at least one maximal open balls in each of the ball of F_λ and such that $G_1(c) \cap G_2(c) = \emptyset$. For every μ with $[F_\mu]_p \in X_\lambda$, let $f_\mu(x) := \{(b - b_1)/(b_2 - b_1) : b \in F_\mu(x), b_i \in G_i(x) \text{ and } b, b_1, b_2 \text{ are in the same ball of } F_\lambda(x)\} \in \mathbf{k}^{[<\infty]}$. Note that $F_\mu(c) \in \text{acl}_0(f_\mu(c)G_1(c)G_2(c))$. Using symmetric functions, we identify $\mathbf{k}^{[<\infty]}$ with some \mathbf{k}^n .

By Lemma 3.4.1, we find an $\mathcal{L}_0(M)$ -definable finite to one map $h : \{[f_\mu]_p : [F_\mu]_p \in X_\lambda\} \rightarrow \mathbf{k}^m$. Then $[F_\mu]_p \in \text{acl}_0([G_1]_p[G_2]_p[f_\mu]_p) \subseteq \text{acl}_0(Mh(\mu))$ and $h(\mu) \in \text{dcl}_0(M[f_\mu]_p) \subseteq \text{dcl}_0(M[F_\mu]_p)$. \square

Since $\mathbf{k}(\text{dcl}_0(M))$ is the perfect closure of $\mathbf{k}(M)$, by compactness, composing with a power of the Frobenius automorphism, we may assume that $g_\lambda(X_\lambda(M)) \subseteq \mathbf{k}(M)$ and that g_λ is uniform in λ . The (uniform) bound in $(\mathbf{E}_{p, \bar{F}}^\infty)$ now follows from (\mathbf{E}_k^∞) . \square

Lemma 3.4.5. *Assume (\mathbf{E}_k^∞) and:*

$(\mathbf{E}_\Gamma^\infty)$ For any $\mathcal{L}(M)$ -definable $(Y_z)_z \subseteq \Gamma$, there exists $n \in \mathbb{Z}_{\geq 0}$ such that, for all z , $|Y_z| < \infty$ implies $|Y_z| \leq n$.

Then for every $\mathcal{L}_0(M)$ -definable $p \in S_x^0(M_0)$, \mathcal{L}_0 -definable $(F_\lambda)_{\lambda \in \Lambda} : \mathbf{K}^x \rightarrow \mathbf{B}^{[r]}$, and $\mathcal{L}(M)$ -definable $(Y_z)_z \subseteq [F_\Lambda]_p$, there exists $n \in \mathbb{Z}_{>0}$ such that $|Y_z| < \infty$ implies $|Y_z| \leq n$.

In particular, $(\mathbf{E}_{p, \bar{F}}^\infty)$ holds.

Proof. Let $r_\lambda(x) := \text{rad}(F_\lambda(x))$. By Lemma 3.4.1, there exists an $\mathcal{L}_0(M)$ -definable finite-to-one map $g : [r_\Lambda]_p \rightarrow \Gamma^m$. Composing by division by a fixed integer, we may assume that $g([r_\Lambda]_p(M)) \subseteq \Gamma(M)$. It now follows that there is a bound on finite $\{[r_\lambda]_p : [F_\lambda]_p \in Y_z\}$. So, cutting each Y_z in finitely many

pieces (and getting rid of the infinite ones), we may assume that $[\text{rad}(F_\lambda)]_p$ is constant and the balls are of the same type, as $[F_\lambda]_p$ ranges through Y_z . Similarly, we may assume that the set of distances between balls in $F_\lambda(x)$ and $F_\mu(x)$, with $[F_\lambda]_p, [F_\mu]_p \in Y_z$ has size bounded by some integer k . We now proceed by induction on k .

Let $\gamma_z(x)$ be the smallest such distance, $G_{\lambda,z}(x)$ be the set of closed balls of radius $\gamma_z(x)$ around $F_\lambda(x)$ and $Z_z := \{[G_{\lambda,z}]_p : [F_\lambda]_p \in Y_z\}$. Then the set of distances between balls in Z_z has size at most $k-1$ and we find a bound by induction. In particular, removing some more infinite Y_z , we find an $H_z : \mathbf{K}^n \rightarrow \mathbf{B}^{[\lt \infty]}$ such that every maximal open ball of $H_z(x)$ contains at most one ball of $F_\lambda(x)$ as $[F_\lambda]_p$ varies through Y_z . The bound now follows from Lemma 3.4.3. \square

3.5. The higher arity case. We can now proceed with the induction:

Proposition 3.5.1. *Assume (\mathbf{C}_V) , (\mathbf{C}_Γ) , (\mathbf{E}_k^∞) and $(\mathbf{E}_\Gamma^\infty)$. Let $X \subseteq \mathbf{K}^x$ be strict pro- $\mathcal{L}(A)$ -definable, where $A = \text{acl}^{\text{eq}}(A) \subseteq M^{\text{eq}}$ and x is countable. Let $\Delta(x;t)$ be a finite set of \mathcal{L}_0 -formulas, $p \in S_x^\Delta(M)$ be $\mathcal{L}(A)$ -quantifiable over \mathcal{L} and consistent with X and $z \subseteq x$. Then, there exists an $\mathcal{L}_0(\mathcal{G}(A))$ -definable $q \in S_z^0(M_0)$ such that $q|_M$ is consistent with p and X .*

Proof. We proceed by induction on $|z|$. In particular, we may assume that for any set $\Delta(x;t)$ of \mathcal{L}_0 -formulas, $p \in S_x^\Delta(M)$ which is $\mathcal{L}(A)$ -quantifiable over \mathcal{L} and finite strictly smaller $w \subset z$, $p|_w$ can be extended to an $\mathcal{L}_0(\mathcal{G}(A))$ -definable $q \in S_w^0(M_0)$.

Claim 3.5.2. *Let $\Delta(x;t)$ be a finite set of \mathcal{L}_0 -formulas, $p \in S_x^\Delta(M)$ be $\mathcal{L}(A)$ -quantifiable over \mathcal{L} and consistent with X , finite $z \subseteq x$ and $\Phi(z;s)$ be a finite set of \mathcal{L}_0 -formulas. Then, there exists a finite set $\Theta(z;t)$ containing Φ and $q \in S_x^{\Delta, \Theta}(M)$ which is $\mathcal{L}(A)$ -quantifiable over \mathcal{L} and consistent with p and X .*

Proof. We proceed by induction on $|z|$. Assume $z = wy$ with $|y| = 1$. By Proposition 3.2.6, we find a finite good presentation $(\Psi(w;t), F(w))$ for Φ . By induction, we find $\Xi(w;u) \supseteq \Psi$ and $q \in S_x^{\Delta, \Xi}(M)$ which is $\mathcal{L}(A)$ -quantifiable over \mathcal{L} and consistent with p and X . Since $w \subset z$, as stated in the first paragraph of the proof, $q|_w$ extends to a complete $\mathcal{L}_0(\mathcal{G}(A))$ -definable $\mathcal{L}_0(M_0)$ -type.

By Corollary 3.2.15 and Lemma 3.4.5, we now find an $\mathcal{L}(A)$ -definable $r \in S_x^{\Delta, \Xi, F}(M)$ which is consistent with q and X . By Lemmas 3.3.1 and 3.4.3, r is $\mathcal{L}(A)$ -quantifiable over \mathcal{L} . \square

Let $(\varphi_i(z; t_i))_{i \in \omega}$ enumerate all \mathcal{L}_0 -formulas. By Claim 3.5.2, we find Θ_i containing φ_i and $q_i \in S_z^{\Delta, \Theta_{\leq i}}(M)$, which is $\mathcal{L}(A)$ -quantifiable over \mathcal{L} and consistent with $p \cup \bigcup_{j < i} q_j$ and X . Then $\bigcup_i q_i \in S_z^0(M)$ is $\mathcal{L}(A)$ -definable and consistent with p and X . By Proposition 3.3.4 and Lemma 3.3.8, q extends to a complete $\mathcal{L}_0(\mathcal{G}(A))$ -definable $\mathcal{L}_0(M_0)$ -type. \square

This result is already non-trivial when $X = \mathbf{K}^x$ and $T = \text{ACVF}$:

Corollary 3.5.3. *Let $\Psi(x; t)$ be a set of \mathcal{L}_0 -formulas and $A = \text{acl}_0(A) \leq M_0$. Any $\mathcal{L}_0(A)$ -quantifiable $p \in S_x^\Psi(M_0)$ can be extended to an $\mathcal{L}_0(A)$ -definable $q \in S_x^0(M_0)$. \square*

If we do not assume $(\mathbf{E}_\Gamma^\infty)$, and try to replace the use of Corollary 3.2.15 by that of Lemma 3.2.18, the above induction fails. We can, nevertheless, recover a local version of the result:

Proposition 3.5.4. *Let $n \in \mathbb{Z}_{>0} \cap \mathbf{K}^\times(M)$ and $X \subseteq \mathbf{K}^x$ be $\mathcal{L}(A)$ -definable, where $A = \text{acl}^{\text{eq}}(A) \subseteq M^{\text{eq}}$. Assume (\mathbf{C}_V) in both M and the pair (M_0, M) , (\mathbf{C}_Γ) , $(\mathbf{E}_\mathbf{k}^\infty)$, $(\mathbf{I}_\mathbf{k})$, (\mathbf{FR}_n) . Also assume that:*

$(\mathbf{P}_{\pi(X)}^n)$ *For every projection $Y \subseteq \mathbf{K}^{zy}$ of X , with $|y| = 1$, $N \geq M$ and $a \in N^y$, Y_a is n -prepared by some finite $\mathcal{L}_0(Ma)$ -definable set $C \subseteq \mathbf{K}$.*

Then, for every finite set $\Psi(x, t)$ of \mathcal{L}_0 -formulas, there exists an $\mathcal{L}_0(\mathcal{G}(A))$ -definable $p \in S_x^\Psi(M)$ consistent with X .

Proof. Let $A_P = \text{acl}_{\mathcal{L}_P}^{\text{eq}}(A)$. We say that $\Theta(zy, s)$, where $|y| = 1$, is a hereditarily good presentation if Θ is of the form $\Phi(z, t) \cup \{y \in F_\lambda(z)\}$ for some good presentation (Φ, F) where Φ is itself an hereditarily good presentation.

Claim 3.5.5. *Let $\Theta(x, s)$ be a hereditarily good presentation and $q \in S_x^\Theta(M_0)$ be $\mathcal{L}_0(\mathcal{G}(A_P))$ -definable. Then for $\mathcal{L} \in \{\mathcal{L}_0, \mathcal{L}_P\}$, q is $\mathcal{L}(\mathcal{G}(A_P))$ -quantifiable over \mathcal{L} — in particular it has a complete $\mathcal{L}_0(\mathcal{G}(A_P))$ -definable extension to $S_x^0(M_0)$.*

Proof. We proceed by induction on $|x|$. Since $(\mathbf{E}_\mathbf{k}^\infty)$ holds both in M_0 and, by Lemma 2.3.19, in the pair (M_0, M) , quantifiability follows from Lemmas 3.3.1 and 3.4.3 — applied respectively to M_0 and to the pair (M_0, M) . The existence of a complete definable extension follows by Corollary 3.5.3 applied in M_0 . \square

We now prove, by induction on $x = zy$, the existence of $p \in S_x^\Psi(M_0)$ which is $\mathcal{L}_0(\mathcal{G}(A_P))$ -definable and consistent with X . By compactness, there exists $(G_\omega)_{\omega \in \Omega} : \mathbf{K}^z \rightarrow \mathbf{K}^{[<\infty]}$ such that the family $(X_z)_z$ is n -prepared by G . Let $d \in \mathbb{Z}_{>0}$ bound the degree of any polynomial appearing in Ψ and G . By Proposition 3.2.6, and induction, we find a finite hereditarily good presentation $(\Theta(z, s), F(z))$ for $\varphi_d(x, uv) := v(\sum_{|I|<d} u_I x^I) \geq v(\sum_{|I|<d} v_I x^I)$. By induction, there exists $q \in S_z^\Theta(M_0)$ which is $\mathcal{L}_0(\mathcal{G}(A_P))$ -definable and consistent with X . By Claim 3.5.5, q is $\mathcal{L}_P(\mathcal{G}(A_P))$ -quantifiable over \mathcal{L}_P . By Lemma 3.2.18, there exists an $\mathcal{L}_P(A_P)$ -definable $p \in S_x^{\Theta, F}(M_0)$ consistent with X . By hypothesis, (\mathbf{C}_V) holds in (M_0, M) , and so does (\mathbf{C}_Γ) , by Corollary 2.3.17. By Proposition 3.3.4, it follows that $p|_{\varphi_d}$ is $\mathcal{L}_0(\mathcal{G}(A_P))$ -definable — and hence so is $p|_\Psi$.

Let $a \in A_P$ be the canonical basis of $p|_{\varphi_d}$. Since $M_0 = M^a \subseteq \text{acl}_0(M)$, we have $a \in \text{acl}_0(M)$. Let $c \in \text{dcl}_0(M)$ be a code of the finite $\mathcal{L}_0(M)$ -orbit

of a — which is included in its finite $\mathcal{L}_P(A)$ -orbit. Let f be \mathcal{L}_0 -definable such that $c \in f(M)$ and $e \in M^{\text{eq}}$ be a code of $f^{-1}(c)$. The $\mathcal{L}_P(A)$ -orbit of c consists of finite subsets of the $\mathcal{L}_P(A)$ -orbit of a and is therefore finite. Hence, so is the $\mathcal{L}(A)$ -orbit of e ; *i.e.*, $e \in \text{acl}^{\text{eq}}(A) = A$. It follows that $p|_{\varphi_a, M} \subseteq \bigcap_{\sigma \in \text{Aut}(M_0/M)} \sigma(p|_{\varphi_a})$ is $\mathcal{L}(A)$ -definable. By Proposition 3.3.4, it is in fact $\mathcal{L}_0(\mathcal{G}(A))$ -definable — and hence so is $p|_{\Psi, M}$. \square

This local result does imply the existence of a global invariant type:

Corollary 3.5.6. *Let $n \in \mathbb{Z}_{>0} \cap \mathbf{K}^\times(M)$ and $X \subseteq \mathbf{K}^x$ be strict pro- $\mathcal{L}(A)$ -definable, where $A = \text{acl}^{\text{eq}}(A) \subseteq M^{\text{eq}}$. Assume (\mathbf{C}_V) in both M and the pair (M_0, M) , (\mathbf{C}_Γ) , $(\mathbf{E}_\mathbf{k}^\infty)$, $(\mathbf{I}_\mathbf{k})$, (\mathbf{FR}_n) and:*

$(\mathbf{P}_{\pi(X)}^n)$ *For every projection $Y \subseteq \mathbf{K}^{zy}$ of X onto finitely many coordinates, with $|y| = 1$, every $N \geq M$ and every $a \in \mathbf{K}^z(N)$, Y_a is n^ℓ -prepared by some $\mathcal{L}_0(Ma)$ -definable set $C \subseteq \mathbf{K}(N)$, for some $\ell \in \mathbb{Z}_{\geq 0}$.*

Then there exists an $\text{Aut}(M/\mathcal{G}(A))$ -invariant $p \in S_x^0(M)$ consistent with X .

Note that (\mathbf{FR}_n) implies (\mathbf{FR}_{n^ℓ}) for every $\ell \in \mathbb{Z}_{\geq 0}$. Also if M is a finitely ramified henselian field, (\mathbf{C}_V) holds in both M and (M_0, M) , since both theories have maximally complete models, and $(\mathbf{P}_{\pi(X)}^n)$ holds by Remark 3.2.17.

Proof. For every (finite) set $\Psi(x, t)$ of \mathcal{L}_0 -formulas, the set of $p \in S_x^0(M)$ which are consistent with X and whose Ψ -type is $\text{Aut}(M/\mathcal{G}(A))$ -invariant is closed. It is non-empty, by Proposition 3.5.4. By compactness, the intersection of all these sets, which coincides with the set of $\text{Aut}(M/\mathcal{G}(A))$ -invariant $p \in S_x^0(M)$ consistent with X , is also non-empty. \square

Remark 3.5.7. If T is a $\mathbf{k}\text{-}\Gamma$ -enrichment of Hen_0 , then the pair (M_0, M) is elementarily equivalent to one where both \mathbf{K} and \mathbf{PK} are maximally complete. Hence (\mathbf{C}_B) — and therefore (\mathbf{C}_V) — holds both in M and in the pair (M_0, M) .

4. INVARIANT COMPLETIONS

Notation. In this section. let T be an \mathbf{RV} -enrichment of the theory of characteristic zero henselian fields.

4.1. Main results. Our goal in this section is to describe the behaviour of global \mathcal{L} -types whose underlying \mathcal{L}_0 -type is invariant. A crucial point is that Fact 2.2.6 can be reformulated in the following manner: for every $A \leq M \models T$,

$$\text{tp}_0(A) \cup \text{tp}(\mathbf{RV}(A)) \vdash \text{tp}(A).$$

Therefore, the main point of this section is to better understand $\text{tp}(\mathbf{RV}(A))$ and then deduce properties of $\text{tp}(A)$. In particular, we will show that $\mathbf{RV}(A)$ is generated by a small canonical set. This will allow us to conclude that a global type whose underlying quantifier free type is invariant is itself invariant over \mathbf{RV} (*cf.* Corollary 4.3.17). However, a better control of the

parameters requires more auxiliary sorts. Recall that

$$\mathbf{Lin}_A := \bigsqcup_{\substack{s \in \mathbf{S}(\text{dcl}_0(A)) \\ \ell \in \mathbb{Z}_{>0}}} s/\ell \text{ms.}$$

In this section, we prove:

Theorem 4.1.1. *Assume*

(**I_k**) *The residue field \mathbf{k} is infinite.*

Let $M \preceq N \models T$ sufficiently saturated and homogeneous, $A \subseteq \mathcal{G}(M)$ and $a \in \mathbf{K}(N)$ such that $\text{tp}_0(a/M)$ is $\text{Aut}(M/A)$ -invariant. Then the type $\text{tp}(a/M)$ is $\text{Aut}(M/\text{ARV}(M)\mathbf{Lin}_A(M))$ -invariant.

4.2. Invariance and stably embedded sets. Note that we consider invariance over large subsets of our model — that happen to be the points of some stably embedded definable sets. This gives rise to some subtle issues and two notions of invariance. When $D = \bigcup_i D_i$ is ind- \mathcal{L} -definable, we denote by D^{eq} the ind- \mathcal{L} -definable union of all \mathcal{L} -interpretable sets X that admit an \mathcal{L} -definable surjection $\prod_j D_{i_j} \rightarrow X$.

Definition 4.2.1. Let M be an \mathcal{L} -structure, $C \subseteq M$, D be a (ind-) \mathcal{L} -definable set and p be a partial $\mathcal{L}(M)$ -type. We say that p :

- is $\text{Aut}(M/C)$ -invariant if for every $\sigma \in \text{Aut}(M/C)$, p and $\sigma(p)$ are equivalent.
- has $\text{Aut}(M/C)$ -invariant D -germs if it is $\text{Aut}(M/C)$ -invariant and so is the p -germ of every $\mathcal{L}(M)$ -definable map $f : p \rightarrow D^{\text{eq}}$;
- is $\text{Aut}(M/D)$ -invariant if it has $\text{Aut}(M/D(M))$ -invariant D -germs.

We will only apply these notions for M saturated, p a complete Δ -type for some set of \mathcal{L} -formulas Δ and C equal to the M -points of a stably embedded (ind-) \mathcal{L} -definable set, D stably embedded — an ind- \mathcal{L} -definable set $D = \bigcup_i D_i$ is stably embedded if any definable $X \subseteq \prod_j D_{i_j}$ is definable with parameters from D .

- Remark 4.2.2.**
- (1) A type might be $\text{Aut}(M/D(M))$ -invariant but not $\text{Aut}(M/D)$ -invariant. For example, let $M \models \text{ACVF}$ and b be a closed ball of M without any $\text{acl}({}^r b^1 \mathbf{RV}_1(M))$ -definable subballs. Then any two $a_1, a_2 \in b(M)$ have the same type over $\text{acl}({}^r b^1 \mathbf{RV}_1(M))$. However, for every $x \in b$, $\text{rv}_1(x - a_1) = \text{rv}_1(x - a_2)$ implies that the a_i are in the same maximal open subball of b . It follows that the generic of b over M is $\text{Aut}(M/{}^r b^1)$ -invariant but not $\text{Aut}(M/{}^r b^1 \mathbf{RV}_1)$ -invariant.
 - (2) A type $p \in \mathbf{S}(M)$ is $\text{Aut}(M/C)$ -invariant if and only if for every realization $a \models p$ in a sufficiently homogeneous $N \succeq M$, any $\sigma \in \text{Aut}(M/C)$ extends to an element of $\text{Aut}(N/Ca)$.
 - (3) On the other hand, a type $p \in \mathbf{S}(M)$ has $\text{Aut}(M/C)$ invariant D -germs, where D is stably embedded, if and only if for every $a \models p$ in a sufficiently saturated $N \succeq M$, any $\sigma \in \text{Aut}(M/C)$ extends to an element of $\text{Aut}(N/CD(N)a)$ — cf. the proof of Lemma 4.2.4.

- (4) The space of types with $\text{Aut}(M/C)$ -invariant D -germs is closed: for any $\sigma \in \text{Aut}(M/C)$ and $\mathcal{L}(M)$ -definable map $f : p \rightarrow D^{\text{eq}}$, no types in the open set “ $[f]_p \neq [f^\sigma]_p$ ” has $\text{Aut}(M/C)$ -invariant D -germs.

Let us now recall the following folklore result on stable embeddedness which states that we can recover the usual characterisation of types and hence of definable closure (equivalently internality) from invariance over a stably embedded definable set:

Lemma 4.2.3. *Let M be saturated sufficiently large, D be (ind-) \mathcal{L} -definable stably embedded and $e \in M$. If e is fixed by every $\sigma \in \text{Aut}(M/D(M))$, then $e \in \text{dcl}(D(M))$.*

Proof. Let $e' \in M$ be such that $e \equiv_{D(M)} e'$. By [TZ12, Lemma 10.1.5] — more precisely its extension *mutatis mutandis* to stably embedded ind-definable sets — we can find $\sigma \in \text{Aut}(M/D(M))$ such that $e' = \sigma(e) = e$. Since D is stably embedded, there exists a small $A \subseteq D(M)$ such that, $\text{tp}(e/A) \vdash \text{tp}(e/D(M))$. So both types have a single realisation in M , i.e., $e \in \text{dcl}(A) \subseteq \text{dcl}(D(M))$. \square

One advantage of the stronger notion of invariance is transitivity:

Lemma 4.2.4. *Let $M \preceq N$ be \mathcal{L} -structures with N saturated and sufficiently large, $C \subseteq M$ (potentially large), D be an (ind-) \mathcal{L} -definable stably embedded set, $p \in S(M)$ have $\text{Aut}(M/C)$ -invariant D -germs, $a \models p$ in N and $q \in S(N)$ be $\text{Aut}(N/CD(N)a)$ -invariant. Then $q|_M$ is $\text{Aut}(M/C)$ -invariant. If, moreover, q has $\text{Aut}(N/CD(N)a)$ -invariant E -germs, for some (ind-) \mathcal{L} -definable set E , then $q|_M$ has $\text{Aut}(M/C)$ -invariant E -germs.*

Proof. Fix $\sigma \in \text{Aut}(M/C)$ and let τ be the partial \mathcal{L} -elementary isomorphism $M(a) \rightarrow M(a)$ induced by σ . Since σ fixes the germs of every definable map from p to D^{eq} , τ induces the identity on $D^{\text{eq}}(\text{dcl}(M(a)))$. Since D is stably embedded, cf. [TZ12, Lemma 10.1.5], τ extends to an element of $\text{Aut}(N/CD(N)a)$, also denoted τ , which thus fixes q . It follows that $q|_M$ is fixed by $\tau|_M = \sigma$.

If, moreover, q has $\text{Aut}(N/CD(N)a)$ -invariant E -germs, then $\sigma = \tau|_M$ fixes the $q|_M$ -germ of any $\mathcal{L}(M)$ -definable function into E^{eq} . \square

The core of our proof of Theorem 4.1.1 is the following variation on transitivity:

Lemma 4.2.5. *Let $M \preceq N \models T$, $C \subseteq M$ potentially large, $a \in \mathbf{K}^x(N)$ a (potentially infinite) tuple and $\rho : \mathbf{K}^x \rightarrow \mathbf{RV}$ be pro- $\mathcal{L}_0(M)$ -definable. Assume that $\text{rv}_\infty(M(a)) \subseteq \text{dcl}_0(C\rho(a))$ and that $p := \text{tp}_0(a/M)$ and $[\rho]_p$ are $\text{Aut}(M/C)$ -invariant. Then $\text{tp}(a/M)$ has $\text{Aut}(M/C)$ -invariant \mathbf{RV} -germs.*

Proof. Pick $\sigma \in \text{Aut}(M/C)$. Let $N_0 \models \text{ACVF}$ containing N be saturated and sufficiently large. Let $\tau : M(a) \rightarrow M(a)$ be the \mathcal{L}_0 -isomorphism induced by σ . Note that $\tau(\rho(a)) = \rho^\sigma(a) = \rho(a)$ and hence $\tau|_{(\text{rv}_\infty(M(a)))}$ is the identity. We may thus extend τ to a partial elementary map which is the identity on

$\mathbf{RV}(N_0)$. So, by stable embeddedness of \mathbf{RV} , τ extends to some element of $\text{Aut}(N_0/C\mathbf{RV}(N_0)a)$, also denoted τ . By Fact 2.2.6, $\text{tp}(Ma) = \text{tp}(\sigma(M)a)$, *i.e.*, $\sigma(p) = p$. Moreover, any $\mathcal{L}(Ma)$ -definable $X \subseteq \mathbf{RV}^n$ is $\mathcal{L}(\text{rv}_\infty(M(a)))$ -definable and hence $X(N) = \tau(X(N)) = X^\tau(N)$, equivalently, σ fixes the p -germ of any $\mathcal{L}(M)$ -definable function into \mathbf{RV}^{eq} . \square

4.3. Computing leading terms. In view of Lemma 4.2.5, given any A -invariant type $\text{tp}_0(a/M)$, we want to find a pro- $\mathcal{L}_0(M)$ -definable map ρ such that $\rho(a)$ generates $\text{rv}_\infty(M(a))$ and $[\rho]_p$ is $\text{Aut}(M/A)$ -invariants. When $A \preceq M$ and A is sufficiently large, this is done in Corollary 4.3.16. As previously stated, for general small A , dealing with closed balls forces us to also consider maps into certain A -definable \mathbf{k} -vector spaces. The goal then becomes to build a “nice” model of T containing A and proceed by transitivity.

The technical core of the proof consists in a generalisation to relative arity one of the classical description of 1-types in henselian fields, cf. Lemmas 4.3.8, 4.3.10 and 4.3.13.

Let us start with three leading term computations that we will need later.

Lemma 4.3.1. *Let $M \models \text{ACVF}$, $L \preceq K = \mathbf{K}(M)$, $\text{rv}_1(L) \preceq R \preceq \text{rv}_1(K)$, $b \in \mathbf{B}(K)$, $g \in \mathbf{K}(\text{dcl}_0(LR))$, $c \in K$ and $P := \prod_{i < d} (x - e_i) \in \mathbf{K}(\text{dcl}_0(LR))[x]$.*

- (1) *If $c, g \in b$ and $e_i \notin b$, for all i , then $\text{rv}_1(P(c)) = \text{rv}_1(P(g)) \in \text{dcl}_0(R)$.*
- (2) *If $c \notin b$ and $g, e_i \in b$, for all i . then $\text{rv}_1(P(c)) = \text{rv}_1(c - g)^d$.*
- (3) *Assume that b is closed, that $c, g, e_i \in b$, for all i , and that the maximal open subball of b around c does not contain g nor any e_i . Then*

$$\text{rv}_1(P(c)) = \bigoplus_{i \leq d} \text{rv}_1(P_i(g)) \text{rv}_1(c - g)^i \in \text{dcl}_0(R \text{rv}_1(c - g)),$$

where $P(y + x) = \sum_i P_i(y)x^i$. In particular, the sum is well-defined.

In fact, computation (2) is an easy particular case of computation (3) — consider the smallest closed ball containing c , g and the e_i .

Proof. Let us first prove (1). We have $\text{rv}_1(P(c)) = \prod_i \text{rv}_1(c - e_i) = \prod_i \text{rv}_1(g - e_i) = \text{rv}_1(P(g)) \in \mathbf{RV}_1(\text{dcl}_0(LR)) \subseteq \text{dcl}_0(\text{rv}_1(L)R) = \text{dcl}_0(R)$, where the inclusion follows from quantifier elimination for ACVF in $\mathcal{L}_{\mathbf{RV}}$. As for (2), we have $\text{rv}_1(P(c)) = \prod_i \text{rv}_1(c - e_i) = \text{rv}_1(c - g)^d$. Finally, in the case of (3), let $Q(x) = P((c - g)x + g)/(c - g)^d$. The roots $(e_i - g)/(c - g)$ of Q are in \mathcal{O} .

Thus $Q \in \mathcal{O}[x]$, $Q_i(0) = P_i(g)(c-g)^{i-d} \in \mathcal{O}$ and $v(Q(1)) = 0$. We have:

$$\begin{aligned}
\mathrm{rv}_1(P(c)) &= \mathrm{rv}_1(c-g)^d \mathrm{res}(Q(1)) \\
&= \mathrm{rv}_1(c-g)^d \left(\sum_i \mathrm{res}(Q_i(0)) \right) \\
&= \mathrm{rv}_1(c-g)^d \left(\bigoplus_i \mathrm{rv}_1(Q_i(0)) \right) \\
&= \bigoplus_{i \leq d} \mathrm{rv}_1(P^{(i)}(g)) \mathrm{rv}_1(c-g)^i \\
&\in \mathbf{RV}_1(\mathrm{dcl}_0(LR\mathrm{rv}_1(c-g))) \\
&\subseteq \mathrm{dcl}_0(R\mathrm{rv}_1(c-g)),
\end{aligned}$$

where the third equality follows from the fact that $v(\sum_i Q_i(0)) = v(Q(1)) = 0 \leq \min_i \{v(Q_i(0))\} \leq v(\sum_i Q_i(0))$. \square

Remark 4.3.2. In mixed characteristic, we will be applying this result to the least equicharacteristic zero coarsening, yielding a computation for rv_∞ and not just rv_1 .

Essentially every computation of leading terms reduces to the above cases by the following lemma.

Lemma 4.3.3. *Let $M \models \mathrm{ACVF}$, $L \leq K = \mathbf{K}(M)$, $c \in K$ and $\mathrm{rv}_\infty(L) \leq R = \mathbf{RV}(\mathrm{dcl}_0(R)) \leq \mathrm{rv}_\infty(K)$. The following are equivalent:*

- (1) $\mathrm{rv}_\infty(L(c)) \subseteq R$;
- (2) For every monic irreducible $P \in \mathbf{K}(\mathrm{dcl}_0(LR))$, $\mathrm{rv}_\infty(P(c)) \in R$.

Moreover if $P \in \mathbf{K}(\mathrm{dcl}_0(LR))[x]$ is irreducible, then its roots are either all inside or outside any $B \in \mathbf{B}^{[<\infty]}(\mathrm{dcl}_0(LR))$.

Proof. By (1), $\mathrm{rv}_\infty(P(c)) \in \mathbf{RV}(\mathrm{dcl}_0(LRc)) \subseteq \mathbf{RV}(\mathrm{dcl}_0(\mathrm{rv}_\infty(L(c))R)) = R$. The converse is a consequence of the fact that rv_∞ is a multiplicative morphism and any polynomial over L is a product of (an element of L and) monic irreducible polynomials over $\mathbf{K}(\mathrm{dcl}_0(LR))$.

As for the moreover statement, let $Q = \prod_{e \in B^v} (x-e) \in \mathbf{K}(\mathrm{dcl}_0(LR))[x]$ where the e range over the roots of P (with multiplicity). If P is irreducible, then $Q = P$ or $Q = 1$. \square

One last important ingredient — also ubiquitous in the development of motivic integration (e.g., in [HK06]) — is the fact that, in characteristic zero, finite sets of points (and of balls in equicharacteristic zero) can be canonically parametrised by \mathbf{RV} . Recall the definition of $b[n]$ and $B[n]$ from Definition 3.2.16.

Lemma 4.3.4. *For every $r \in \mathbb{Z}_{>0}$, there exists $m \in \mathbb{Z}_{>0}$ such that for every characteristic zero valued field L and $B \in \mathbf{B}^{[r]}(L)$ with $|B[m]| = r$, there exists an $\mathcal{L}_0(\ulcorner B \urcorner)$ -definable injection $\nu : B \rightarrow \mathbf{RV}^m$.*

Proof. We proceed by induction on r . If $r = 1$, take $m = 1$ and ν to be constant equal to $1 \in \mathbf{RV}_1$. If $|B| > 1$, we may assume that $|B[m]| = r$

for all m , the Lemma will follow by compactness. Also, assuming that $L \models \text{ACVF}$ is sufficiently saturated and homogeneous, it suffices to find an $\text{Aut}(L/\ulcorner B \urcorner)$ -invariant injection $\nu : B \rightarrow (\mathbf{RV}_\infty)^n$. Indeed, since B is finite some projection to \mathbf{RV}^n is already injective and it must be definable. Finally, let $\gamma := \max\{v(b_1 - b_2) : b_i \in B \text{ distinct}\}$. Since $|B[m]| = r$ for all m , $\gamma < \text{rad}(B) + v(\mathbb{Z})$ and B can be injected in the set of open v_∞ -balls of radius γ/Δ_∞ . So we may assume that the residue characteristic of L is zero. Let B' be the set of closed balls of radius γ around the balls of B . By construction, we have $|B'| < |B| = r$. For every $b' \in B'$, let $B_{b'} := \{b \in B : b \subseteq b'\}$. Note that, by hypothesis, $\text{res}_{b'}(b) \in \mathbf{R}_{b'} = \{\text{maximal open subballs of } b'\}$ uniquely determines b inside B . Let $c_{b'} \in \mathbf{R}_{b'}$ denote the average of the $\text{res}_{b'}(b)$ as b ranges over $B_{b'}$. By induction, we find an $\mathcal{L}_0(\ulcorner B \urcorner)$ -definable injection $\mu : \{c_{b'} : b' \in B'\} \rightarrow \mathbf{RV}^n$. For every $b \in B$, let $\nu(b) := (rv_1(b - c_{b'}), \mu(c_{b'}))$ where $b \subseteq b' \in B'$. Then $\nu : B \rightarrow \mathbf{RV}^{n+1}$ is an $\mathcal{L}_0(\ulcorner B \urcorner)$ -definable injection. \square

Lemma 4.3.5. *For every $r \in \mathbb{Z}_{>0}$ there exists an $m \in \mathbb{Z}_{>0}$ such that for every characteristic zero valued field L and every $B \in \mathbf{B}^{[r]}(L)$ with $|B[m]| = r$, there exists $g \in \mathbf{K}^{[r]}(L)$ with exactly one point inside each ball of $B[m]$.*

Proof. Let us start with a weaker version of the result:

Claim 4.3.6. *For every $b \in \mathbf{B}(\text{acl}_0(L))$, $b(\text{acl}_0(L)) \neq \emptyset$.*

Proof. We may assume that L is algebraically closed. If $v(L) \neq 0$, $L \models \text{ACVF}$ and hence, by model completeness, $b(L) \neq \emptyset$. If $v(L) = 0$, $\text{rad}(b) \in \Gamma(\text{dcl}_0(L)) = \{0\}$. If $0 \in b$, we are done. Otherwise, $v(b) = 0$ and $b \subseteq \mathcal{O}$. So b is open and it is (interdefinable with) a residue element. But $\mathbf{k}(\text{acl}_0(L)) = \text{res}(L)$ and thus $b(L) \neq \emptyset$. \square

Claim 4.3.7. *For every $B \in \mathbf{B}^{[r]}(\text{dcl}_0(L))$, there exists $m \in \mathbb{Z}_{>0}$ and $g \in \mathbf{K}^{[r]}(\text{dcl}_0(L))$ such that, if $|B[m]| = r$, there is exactly one point of g inside each ball of $B[m]$.*

Proof. We may assume that B is irreducible over L — i.e., for any non-empty $\mathcal{L}_0(L)$ -definable $C \subseteq B$, $C = B$. For every $b \in B$, let $d \in b(\text{acl}_0(L))$. Let D be $\mathcal{L}_0(L)$ -definable and irreducible over L containing d and let $g_b \in \text{acl}_0(L)$ be the average of $D \cap b$. Then $g_b \in b[m]$, where $m := |D \cap b|$. Let g be finite $\mathcal{L}_0(L)$ -definable set irreducible over L containing g_b . Since $|B[m]| = r = |B|$, we get $g \cap b[m] = \{g_b\}$. By irreducibility, each ball of $B[m]$ contains exactly one element of g . \square

The lemma follows by compactness. \square

Let $\mathbf{B}_x^{[r]}$ denote the (ind- \mathcal{L}_0 -definable) set of \mathcal{L}_0 -definable maps $F : \mathbf{K}^x \rightarrow \mathbf{B}^{[r]}$ and $\mathbf{B}_x^{[<\infty]}$ denote the (ind- \mathcal{L}_0 -definable) set $\bigcup_r \mathbf{B}_x^{[r]}$. Similarly we denote $\mathbf{K}_x^{[r]}$ the (ind- \mathcal{L}_0 -definable) set of \mathcal{L}_0 -definable maps $F : \mathbf{K}^x \rightarrow \mathbf{K}^{[r]}$ and $\mathbf{K}_x^{[<\infty]}$ the (ind- \mathcal{L}_0 -definable) set $\bigcup_r \mathbf{K}_x^{[r]}$.

Notation. We fix $M \leq N \models T$, $A \subseteq M^{\text{eq}}$, $a \in \mathbf{K}^x(N)$ a potentially infinite tuple and $c \in \mathbf{K}(N)$ a single element. Assume that $p(xy) = \text{tp}_0(ac/M)$ is $\text{Aut}(M/A)$ -invariant and let $q := \text{tp}_0(a/M)$. For every $F, G \in \mathbf{B}_x^{[<\infty]}(M)$, we write $F \leq_q G$ if $q(x) \vdash F^\cup(x) \subseteq G^\cup(x)$. Finally, let $E := \{F \in \mathbf{B}_x^{[<\infty]}(M) : p(x, y) \vdash y \in F^\cup(x)\}$.

In the following Lemmas 4.3.8, 4.3.10 and 4.3.13, we will describe how $\mathbf{RV}(M(ac))$ is generated depending on the shape of E .

Lemma 4.3.8 (Finite sets). *Assume E has a least element $f \in \mathbf{K}_x^{[r]}$ for \leq_q . Then, there exists a pro- $\mathcal{L}_0(M)$ -definable map $\rho : \mathbf{K}^{xy} \rightarrow \mathbf{RV}^n$, whose p -germ is $\text{Aut}(M/A)$ -invariant such that $\text{rv}_\infty(M(ac)) \subseteq \text{dcl}_0(\text{rv}_\infty(M(a))\rho(ac))$.*

Proof. Let $\rho(ac) = \nu_a(c)$, where $\nu_a : f(a) \rightarrow \mathbf{RV}^n$ is the $\mathcal{L}_0(f(a))$ -definable injection of Lemma 4.3.4. By invariance of p , for every $\sigma \in \text{Aut}(M/A)$, $c \in f^\sigma(a) \cap f(a)$. By minimality, we have $f^\sigma(a) = f(a)$ and hence $[f]_q$ — and thus $[\rho]_p$ — is $\text{Aut}(M/A)$ -invariant. Moreover, since $c \in \text{dcl}_0(Ma\rho(ac))$, we have $\text{rv}_\infty(M(ac)) \subseteq \mathbf{RV}(\text{dcl}_0(Ma\rho(ac))) \subseteq \text{dcl}_0(\text{rv}_\infty(Ma)\rho(ac))$. \square

We now assume that $E \cap \mathbf{K}_x^{[<\infty]} = \emptyset$.

Lemma 4.3.9. *There exists a pro- $\mathcal{L}_0(M)$ -definable map $\nu : \mathbf{K}^{xy} \rightarrow \mathbf{RV}^\xi$, with ξ potentially infinite, whose p -germ is $\text{Aut}(M/A)$ -invariant such that, for every $F \in E$, for some $m \in \mathbb{Z}_{>0}$ uniformly bounded in $|F(a)|$, the ball $b \in F(a)[m]$ containing c is $\mathcal{L}_0(Ma\nu(ac))$ -definable, and $\nu(ac) \in \text{acl}_0(Ma)$.*

Proof. For every $r \in \mathbb{Z}_{>0}$, let $m_r \in \mathbb{Z}_{>0}$ be as in Lemma 4.3.4. Let $F \in E \cap \mathbf{B}_x^{[r]}$ be irreducible over q and such that $|F[m_r]| = r$ — if such an F does not exist let $\nu_r(x) = 1$. By irreducibility, for every $G \in E \cap \mathbf{B}_x^{[r]}$ with $G \leq_q F$, every ball in $F(a)$ contains exactly one ball of $G(a)$. In particular, neither $\gamma = \max\{v(b_1 - b_2) : b_i \in F(a) \text{ distinct}\}$ nor $B_r(a)$, the set of open balls of radius $\gamma + v(m_r)$ around balls of $F(a)$, depend on the choice of F . It follows that $[B_r]_q$ is $\text{Aut}(M/A)$ -invariant. By construction, inclusion induces an injection $F(a) \rightarrow B_r(a)$ and that $|B_r(a)| = |B_r(a)[m_r]|$.

As in Lemma 4.3.4, let $\nu_r : B_r(a) \rightarrow \mathbf{RV}^n$ be an $\mathcal{L}_0(B_r(a))$ -definable injection. Let $\nu_r(ac) = \nu_r(b)$ where $c \in b \in B_r(a)$. Note that $\nu_r(ac) \in \text{acl}_0(Ma)$. The element of $F(a)$ containing c is uniquely determined by b , and hence by $\nu_r(ac)$; and $[\nu_r]_p$ is $\text{Aut}(M/A)$ -invariant by construction.

Let us now fix any $F \in E$ that we can assume irreducible. Let $M = \max\{m_s : s \leq |F(a)|\}$. The sequence $|F(a)[M^k]| \geq 1$ is decreasing, bounded by $|F(a)|$ and hence, there exists $k \leq |F(a)|$ such that $|F(a)[M^k][M]| = |F(a)[M^{k+1}]| = |F(a)[M^k]|$. Let $r := |F[M^k]|$. By the previous paragraphs and the choice of M , the ball $b \in F(a)[M^k]$ containing c is $\mathcal{L}_0(Ma\nu_r(ac))$ -definable. It follows that $\nu = (\nu_r)_{r \geq 1}$ has the required properties. \square

We now wish to consider the case where, either E induces a strict intersection in the least equicharacteristic zero coarsening v_∞ (case (1)), or c is generic over Ma in a finite set of open v_∞ -balls (case (2)):

Lemma 4.3.10 (Open and strict balls). *Assume that one of the following holds:*

- (1) *For all $F \in E$, there exists $G \in E$ with $G[m] <_q F$, for any $m \in \mathbb{Z}_{>0}$;*
- (2) *There exists an $r \in \mathbb{Z}_{>0}$ such that for every $F \in E$ and $m \in \mathbb{Z}_{>0}$, there exists an open $G \in E \cap \mathbf{B}_x^{[r]}$ with $G[m] \leq_q F$.*

Then, there is a pro- $\mathcal{L}_0(M)$ -definable map $\rho : \mathbf{K}^{xy} \rightarrow \mathbf{RV}^\xi$ whose p -germ is $\text{Aut}(M/A)$ -invariant and such that $\text{rv}_\infty(M(ac)) \subseteq \text{dcl}_0(\text{rv}_\infty(M(a))\rho(ac))$.

Note that the cases (1) and (2) are not mutually exclusive.

Proof. Let $\nu : \mathbf{K}^{xy} \rightarrow \mathbf{RV}^\xi$ be as in Lemma 4.3.9.

Claim 4.3.11. *For every $F \in E$, the ball $b \in F(a)$ containing c is in $\text{dcl}_0(Ma\nu(ac))$.*

Proof. Assume that there exist $G \in E$ with $G[m] \leq_q F$, for every $m \in \mathbb{Z}_{>0}$. Then, by Lemma 4.3.9 applied to G , the ball $b' \in G(a)[m]$ containing c is $\mathcal{L}_0(Ma\nu(ac))$ -definable. The claim follows since $b' \subseteq b$.

Otherwise, by case (2), we can find a minimal r such that for every $F \in E$ and $m \in \mathbb{Z}_{>0}$, there exists an open $G \in E \cap \mathbf{B}_x^{[r]}$ with $G[m] \leq_q F$. Then for m sufficiently large, depending on r , by Lemma 4.3.9, the ball $b' \in G[m]$ containing c is $\mathcal{L}_0(Ma\nu(ac))$ -definable. The claim follows since $b' \subseteq b$. \square

If there does not exist $g \in \mathbf{K}_x^{[\leq \infty]}(M)$ such that $\emptyset <_q g \leq_q E$, let $\rho(ac) = \nu(ac)$. If such a g exists, we may assume that it is irreducible and, then, the cardinality of the $F \in E$ irreducible over q is bounded by $|g(a)|$. Let $F \in E$ be irreducible over q of maximal cardinality r and let $b(ac) \in F(a)$ contain c . Note that the partition of $g(a)$ induced by F , and in particular $h(ac) := g(a) \cap b(ac)$, does not depend on F . Moreover, since there is some (irreducible) $G \in E \cap \mathbf{B}_x^{[r]}$ with $G[|h(ac)|] \leq_q F$, the average of $h(ac)$ is in $b(ac)$. So, replacing $h(ac)$ by its average, we may assume that $h(ac)$ is a singleton. Then, $h(ac) \in \text{dcl}_0(g(a)b(ac)) \subseteq \text{dcl}_0(Ma\nu(ac))$ by Claim 4.3.11. Let $\rho(ac) = (\nu(ac), \text{rv}_\infty(c - h(ac)))$.

For every $\sigma \in \text{Aut}(M/A)$, applying the previous argument to g and $F^{\sigma^{-1}} \in E$, we have $h(ac) \in b^{\sigma^{-1}}(ac)$, and hence, by invariance of p , $h^\sigma(ac) \in b(ac)$. Note that the smallest (closed) ball containing $h(ac)$ and $h^\sigma(ac)$ is algebraic over Ma and let $D(a)$ be its finite orbit over Ma . We have $D \leq_q E$. If $D[m] \in E$, for some $m \in \mathbb{Z}_{>0}$, then there is $G \in E$ open (irreducible) such that $G[m] \leq_q D[m]$ and hence $D >_q G \in E$. But then either $h(ac) \notin G(ac)$ or $h^\sigma(ac) \notin G$. By invariance of p , we may assume that $h(ac) \notin G(ac)$, contradicting the fact that $h(ac)$ does not depend on the choice of F . Thus $D(a)[m] \notin E$ and so $c \notin D(a)[m]^\cup$. It follows that $\text{rv}_\infty(c - h(ac)) = \text{rv}_\infty(c - h^\sigma(ac))$. We have just proved the $\text{Aut}(M/A)$ -invariance of $[\rho]_p$.

Now, to prove that $\text{rv}_\infty(M(ac)) \subseteq \text{dcl}_0(\text{rv}_\infty(M(a))\rho(ac))$, by Lemma 4.3.3, it suffices to prove that, for every irreducible $P \in \mathbf{K}(\text{dcl}_0(Ma\nu(ac)))[x]$, $\text{rv}_\infty(P(c)) \in \text{dcl}_0(\text{rv}_\infty(M(a))\rho(ac))$. Recall that, by Lemma 4.3.9, $\nu(ac) \in$

$\text{acl}_0(Ma)$. Let $z(a)$ be the finite set, irreducible over $M(a)$, containing the set Z of roots of P . If $z \leq_q E$, then, as above c avoids $d(ac)[m]$, where $d(ac)$ is the smallest closed ball around $Z \cup \{h(ac)\}$. By Lemma 4.3.1.(2), taking into account Remark 4.3.2, $\text{rv}_\infty(P(c)) = \text{rv}_\infty(c - h(ac))^d \in \text{dcl}_0(\rho(ac))$. Otherwise, there is some $F \in E$ such that $Z \cap F(a)^\cup = \emptyset$. Then, by Lemma 4.3.1.(1), $\text{rv}_\infty(P(c)) \in \text{dcl}_0(\text{rv}_\infty(M(a))\nu(ac))$. \square

Remark 4.3.12. There are actually two distinct possible behaviours in Lemma 4.3.10:

- If there does not exist $g \in \mathbf{K}_x^{[<\infty]}$ such that $g \leq_q F$ for every $F \in E$, then $\text{rv}_\infty(M(ac)) \subseteq \text{dcl}_0(\text{rv}_\infty(M(a))\nu(ac)) \subseteq \text{acl}_0(\text{rv}_\infty(M(a)))$;
- If such a g exists, then $\nu(M(ac)) \notin \text{acl}_0(\nu(M(a)))$.

The last remaining case to consider is when c is generic over Ma in some closed v_∞ -ball. For every $B \in \mathbf{B}^{[<\infty]}$, we define $\mathbf{R}_{B,m} := \{b' \subseteq B^\cup : b' \text{ open ball of radius } \text{rad}(B) + v(m)\}$ and $\mathbf{R}_{B,\infty} = \varprojlim_m \mathbf{R}_{B,m}$. For every $x \in B^\cup$, let $\text{res}_{B,m}(x)$ denote the unique element of $\mathbf{R}_{B,m}$ containing x and $\text{res}_{B,\infty} : B^\cup \rightarrow \mathbf{R}_{B,\infty}$ be the induced map.

Lemma 4.3.13 (Closed balls). *Assume that there exists an $F \in E$ such that for every $g \in \mathbf{K}_x^{[<\infty]}(M)$ with $g \leq_q F$, $c \notin \text{res}_{F(a),\infty}(g(a))^\cup$. Let $b \in F(a)$ contain c , $\xi \in \mathbf{R}\mathbf{V}^n$ such that $b \in \text{dcl}_0(Ma\xi)$ and $G \in \text{res}_{b,\infty}(\text{dcl}_0(Ma\xi))$. Then $\text{rv}_\infty(M(ac)) \subseteq \text{dcl}_0(\text{rv}_\infty(M(a))\xi\text{rv}_\infty(\text{res}_{F(a),\infty}(c) - G(a\xi)))$.*

In later applications of this lemma, we will take $\xi = \nu(ac)$ as given by Lemma 4.3.9.

Proof. Note that for any $m \in \mathbb{Z}_{>0}$, the hypothesis on F remains true of $F[m]$. So, replacing F by some $F[m]$, with $|F[m](a)|$ minimal, we may assume that $|F[m](a)|$ is constant. By Lemma 4.3.5, we can now find $f \in \mathbf{K}_x^{[<\infty]}(M)$ such that $f(a)$ has exactly one point in every ball of $F(a)$. By hypothesis, $c \notin \text{res}_{F(a),\infty}(f(a))$. Let $h \in \text{dcl}_0(Ma\xi)$ denote the unique element of $f(a) \cap b$. Since $\text{rad}(F(a))/\Delta_\infty = v_\infty(c - h) = v_\infty(c - G(a, \xi)) \leq v_\infty(G(a, \xi) - h)$, we have $\text{rv}_\infty(c - h) = \text{rv}_\infty(\text{res}_{F(a),\infty}(c) - G(a\xi)) \oplus \text{rv}_\infty(G(a\xi) - h)$. So it suffices to prove that $\text{rv}_\infty(M(ac)) \subseteq \text{dcl}_0(\text{rv}_\infty(M(a))\xi\text{rv}_\infty(c - h))$.

By Lemma 4.3.3, it further suffices to prove that, for every irreducible $P \in \mathbf{K}(\text{dcl}_0(Ma\xi))[x]$, $\text{rv}_\infty(P(c)) \in \text{dcl}_0(\text{rv}_\infty(M(a))\xi\text{rv}_\infty(c - h))$. If, for every $m \in \mathbb{Z}_{>0}$, no root of P is in $b[m]$, then, by Lemma 4.3.1.(1), $\text{rv}_\infty(P(c)) \in \text{dcl}_0(\text{rv}_\infty(M(a))\xi)$. Otherwise, let $m \in \mathbb{Z}_{>0}$ be such that every root of P is in $b[m]$. Since $\mathbf{K}(\text{acl}_0(Ma\xi)) \subseteq \text{acl}_0(Ma)$, let $z(a)$ be finite irreducible over $M(a)$ containing the roots of P . By hypothesis, $c \notin \text{res}_{F(a)[m]}(z(a))$. By Lemma 4.3.1.(3), $\text{rv}_\infty(P(c)) \in \text{dcl}_0(\text{rv}_\infty(M(a))\text{rv}_\infty(c - h))$. \square

Notation. Let $\widehat{A} \subseteq \mathbf{K}(M)$ contain a realisation of every $\mathcal{L}(A)$ -type and assume that M is sufficiently saturated and homogeneous.

We can now wrap up the relative arity one case:

Proposition 4.3.14. *There exists a pro- $\mathcal{L}_0(\widehat{A})$ -definable map $\rho : \mathbf{K}^{xy} \rightarrow \mathbf{RV}^\xi$ such that $\text{rv}_\infty(M(ac)) \subseteq \text{dcl}_0(\text{rv}_\infty(M(a))\rho(ac))$.*

Proof. Note first that any $\text{Aut}(M/A)$ -invariant p -germ of $\mathcal{L}_0(M)$ -definable functions is represented by an $\mathcal{L}_0(\widehat{A})$ -definable function — it suffices to consider a realisation in \widehat{A} of the type of the parameters over A .

If $E \cap \mathbf{K}_x^{[\lt \infty]} \neq \emptyset$, we apply Lemma 4.3.8. So let us assume that $E \cap \mathbf{K}_x^{[\lt \infty]} = \emptyset$. If for every $F \in E$, there exists $G \in E$ with $G[m] \leq_q F$, for every $m \in \mathbb{Z}_{>0}$, we are in case (1) of Lemma 4.3.10 and we can conclude. So we may assume that there exists F such that for every $G \in E$, $F \leq_q G[m]$, for some $m \in \mathbb{Z}_{>0}$. If there exists $g \in \mathbf{K}_x^{[\lt \infty]}$ with $g \leq_q F$ and $c \in \text{res}_{F(a), \infty}(g(a))$, then, for all $H \in E$ and $n \in \mathbb{Z}_{>0}$, $H \leq_q F[n]$, for some $m \in \mathbb{Z}_{>0}$. Let $G(a) := \text{res}_{F(a), mn}(g(a))$, then $c \in G[n] \leq_q H$ and hypothesis (2) of Lemma 4.3.10 holds with $r = |g|$. So we may assume that no such g exist, *i.e.*, the hypotheses of Lemma 4.3.13 hold. As previously, we may assume that $|F[m]|$ is constant. Let ν be as in Lemma 4.3.9; we may assume that ν is $\mathcal{L}_0(\widehat{A})$ -definable. Let $b(ac) \in F(a)$ contain c .

Claim 4.3.15. *There exists $G(ac) \in \text{res}_{b(a)[m], \infty}(\text{dcl}_0(\widehat{A}a\nu(ac)))$, for some $m \in \mathbb{Z}_{>0}$.*

Proof. By construction of \widehat{A} , there exists $\sigma \in \text{Aut}(M/A)$ such that F^σ is $\mathcal{L}_0(\widehat{A})$ -definable. By $\text{Aut}(M/A)$ -invariance of p , we have $F^\sigma \in E$ and hence $F^\sigma \leq_q F[m]$, for some $m \in \mathbb{Z}_{>0}$. So, up to replacing F by F^σ , we may assume F is $\mathcal{L}_0(\widehat{A})$ -definable. By Lemma 4.3.5, and replacing F by some $F[m]$, we find $g \in \mathbf{K}_x^{[\lt \infty]}(\widehat{A})$ with exactly one element in each ball of F . It then suffices to consider the only element of $\mathbf{k}_{F(a), \infty}(g(a))$ contained in $b(ac)$. \square

By Lemma 4.3.13, $\text{rv}_\infty(M(ac)) \subseteq \text{dcl}_0(\text{rv}_\infty(M(a))\nu(ac)\text{rv}_\infty(\text{res}_{F(a), \infty}(c) - G(ac))) \subseteq \text{dcl}_0(\text{rv}_\infty(Ma)\widehat{A}ac)$. \square

Corollary 4.3.16. *There exists a pro- $\mathcal{L}_0(\widehat{A})$ -definable map $\rho : \mathbf{K}^x \rightarrow \mathbf{RV}^\xi$, such that $\text{rv}_\infty(M(a)) \subseteq \text{dcl}_0(\text{rv}_\infty(M)\rho(a))$.*

Proof. We proceed by induction on an enumeration of a . The induction step is Proposition 4.3.14 and the limit case is trivial. \square

Corollary 4.3.17. *The type $\text{tp}(a/M)$ is $\text{Aut}(M/\widehat{A}\mathbf{RV})$ -invariant.*

Proof. It follows from Corollary 4.3.16 and Lemma 4.2.5. \square

4.4. Invariant resolutions.

Notation. Let $M \preceq N \models T$ both be sufficiently saturated and homogeneous and $A \subseteq \mathcal{G}(M)$.

By transitivity, there remains to build a sufficiently saturated model containing A whose type is invariant.

Lemma 4.4.1. *Assume that $A \subseteq \mathbf{K}(M)$ and let $R \subseteq \mathbf{RV}(M)$. There exists $C \subseteq \mathbf{K}(N)$ and a pro- $\mathcal{L}_0(M)$ -definable map $\rho : \mathbf{K}^{|C|} \rightarrow \mathbf{RV}^\xi$ such that $R \subseteq \text{rv}_\infty(A(C)) \subseteq \text{dcl}_0(AR)$, $q := \text{tp}_0(C/M)$ and $[\rho]_q$ are $\text{Aut}(M/AR)$ -invariant and $\text{rv}_\infty(M(C)) \subseteq \text{dcl}_0(\text{rv}_\infty(M)\rho(C))$.*

Proof. We proceed by induction on an enumeration of R . Assume the property holds of R for some C and ρ and pick any $\zeta \in \mathbf{RV}_\infty(M)$. If $\zeta \in \text{rv}_\infty(\text{acl}_0(AC))$, let $c \in \mathbf{K}(\text{acl}_0(AC))$ be such that $\text{rv}_\infty(c) = \zeta$. Let D be a minimal finite $\mathcal{L}_0(AC\zeta)$ -definable set containing c . Replacing c by the average of D , we may assume that $c \in \text{dcl}_0(AC\zeta) \subseteq \text{dcl}_0(MC) \subseteq N$. We have $R\zeta \subseteq \text{rv}_\infty(A(Cc)) \subseteq \mathbf{RV}(\text{dcl}_0(AC\zeta)) \subseteq \text{dcl}_0(\text{rv}_\infty(A(C))\zeta) \subseteq \text{dcl}_0(AR\zeta)$, $\text{tp}_0(Cc/M)$ is $\text{Aut}(M/AR\zeta)$ -invariant and, since $c \in \text{dcl}_0(MC) = M(C)^{\text{h}}$, $\text{rv}_\infty(M(Cc)) = \text{rv}_\infty(M(C)) \subseteq \text{dcl}_0(\text{rv}_\infty(M)\rho(C))$.

If $\zeta \notin \text{rv}_\infty(\text{acl}_0(AC))$, let $c \in N$ be generic in $\text{rv}_\infty^{-1}(\zeta)$ over M . Then $p := \text{tp}_0(Cc/M)$ is $\text{Aut}(M/AR\zeta)$ -invariant. By Lemma 4.3.10, we find a pro- $\mathcal{L}_0(M)$ -definable map $\rho' : \mathbf{K}^{|C|+1} \rightarrow \mathbf{RV}_\infty^\xi$ such that $\text{rv}_\infty(M(Cc)) \subseteq \text{dcl}_0(\text{rv}_\infty(M(C))\rho'(Cc)) \subseteq \text{dcl}_0(\text{rv}_\infty(M)\rho(C)\rho'(Cc))$ and whose p -germ is $\text{Aut}(M/AR\zeta)$ -invariant. Moreover, no root of any $P \in \mathbf{K}(\text{dcl}(AC))[x]$ is in $\text{rv}_\infty^{-1}(\zeta)$. For any $g \in \text{rv}_\infty^{-1}(\zeta)$, by Lemma 4.3.1.(1), $\text{rv}_\infty(P(c)) = \text{rv}_\infty(P(g))$ does not depend on g and is thus in $\mathbf{RV}(\text{dcl}_0(AC\zeta)) \subseteq \text{dcl}_0(\text{rv}_\infty(AC)\zeta) \subseteq \text{dcl}_0(AR\zeta)$. By Lemma 4.3.3, $\text{rv}_\infty(A(Cc)) \subseteq \text{dcl}_0(AR\zeta)$. \square

Corollary 4.4.2. *Assume that $A \subseteq \mathbf{K}(M)$. There exists $A \subseteq C \preceq N$ and a pro- $\mathcal{L}_0(M)$ -definable map $\rho : \mathbf{K}^{|C|} \rightarrow \mathbf{RV}^\xi$, such that $p := \text{tp}_0(C/M)$ is $\text{Aut}(M/AR\mathbf{V}(M))$ -invariant, $[\rho]_p$ is $\text{Aut}(M/AR\mathbf{V}(M))$ -invariant and $\text{rv}_\infty(MC) \subseteq \text{dcl}_0(\text{rv}_\infty(M)\rho(C))$.*

In particular, by Lemma 4.2.5, $\text{tp}(C/M)$ is $\text{Aut}(M/AR\mathbf{V})$ -invariant.

Proof. Let $A \subseteq M_1 \preceq M$, with M_1 small. Applying Lemma 4.4.1 to $\text{rv}_\infty(M_1)$, we find $C \subseteq N$ and ρ such that $\text{rv}_\infty(M_1) \subseteq \text{rv}_\infty(A(C)) \subseteq \text{dcl}_0(\text{Arv}_\infty(M_1)) \cap \text{rv}_\infty(N) \subseteq \text{rv}_\infty(M_1)$, $p := \text{tp}_0(C/M)$ and $[\rho]_p$ are $\text{Aut}(M/\text{Arv}_\infty(M_1))$ -invariant and $\text{rv}_\infty(M(C)) \subseteq \text{dcl}_0(\text{rv}_\infty(M)\rho(C))$. By replacing C with $\mathbf{K}(\text{dcl}_0(AC))$, we may assume $A \subseteq C = C^{\text{h}}$. Since $\text{rv}_\infty(C) = \text{rv}_\infty(A(C)) = \text{rv}_\infty(M_1) \preceq \text{rv}_\infty(N)$ and C is a characteristic zero henselian field, it follows from Fact 2.2.6 that $C \preceq N$. \square

Corollary 4.4.3. *Assume that $A \subseteq \mathbf{K}(M)$. There exists $\widehat{A} \preceq N$ containing a realisation of every $\mathcal{L}(A)$ -type such that $\text{tp}(\widehat{A}/M)$ is $\text{Aut}(M/AR\mathbf{V})$ -invariant.*

Proof. Corollary 4.4.2 implies that, for every $\mathcal{L}(A)$ -definable set X , there exists an $\text{Aut}(M/AR\mathbf{V})$ -invariant type concentrating on X — since the C from Corollary 4.4.2 is a model containing A . Since the set of $\text{Aut}(M/AR\mathbf{V})$ -invariant types is closed, it follows by compactness that any $\mathcal{L}(A)$ -type has an $\text{Aut}(M/AR\mathbf{V})$ -invariant extension. The corollary follows by the standard construction relying on transitivity, Lemma 4.2.4 — for the limit steps,

note that $\text{Aut}(M/\text{ARV})$ -invariance is finitary: $\text{tp}(c/M)$ is $\text{Aut}(M/\text{ARV})$ -invariant if and only if for every finite $c_0 \subseteq c$, $\text{tp}(c_0/M)$ is $\text{Aut}(M/\text{ARV})$ -invariant. \square

Recall that, by Convention 2.1, $\mathbf{Lin}_A(M)$ denotes the set of cosets $c + \ell \mathbf{ms}$ where $s \in \mathbf{S}(\text{dcl}_0(A))$ has a basis in M and $c \in s(M)$.

Lemma 4.4.4. *Assuming:*

(\mathbf{I}_k) *The residue field \mathbf{k} is infinite in models of T ,*

there exists $C \subseteq \mathbf{K}(N)$ and an $\mathcal{L}_0(A)$ -definable map $\rho : \mathbf{K}^{|C|} \rightarrow \mathbf{Lin}_A^\xi$ such that, for all $n \in \mathbb{Z}_{>0}$, $\mathbf{S}_n(A) \subseteq s_n(C)$, $\text{tp}_0(C/M)$ is $\mathcal{L}_0(A)$ -definable and $\text{rv}_\infty(M(C)) \subseteq \text{dcl}_0(\text{rv}_\infty(M)\mathbf{Lin}_A(M)\rho(C))$.

In mixed characteristic, we may further assume that $\bigcup_{n>0} \mathbf{T}_n(A) \subseteq \text{dcl}_0(C)$.

Proof. Fix $s \in \mathbf{S}_n(A)$ and let $\beta \in \text{GL}_n(M)$ be a basis of s . Then, any $\alpha \vDash \beta \cdot (\eta_{\mathcal{O}|_M})^{\otimes n^2}$ — which is realised in N by (\mathbf{I}_k) — where $\eta_{\mathcal{O}}$ is the generic (quantifier free) type of \mathcal{O} , is a basis of s . Note that $\text{tp}_0(\alpha/M) = \beta \cdot \eta_{\mathcal{O}}^{\otimes n^2}$ only depends on s and is indeed $\mathcal{L}_0(A)$ -definable. Let $\bar{\alpha}$, respectively $\bar{\beta}$, be the basis of $s/\mathfrak{m}_\infty s$ — seen as a pro-definable set — induced by α , respectively β . The matrix of \mathbf{k}_∞ -coefficients of α in the basis β is $\text{res}_\infty(\beta^{-1} \cdot \alpha)$ where $\beta^{-1} \cdot \alpha \vDash (\eta_{\mathcal{O}|_M})^{\otimes n^2}$. It follows that $\text{rv}_\infty(M(\alpha)) = \text{rv}_\infty(M(\beta^{-1} \cdot \alpha)) \subseteq \text{dcl}_0(\mathbf{RV}(M)\text{res}_\infty(\beta^{-1} \cdot \alpha)) \subseteq \text{dcl}_0(\mathbf{RV}(M)\bar{\beta}\bar{\alpha})$. The first part of the statement follows by iterating the above construction independently for every $s \in \mathbf{S}_n(A)$.

Let us now assume that we are in mixed characteristic. For every $c \vDash \eta_{\mathfrak{m}|_M}$, $\text{rv}_\infty(M(Cc)) \subseteq \text{dcl}_0(\text{rv}_\infty(M(C))\text{res}_\infty(c))$. This also holds for all maximal open subballs of \mathcal{O} . So, enlarging C further, we may also assume that $\mathbf{k}(\text{dcl}_0(AC)) \cap M \subseteq \text{res}(C)$. Then, for every $e \in \mathbf{T}_n(A)$, $s = \tau_n(e) \in \mathbf{S}(A)$ has a basis in C and every coordinate of e in that basis is the residue of an element of C . It then follows that $e \in t_n(C)$. \square

Lemma 4.4.5. *In equicharacteristic zero, assuming that, for all $n \in \mathbb{Z}_{>0}$, $\tau_n(\mathbf{T}_n(A)) \subseteq s_n(\mathbf{K}(A))$. Then there exists $C \subseteq \mathbf{K}(N)$ and an $\mathcal{L}_0(M)$ -definable map $\rho : \mathbf{K}^{|C|} \rightarrow \mathbf{RV}^\xi$ such that $A \subseteq \text{dcl}_0(C)$, $q := \text{tp}_0(C/M)$ is $\text{Aut}(M/A)$ -invariant, $[\rho]_q$ is $\text{Aut}(M/A)$ -invariant and $\text{rv}_\infty(M(C)) \subseteq \text{dcl}_0(\text{rv}_\infty(M)\rho(C))$.*

Proof. For every $e \in \mathbf{T}_n(A)$, by hypothesis, $s := \tau(e)$ has a basis in $\mathbf{K}(A)$. It follows that $s/\mathfrak{m}s$ also has a basis of $\text{dcl}_0(A)$ -points and hence is $\mathcal{L}_0(A)$ -definably isomorphic to \mathbf{k}^n . By Lemma 4.4.1 applied to $R := \mathbf{k}(\text{dcl}_0(A)) \cap M \subseteq \text{dcl}_0(A)$, we find $\mathbf{K}(A) \subseteq C \subseteq \mathbf{K}(N)$ such that $\mathbf{k}(\text{dcl}_0(A)) \cap M \subseteq \text{res}(C)$, $q := \text{tp}_0(C/M)$ is $\text{Aut}(M/A)$ -invariant, $[\rho]_q$ is $\text{Aut}(M/A)$ -invariant and $\text{rv}_\infty(M(C)) \subseteq \text{dcl}_0(\text{rv}_\infty(M)\rho(C))$. Then, we have $A = \mathbf{K}(A) \cup \bigcup_n \mathbf{S}_n(A) \cup \bigcup_n \mathbf{T}_n(A) \subseteq \text{dcl}_0(C)$. \square

Corollary 4.4.6. *Assume that (\mathbf{I}_k) holds. Then there exists $C \subseteq \mathbf{K}(N)$ such that $\text{tp}_0(C/M)$ has $\text{Aut}(M/\text{ARV}(M)\mathbf{Lin}_A(M))$ -invariant \mathbf{RV} -germs and $A \subseteq \text{dcl}_0(C)$.*

Proof. In mixed characteristic, this follows immediately from Lemmas 4.2.5 and 4.4.4. In residue characteristic zero, it follows from Lemmas 4.4.4 and 4.4.5 and transitivity, cf. Lemma 4.2.4. \square

Proof of Theorem 4.1.1. By Corollary 4.4.6, we find $C \subseteq \mathbf{K}(N)$ such that $A \subseteq \text{dcl}_0(C)$ and $\text{tp}(C/M)$ has $\text{Aut}(M/\text{ARV}(M)\mathbf{Lin}_A(M))$ -invariant \mathbf{RV} -germs. Let $M \preceq M_1 \preceq N$ be sufficiently saturated and homogeneous and contain C . By Corollary 4.4.3, we find $\widehat{C} \preceq N$ containing a realisation of every $\mathcal{L}(C)$ -type such that $\text{tp}(\widehat{C}/M_1)$ is $\text{Aut}(M_1/\text{CRV})$ -invariant. Let $M_1 \preceq M_2 \preceq N$ be sufficiently saturated and homogeneous and contain \widehat{C} . Let $p := \text{tp}_0(a/M)$, which is $\text{Aut}(M/A)$ -invariant by assumption.

Claim 4.4.7. $\text{tp}(a/M) \cup p|_{M_2}$ is consistent.

Proof. Let $\varphi(x, m)$ be some $\mathcal{L}(M)$ -formula such that $N \models \varphi(a, m)$ and let $\psi(x, d) \in p|_{M_2}$. Let $\sigma \in \text{Aut}(N/Amd)$ be such that $\sigma(d) \in M$. Then $N \models \psi(a, \sigma(d))$ and hence $\sigma^{-1}(a) \models \psi(x, d) \wedge \varphi(x, m)$. \square

Let $a' \models \text{tp}(a/M) \cup p|_{M_2}$. Then $\text{tp}_0(a'/M_2) = p|_{M_2}$ is $\text{Aut}(M_2/C)$ -invariant and thus, by Corollary 4.3.17, $\text{tp}(a'/M_2)$ is $\text{Aut}(M_2/\widehat{C}\mathbf{RV})$ -invariant. Since $\text{tp}(\widehat{C}/M_1)$ is $\text{Aut}(M_1/\text{CRV})$ -invariant, by transitivity, cf. Lemma 4.2.4, $\text{tp}(a'/M_1)$ is $\text{Aut}(M_1/\text{CRV})$ -invariant. By transitivity, since $\text{tp}(C/M)$ has $\text{Aut}(M/\text{ARV}(M)\mathbf{Lin}_A(M))$ -invariant \mathbf{RV} -germs, $\text{tp}(a/M) = \text{tp}(a'/M)$ is $\text{Aut}(M/\text{ARV}(M)\mathbf{Lin}_A(M))$ -invariant. \square

Let us conclude this section by relating Theorem 4.1.1 to imaginaries:

Proposition 4.4.8. *Let $T \supseteq \text{Hen}_0$ be an \mathcal{L} -theory, such that:*

- (D) *For every strict pro- $\mathcal{L}(A)$ -definable $X \subseteq \mathbf{K}^x$, with $A = \text{acl}^{\text{eq}}(A) \subseteq M^{\text{eq}} \models T^{\text{eq}}$, there exist an $\text{Aut}(M/\mathcal{G}(A))$ -invariant $p \in S_x^0(M)$ consistent with X ;*
- (Q \mathbf{K}) *For every tuple $a \in \mathbf{K}(M)$, with $M \models T$, $\text{tp}_1(f(a)) \vdash \text{tp}(a)$, where $f : \mathbf{K} \rightarrow \mathbf{K}^x$ is pro- \mathcal{L} -definable and $\mathcal{L}_0 \subseteq \mathcal{L}_1 \subseteq \mathcal{L}$ such that \mathcal{L}_1 is an \mathbf{RV} -enrichment of \mathcal{L}_0 ;*
- (I \mathbf{k}) *The residue field \mathbf{k} is infinite;*
- (SE) *\mathbf{RV} and $\mathbf{R} = \bigcup_{\ell} \mathcal{O}/\ell\mathbf{m}$ are stably embedded.*

Let $M \models T$, $e \in M^{\text{eq}}$ and $A = \text{acl}^{\text{eq}}(e)$. Then

$$e \in \text{dcl}^{\text{eq}}(\mathbf{K}(A) \cup (\mathbf{RV} \cup \mathbf{Lin}_{\mathcal{G}(A)})^{\text{eq}}(A)).$$

Proof. Let $M \models T$ be saturated and sufficiently large, $e \in M^{\text{eq}}$ and $A = \text{acl}^{\text{eq}}(e)$. Then $e = g(a)$ for some \mathcal{L} -definable map g and tuple $a \in \mathbf{K}(M)$. Let $Y = g^{-1}(e)$ and $X = f(Y)$, which is a strict pro- $\mathcal{L}(A)$ -definable set. By (D), there exists an $\text{Aut}(M/\mathcal{G}(A))$ -invariant $p \in S_x^0(M)$ consistent with X . We may assume that $f(a) \models p$. By Theorem 4.1.1, $\text{tp}_1(f(a)/M)$ is $\text{Aut}(M/\mathcal{G}(A)\mathbf{RV}(M)\mathbf{Lin}_{\mathcal{G}(A)}(M))$ -invariant. By (Q \mathbf{K}), $\text{tp}(a/M)$ — and hence $e \in M^{\text{eq}}$ — is also $\text{Aut}(M/\mathcal{G}(A)\mathbf{RV}(M)\mathbf{Lin}_{\mathcal{G}(A)}(M))$ -invariant. Since $\mathbf{Lin}_{\mathcal{G}(A)}$ is a collection of free $\mathcal{O}/\ell\mathbf{m}$ -modules, $\mathbf{Lin}_{\mathcal{G}(A)}$ and, in fact $\mathbf{Lin}_{\mathcal{G}(A)} \cup \mathbf{RV}$, is stably embedded. By Lemma 4.2.3, $e \in \text{dcl}^{\text{eq}}(\mathcal{G}(A) \cup$

$\mathbf{RV}(M) \cup \mathbf{Lin}_{\mathcal{G}(A)}(M)$ — i.e., $e = h(c)$ for some $\mathcal{L}(\mathcal{G}(A))$ -definable map h and tuple $c \in \mathbf{RV}^m(M) \times \mathbf{Lin}_{\mathcal{G}(A)}^n(M)$. Let $Z = h^{-1}(e)$. Then $\ulcorner Z \urcorner \in (\mathbf{RV} \cup \mathbf{Lin}_{\mathcal{G}(A)})^{\text{eq}}(A)$ and

$$\begin{aligned} e \in \text{dcl}^{\text{eq}}(\mathcal{G}(A) \ulcorner Z \urcorner) &\subseteq \text{dcl}^{\text{eq}}(\mathcal{G}(A) \cup (\mathbf{RV} \cup \mathbf{Lin}_{\mathcal{G}(A)})^{\text{eq}}(A)) \\ &\subseteq \text{dcl}^{\text{eq}}(\mathbf{K}(A) \cup (\mathbf{RV} \cup \mathbf{Lin}_{\mathcal{G}(A)})^{\text{eq}}(A)), \end{aligned}$$

since $\mathcal{G}(A) \setminus \mathbf{K}(A) \subseteq \mathbf{Lin}_{\mathcal{G}(A)}^{\text{eq}}(A)$. \square

5. IMAGINARIES IN SHORT EXACT SEQUENCES

In this section we will establish results which yield a relative understanding of imaginaries in certain pure short exact sequences of modules.

5.1. The core case. We start with a well known lemma. We include a proof for convenience.

Lemma 5.1.1. *Let D and C be stably embedded (ind-)definable sets in $D \cup C$, such that $D \perp C$.*

- (1) *Assume that both D and C (considered with the full induced structures) weakly eliminate imaginaries. Then $D \cup C$ weakly eliminates imaginaries.*
- (2) *Assume that both D and C eliminate imaginaries and that in C one has $\text{dcl} = \text{acl}$. Then $D \cup C$ eliminates imaginaries.*

Proof. (1) Let $X \subseteq D^m \times C^n$ be a definable subset. Since $D \perp C$, the equivalence relation \sim on C^n , given by $c \sim c' :\Leftrightarrow X_c = X_{c'}$, has finitely many equivalence classes $Z_1 = c_1 / \sim, \dots, Z_k = c_k / \sim$. As \sim is $\ulcorner X \urcorner$ -definable and C is stably embedded, the Z_i are all $C^{\text{eq}}(\text{acl}^{\text{eq}}(\ulcorner X \urcorner))$ -definable.

For $i = 1, \dots, k$, set $Y_i = X_{c_i} \subseteq D^m$, which is $D^{\text{eq}}(\text{acl}^{\text{eq}}(\ulcorner X \urcorner))$ -definable since D is stably embedded. As D and C weakly eliminate imaginaries, there are finite tuples $d \in D(\text{acl}^{\text{eq}}(\ulcorner X \urcorner))$ and $c \in C(\text{acl}^{\text{eq}}(\ulcorner X \urcorner))$ such that the Y_i are all d -definable and the Z_i are all c -definable. Thus $X = \bigcup_{i=1}^k (Y_i \times Z_i)$ is dc -definable, so in particular $D(\text{acl}^{\text{eq}}(\ulcorner X \urcorner))C(\text{acl}^{\text{eq}}(\ulcorner X \urcorner))$ -definable.

(2) The assumptions on C yield $C^{\text{eq}}(\text{acl}^{\text{eq}}(\ulcorner X \urcorner)) \subseteq \text{dcl}^{\text{eq}}(C(\ulcorner X \urcorner))$, and so the sets Z_1, \dots, Z_k are c -definable for some $c \in C(\ulcorner X \urcorner)$. In particular the Y_i are then all $\ulcorner X \urcorner$ -definable, thus $D^{\text{eq}}(\ulcorner X \urcorner)$ -definable. By elimination of imaginaries in D , we find $d \in D(\ulcorner X \urcorner)$ such that all Y_i are d -definable. We now finish as in (1). \square

Fact 5.1.2. *Let G be a group, and let H_1, \dots, H_N be subgroups of G . Then the left cosets of the H_i form a pre-basis of closed sets for a noetherian topology on G . Moreover, setting $H_I := \bigcap_{i \in I} H_i$ for $I \subseteq \{1, \dots, N\}$, the irreducible closed sets for this topology are precisely the left cosets of those H_I with the property that any proper subgroup of H_I of the form H_J is of infinite index in H_I .*

Proof. This is an easy consequence of Neumann's Lemma. \square

Let R be an integral domain, $\mathcal{L} \supseteq \mathcal{L}_{R\text{-mod}}$ and M an \mathcal{L} -expansion of an infinite torsion free R -module. Let $Z \subseteq M^n$ be an $\mathcal{L}(M)$ -definable set. We set $\dim_R(Z) := \max\{\dim_R(c/M) : c \in Z(N)\}$, where $N \succcurlyeq M$ is sufficiently saturated and $\dim_R(a/B)$ denotes the $Q(R)$ -linear dimension of a over B , for $Q(R)$ the field of fractions of R .

Lemma 5.1.3. *In the above situation, assume $\dim_R(Z) \leq r$. Then there are definable sets $C_1, \dots, C_s \subseteq M^n$ with the following properties:*

- (1) $\bigcup_{i=1}^s C_i$ is $\ulcorner Z \urcorner$ -definable.
- (2) Each C_i is $\text{acl}^{\text{eq}}(\ulcorner Z \urcorner)$ -definable.
- (3) Each C_i is of the form $\gamma_i + H_i$, where H_i is a definable R -submodule of M^n given by a condition $L_i x' = m x''$, for some splitting of the variables $x = x' x''$ with $|x'| = r$ and $|x''| = n - r$, matrix $L_i \in R^{(n-r) \times r}$ and $m \in R \setminus \{0\}$. (In particular, $\dim_R(C_i) = r$ for all i .)
- (4) $Z \subseteq \bigcup_{i=1}^s C_i$

Moreover, if \mathcal{Z} is a definable family such that $\dim_R(\mathcal{Z}_b) \leq r$ for all b , there are finitely many R -submodules H_1, \dots, H_N as in the statement such that for any b the C_i may be chosen among the cosets of the H_k .

Proof. By compactness there are sets C_1, \dots, C_N with $C_i = \delta_i + H_i$ as in (3) such that $Z \subseteq \bigcup_{i=1}^N C_i$. Note that all H_i are \emptyset -definable subgroups.

By Fact 5.1.2, the cosets of the H_i form a pre-basis of closed sets for a noetherian topology on M^n . In particular, in this topology there exists a smallest closed subset W of M^n containing Z , and this W is clearly $\ulcorner Z \urcorner$ -definable. The (finitely many) irreducible components of W are then all $\text{acl}^{\text{eq}}(\ulcorner Z \urcorner)$ -definable. If W_j is such an irreducible component, it is of the form $W_j = \gamma_j + \bigcap_{i \in I_j} H_i$ (where $I_j \neq \emptyset$ in case $r < n$, since $Z \subseteq \bigcup_{i=1}^N \delta_i + H_i$ by assumption). As the H_i are \emptyset -definable, it is easy to see that if we replace each component $W_j = \gamma_j + \bigcap_{i \in I_j} H_i$ by $W'_j = \bigcup_{i \in I_j} \gamma_j + H_i$, then the union of all W'_j is $\ulcorner Z \urcorner$ -definable and each coset $\gamma_j + H_i$ occurring in this union is $\text{acl}^{\text{eq}}(\ulcorner Z \urcorner)$ -definable.

The moreover part follows by compactness. \square

Theorem 5.1.4. *Let R be an integral domain and M be*

$$0 \rightarrow \mathbf{A} \rightarrow \mathbf{B} \rightarrow \mathbf{C} \rightarrow 0$$

a short exact sequence of R -modules, in an \mathbf{A} - \mathbf{C} -enrichment \mathcal{L} of the pure (in the sense of model theory) three sorted sequence of R -modules. Assume the following properties hold:

- (1) \mathbf{A} is a pure submodule of \mathbf{B} (in the sense of module theory),
- (2) \mathbf{C} is torsion free,
- (3) For any $l \in R \setminus \{0\}$, the quotient $\mathbf{C}/l\mathbf{C}$ is finite and the preimage in \mathbf{B} of any coset $c + l\mathbf{C}$ contains an element which is algebraic over \emptyset .

Let $e \in M^{\text{eq}}$. Then, setting $E := \text{acl}^{\text{eq}}(e)$ and $\Delta := \mathbf{C}(E)$, we have

$$e \in \text{dcl}^{\text{eq}}(\mathbf{C}^{\text{eq}}(E)\mathbf{B}_{\Delta}^{\text{eq}}(E)),$$

where \mathbf{B}_{Δ} denotes the union of all fibers \mathbf{B}_{δ} with $\delta \in \Delta$.

We will prove a slight generalization of Theorem 5.1.4, namely the following Theorem 5.1.5.

Theorem 5.1.5. *Let \tilde{R} be a ring and $R = \tilde{R}/I$ an integral domain, with I a finitely generated ideal. Let M be*

$$0 \rightarrow \tilde{\mathbf{A}} \rightarrow \tilde{\mathbf{B}} \rightarrow \tilde{\mathbf{C}} \rightarrow 0$$

a short exact sequence of \tilde{R} -modules, in an $\tilde{\mathbf{A}}\text{-}\tilde{\mathbf{C}}$ -enrichment \mathcal{L} of the pure (in the sense of model theory) three sorted sequence of \tilde{R} -modules. Let $\mathbf{A} = \{a \in \tilde{\mathbf{A}} : Ia = (0)\}$, and let \mathbf{B} and \mathbf{C} be the \tilde{R} -submodules of $\tilde{\mathbf{B}}$ and $\tilde{\mathbf{C}}$ defined similarly. Consider $\mathbf{A}, \mathbf{B}, \mathbf{C}$ as R -modules in the natural way.

Assume the following properties hold:

- (1) $\tilde{\mathbf{A}}$ is a pure \tilde{R} -submodule of $\tilde{\mathbf{B}}$ (in the sense of module theory),
- (2) $\mathbf{C} = \tilde{\mathbf{C}}$,
- (3) \mathbf{C} is a torsion free R -module,
- (4) For any $l \in R \setminus \{0\}$, the quotient $\mathbf{C}/l\mathbf{C}$ is finite and the preimage in $\tilde{\mathbf{B}}$ of any coset $c + l\mathbf{C}$ contains an element which is algebraic over \emptyset .

Let $e \in M^{\text{eq}}$. Then, setting $E := \text{acl}^{\text{eq}}(e)$ and $\Delta := \mathbf{C}(E)$, we have

$$e \in \text{dcl}^{\text{eq}}(\mathbf{C}^{\text{eq}}(E)\tilde{\mathbf{B}}_{\Delta}^{\text{eq}}(E)),$$

where $\tilde{\mathbf{B}}_{\Delta}$ denotes the union of all $\tilde{\mathbf{B}}_{\delta}$ for $\delta \in \Delta$.

Proof. Denote $\iota : \tilde{\mathbf{A}} \rightarrow \tilde{\mathbf{B}}$ and $v : \tilde{\mathbf{B}} \rightarrow \tilde{\mathbf{C}}$ the structural maps.

Note that in particular $\tilde{\mathbf{A}} = \tilde{\mathbf{B}}_0 \subseteq \tilde{\mathbf{B}}_{\Delta}$. By (1), $\tilde{\mathbf{A}}$ and $\tilde{\mathbf{C}}$ are (purely) stably embedded in T , with $\tilde{\mathbf{A}} \perp \tilde{\mathbf{C}}$. Indeed, replacing M by a sufficiently saturated extension, the purity assumption (1) entails that the short exact sequence of \tilde{R} -modules is split, and adding a splitting yields an expansion in which $\tilde{\mathbf{B}}$ is just the product structure of $\tilde{\mathbf{A}}$ and $\tilde{\mathbf{C}}$. Thus the claimed properties hold in an expansion in which $\tilde{\mathbf{A}}$ and $\tilde{\mathbf{C}}$ are unchanged, so the same properties already hold in M . As $\tilde{\mathbf{B}}_{\Delta}$ is internally $\tilde{\mathbf{A}}$ -internal, it follows from Lemma 2.5.18 that $\tilde{\mathbf{B}}_{\Delta}$ is stably embedded in T (over Δ), and one has $\tilde{\mathbf{B}}_{\Delta} \perp \tilde{\mathbf{C}}$.

Since I is finitely generated, \mathbf{A}, \mathbf{B} and \mathbf{C} are definable, and it follows from purity of the initial exact sequence that the induced sequence of R -modules

$$0 \rightarrow \mathbf{A} \rightarrow \mathbf{B} \rightarrow \mathbf{C} \rightarrow 0$$

is also exact.

Let $X \subseteq \tilde{\mathbf{B}}^n$ be $\mathcal{L}(M)$ -definable with $\ulcorner X \urcorner = \text{dcl}^{\text{eq}}(e)$. Assume $\dim_R(v(X)) = r$. Up to passing to some subset of X which is definable over $\text{acl}^{\text{eq}}(\ulcorner X \urcorner) = E$, we may assume, using Lemma 5.1.3, that there are $L \in R^{(n-r) \times r}$, $m \in R \setminus \{0\}$ and $\delta'' \in \mathbf{C}^{n-r}$ such that for $x = x'x''$ with $|x'| = r$, we have $mv(x'') = Lv(x') + \delta''$ for any $x = x'x'' \in X$. We now consider the (fiberwise) action of $(\mathbf{A}^r, +)$ on $\tilde{\mathbf{B}}^n$ given by

$$a \cdot (x', x'') := (x' + ma, x'' + La).$$

Claim 5.1.6. *There is $l \in R \setminus \{0\}$ such that*

$$\dim_R(\{z \in v(X) : X_z \text{ is not } l\mathbf{A}^r\text{-invariant}\}) < r.$$

Proof. Let $\pi(z)$ be the partial type expressing that $z \in v(X)$ and that $\dim_R(z/M) = r$, and let $\tau(y)$ be the partial type expressing that $y \in l\mathbf{A}^r$ for any $l \in R \setminus \{0\}$. Given a solution $(c, a) \models \pi(z) \cup \tau(y)$ in $N \succcurlyeq M$, with $c = c'c''$. Then there is an R -linear map $\theta : \mathbf{C}(N) \rightarrow \mathbf{A}(N)$ which is trivial on $\mathbf{C}(M)$ and such that $\theta(c') = ma$. Indeed, such a map θ may be found as the restriction of an R -linear map from $\mathbf{C}(N) \otimes_R Q(R)$ to $\mathbf{A}(N)$, by choosing a coherent sequence of integer divisors of a .

For $b \in \tilde{\mathbf{B}}(N)$, let $\rho(b) := b + \theta(v(b))$. Then ρ is an \tilde{R} -module-automorphism of $\tilde{\mathbf{B}}(N)$ which induces the identity on $\tilde{\mathbf{B}}(M) \cup \tilde{\mathbf{A}}(N) \cup \mathbf{C}(N)$. It follows from the assumptions that $\rho \in \text{Aut}_{\mathcal{L}}(N/M)$.

In particular, for any $b \in X_c$, we have $\text{tp}(b/M) = \text{tp}(\rho(b)/M)$ and so $\rho(b) \in X_c$. On the other hand, as $c'' = (Lc' + \delta'')/m = Lc'/m + \delta''/m$ and $v(b) = (c', c'')$, using $L \circ \theta = \theta \circ L$ we compute

$$\theta(v(b)) = (\theta(c'), \theta(c'')) = (\theta(c'), \theta(Lc'/m) + \theta(\delta''/m)) = (ma, La),$$

from which it follows that

$$\rho(b) = b + \theta(v(b)) = (b' + \theta(c'), b'' + \theta(c'')) = (b' + ma, b'' + La) = a \cdot (b', b'').$$

Thus X_c is invariant under the action by $\bigcap_{l \in R \setminus \{0\}} l\mathbf{A}^r$. Claim 5.1.6 now follows by compactness. \square

Fix $l \in R \setminus \{0\}$ as in the claim. Arguing by induction on r , we may assume that X_z is $l\mathbf{A}^r$ -invariant for any $z \in v(X)$. In addition, using assumption (4), we may suppose that $v(X) \subseteq ml\mathbf{C}^n$. Indeed, there are only finitely many cosets of $ml\mathbf{C}^n$, all $\text{acl}^{\text{eq}}(\emptyset)$ -definable, so we may assume $X \subseteq v^{-1}(W)$ for a coset W of $ml\mathbf{C}^n$. Replacing X by $X - h$ for some $h \in W \cap \text{acl}(\emptyset)$ if necessary, we may assume $W = ml\mathbf{C}^n$. Let $a \in \tilde{\mathbf{A}}^r$ and $c \in \mathbf{C}^n$. If there exist $b_1 = b'_1 b''_1 \in X_c$ and $b'_0 \in \mathbf{B}^r$ such that $b'_1 = a + mlb'_0$, we set

$$Y_{a,c} := \{b'' - lLb'_0 : (b'_1, b'') \in X\} = X_{(b'_1)} - lLb'_0,$$

where $X_{(b'_1)}$ denotes the fiber $\{b'' \in \tilde{\mathbf{B}}^{n-r} : (b'_1, b'') \in X\}$. Else we set $Y_{a,c} := \emptyset$. Let us first show that in the first case, $Y_{a,c}$ does not depend on the choice of b_1 and b'_0 . Indeed, if $d_1 = d'_1 d''_1 \in X_c$ and $d'_0 \in \mathbf{B}^r$ are such that $d'_1 = a + mld'_0$, then $b'_1 - d'_1 = ml(b'_0 - d'_0)$, so $mlv(b'_0 - d'_0) = 0$, thus $v(b'_0 - d'_0) = 0$, i.e., $b'_0 - d'_0 \in \mathbf{A}^r$. Set $a'_0 := l(b'_0 - d'_0)$. For $d'' \in X_{(d'_1)}$, the invariance of X_c under $l\mathbf{A}^r$ then yields

$$a'_0 \cdot (d'_1, d'') = (d'_1 + ma'_0, d'' + La'_0) = (b'_1, d'' + La'_0) \in X_c,$$

so $d'' + lL(b'_0 - d'_0) = d'' + La'_0 \in X_{(b'_1)}$, showing that $X_{(d'_1)} - lLd'_0 \subseteq X_{(b'_1)} - lLb'_0$. By symmetry, we get the other inclusion $X_{(b'_1)} - lLb'_0 \subseteq X_{(d'_1)} - lLd'_0$, thus $X_{(b'_1)} - lLb'_0 = X_{(d'_1)} - lLd'_0$.

Let $\delta'' = (\delta''_i)_{1 \leq i \leq n-r}$ and let $b' = a + mlb'_0$ be as in the definition of $Y_{a,c}$. Then for $y = b'' - lLb'_0 \in Y_{a,c}$, we compute

$$mv(y) = mv(b'') - mlLv(b'_0) = (Lv(b') + \delta'') - Lv(b') = \delta'',$$

yielding $Y_{a,c} \subseteq \tilde{\mathbf{B}}_{\delta''/m} = \prod_{i=1}^{n-r} \tilde{\mathbf{B}}_{\delta''_i/m}$. It follows that

$$Y \subseteq (\tilde{\mathbf{B}}_{\delta''/m} \times \tilde{\mathbf{A}}^r) \times \mathbf{C}^n.$$

As $\delta''/m \in \mathbf{C}(E) = \Delta$ and $\tilde{\mathbf{B}}_{\Delta} \perp \mathbf{C}$, using Lemma 5.1.1(1), we finish the proof of the theorem once the following claim is established.

Claim 5.1.7. $\ulcorner X \urcorner$ and $\ulcorner Y \urcorner$ are interdefinable.

It is clear by construction that Y is $\ulcorner X \urcorner$ -definable. For the converse, we will use that we have reduced to the case where $v(X) \subseteq ml\mathbf{C}^n$. We may thus reconstruct X from Y as follows:

$$d = d'd'' \in X \Leftrightarrow \exists a \in \tilde{E}^r \exists d'_0 \in \mathbf{B}^r : d' = a + mld'_0 \text{ and } d'' \in Y_{a,v(d)} + lLd'_0$$

This yields the claim. \square

5.2. Some variants. We will now state two variants of Theorem 5.1.5, tailor made for our applications to (enriched) henselian valued fields.

Variante 5.2.1. Let \mathcal{L} be a multisorted language, $\mathcal{A} \sqcup \{\tilde{\mathbf{B}}\} \sqcup \mathcal{C}$ a partition of the sorts of \mathcal{L} and let $\tilde{\mathbf{A}} \in \mathcal{A}$ and $\tilde{\mathbf{C}} \in \mathcal{C}$. Let \tilde{R} be a ring and $R = \tilde{R}/I$ an integral domain, with I a finitely generated ideal. Let

$$(5.1) \quad 0 \rightarrow \tilde{\mathbf{A}} \rightarrow \tilde{\mathbf{B}} \rightarrow \tilde{\mathbf{C}} \rightarrow 0$$

be a short exact sequence of \tilde{R} -modules. Let M be an \mathcal{L} -structure which is an \mathcal{A} - \mathcal{C} -enrichment of the pure (in the sense of model theory) sequence of \tilde{R} -modules (5.1). Assume that the properties (1)-(4) from the statement of Theorem 5.1.5 hold.

Let $e \in M^{\text{eq}}$. Then, setting $E := \text{acl}^{\text{eq}}(e)$ and $\Delta := \mathbf{C}(E)$, we have

$$e \in \text{dcl}^{\text{eq}}(\mathbf{C}^{\text{eq}}(E) \cup (\mathcal{A} \cup \tilde{\mathbf{B}}_{\Delta})^{\text{eq}}(E)),$$

where $\tilde{\mathbf{B}}_{\Delta}$ denotes the union of all $\tilde{\mathbf{B}}_{\delta}$ for $\delta \in \Delta$.

Proof. The proof is a slight variation of the proof of Theorem 5.1.5. Let us indicate the necessary adaptations.

Firstly, it follows from the assumptions that \mathcal{A} and \mathcal{C} are (purely) stably embedded with $\mathcal{A} \perp \mathcal{C}$. Thus, by Lemma 2.5.18, $(\mathcal{A} \cup \tilde{\mathbf{B}}_{\Delta}) \perp \mathcal{C}$ and $\mathcal{A} \cup \tilde{\mathbf{B}}_{\Delta}$ is stably embedded.

Given $e \in M^{\text{eq}}$, we choose an $\mathcal{L}(M)$ -definable set $X \subseteq \mathbf{A}' \times \mathbf{C}' \times \tilde{\mathbf{B}}^n$ with $e = \ulcorner X \urcorner$, where \mathbf{A}' is a finite product of sorts from \mathcal{A} and \mathbf{C}' is a finite product of sorts from \mathcal{C} . For $(a', c') \in \mathbf{A}' \times \mathbf{C}'$, let

$${}_{(a',c')}X \subseteq \tilde{\mathbf{B}}^n$$

be the fiber over (a', c') . Performing the same reductions as in the proof of Theorem 5.1.5, by compactness, we may assume that there is an \emptyset -definable set

$$Y \subseteq \mathbf{A}' \times \mathbf{C}' \times \tilde{\mathbf{B}}_\delta \times \mathbf{C}^n$$

for some finite tuple $\delta \in \Delta$ such that for any $(a', c') \in \mathbf{A}' \times \mathbf{C}'$, $\ulcorner (a', c')X \urcorner$ and $\ulcorner (a', c')Y \urcorner$ are interdefinable, so in particular, $\ulcorner X \urcorner$ and $\ulcorner Y \urcorner$ are interdefinable. The result then follows from Lemma 5.1.1, since $(\mathcal{A} \cup \tilde{\mathbf{B}}_\Delta) \perp \mathcal{C}$. \square

The second variant is designed for applications to henselian valued fields in mixed characteristic.

Variant 5.2.2. *Let \mathcal{L} be a multisorted language, $\mathcal{A} \sqcup \{\tilde{\mathbf{B}}_n : n \in \mathbb{N}\} \sqcup \mathcal{C}$ a partition of the sorts of \mathcal{L} . For any $n \in \mathbb{N}$, let $\tilde{\mathbf{A}}_n \in \mathcal{A}$, and let $\tilde{\mathbf{C}} \in \mathcal{C}$. Let \tilde{R} be a ring and $R = \tilde{R}/I$ an integral domain, with I a finitely generated ideal. Let $\tilde{\mathbf{A}} = (\tilde{\mathbf{A}}_n)_{n \in \mathbb{N}}$, $\tilde{\mathbf{B}} = (\tilde{\mathbf{B}}_n)_{n \in \mathbb{N}}$ be projective systems of \tilde{R} -modules with surjective transition functions, let $\tilde{\mathbf{C}} = (\tilde{\mathbf{C}}_n)_{n \in \mathbb{N}}$ be the projective system with $\tilde{\mathbf{C}}_n = \tilde{\mathbf{C}}$ for all n and identical transition functions. Let*

$$(5.2) \quad 0 \rightarrow \tilde{\mathbf{A}} \rightarrow \tilde{\mathbf{B}} \rightarrow \tilde{\mathbf{C}} \rightarrow 0$$

be a short exact sequence of projective systems of \tilde{R} -modules.

Let M be an \mathcal{L} -structure which is an \mathcal{A} - \mathcal{C} -enrichment of the pure (in the sense of model theory) sequence of projective systems of \tilde{R} -modules (5.2).

Assume that for every $n \in \mathbb{N}$ the exact sequence $0 \rightarrow \tilde{\mathbf{A}}_n \rightarrow \tilde{\mathbf{B}}_n \rightarrow \tilde{\mathbf{C}} \rightarrow 0$ satisfies the properties (1)-(4) from the statement of Theorem 5.1.5.

Let $e \in M^{\text{eq}}$. Then, setting $E := \text{acl}^{\text{eq}}(e)$ and $\Delta := \mathbf{C}(E)$, we have

$$e \in \text{dcl}^{\text{eq}}(\tilde{\mathbf{C}}^{\text{eq}}(E)(\mathcal{A} \cup \tilde{\mathbf{B}}_\Delta)^{\text{eq}}(E)),$$

where $\tilde{\mathbf{B}}_\Delta$ denotes the union of all $(\tilde{\mathbf{B}}_n)_\delta$ for $\delta \in \Delta$ and $n \in \mathbb{N}$.

Proof. Let us first show that \mathcal{A} and \mathcal{C} are (purely) stably embedded in M such that $\mathcal{A} \perp \mathcal{C}$. For this, given $N \in \mathbb{N}$, we consider the structure M_N given by restricting M to the sorts $\mathcal{A} \sqcup \{\tilde{\mathbf{B}}_m : m \leq N\} \sqcup \mathcal{C}$. For $m \leq N$ we denote by $p_{N,m}$ the structural map from $\tilde{\mathbf{B}}_N$ to $\tilde{\mathbf{B}}_m$ and by $q_{N,m}$ the one from $\tilde{\mathbf{A}}_N$ to $\tilde{\mathbf{A}}_m$. For any $m \leq N$, the sequence \tilde{S}_m of \tilde{R} -modules

$$0 \rightarrow \tilde{\mathbf{A}}_m \rightarrow \tilde{\mathbf{B}}_m \rightarrow \tilde{\mathbf{C}} \rightarrow 0$$

is interpretable in the sequence \tilde{S}_N once a predicate for $\ker(q_{N,m}) \leq \tilde{\mathbf{A}}_N$ is added. Thus, M_N may be seen as an \mathcal{A} - \mathcal{C} -enrichment of \tilde{S}_N .

As in the previous proofs, it follows that $\mathcal{A} \cup \tilde{\mathbf{B}}_\Delta$ is stably embedded in M , with $(\mathcal{A} \cup \tilde{\mathbf{B}}_\Delta) \perp \tilde{\mathbf{C}}$. Given $e \in M^{\text{eq}}$, we choose $X \subseteq \mathbf{A}' \times \mathbf{C}' \times \tilde{\mathbf{B}}_k^n$ $\mathcal{L}(M)$ -definable with $e = \ulcorner X \urcorner$, where \mathbf{A}' is a finite product of sorts from \mathcal{A} , \mathbf{C}' is a finite product of sorts from \mathcal{C} and $k \in \mathbb{N}$. Let $N \geq k$ be such that X may be defined using formulas involving only variables from sorts in $\mathcal{A} \cup \mathcal{C} \cup \{\tilde{\mathbf{B}}_i : i \leq N\}$. Since, for $N \geq m$, \tilde{S}_m is interpretable in an \mathcal{A} - \mathcal{C} -enrichment of \tilde{S}_N , we may conclude with Variant 5.2.1. \square

5.3. Imaginaries in RV. Recall that in a finitely ramified henselian valued field, the projective system of short exact sequences:

$$1 \rightarrow \mathbf{R}_n^\times \rightarrow \mathbf{RV}_n^\times \rightarrow \mathbf{\Gamma}^\times \rightarrow 0$$

is stably embedded with the induced structure a $\mathbf{\Gamma}$ - \mathbf{R} -enrichment of the pure short exact sequence of abelian groups. Thus Variant 5.2.2 applies and yields the following elimination of imaginaries:

Proposition 5.3.1. *Let M be a $\mathbf{\Gamma}$ - \mathbf{R} -enriched finitely ramified henselian field, $A \subseteq \mathcal{G}(M)$, $e \in (\mathbf{RV} \cup \mathbf{Lin}_A)^{\text{eq}}(M)$ and $E = \text{acl}^{\text{eq}}(e)$. Assume that:*

- *For every $n, \ell \in \mathbb{Z}_{>0}$, $\mathbf{\Gamma}/\ell\mathbf{\Gamma}$ is finite and the preimage in \mathbf{RV}_n of any coset of $\ell\mathbf{\Gamma}$ contains an element which is algebraic over \emptyset .*

Then $e \in \text{dcl}^{\text{eq}}(\mathbf{\Gamma}^{\text{eq}}(E) \cup (\mathbf{Lin}_A \cup \mathbf{RV}_{\mathbf{\Gamma}(E)})^{\text{eq}}(E))$.

In particular, for $A = \text{acl}^{\text{eq}}(A) \subseteq M^{\text{eq}}$, $(\mathbf{RV} \cup \mathbf{Lin}_{\mathcal{G}(A)})^{\text{eq}}(A) \subseteq \text{dcl}^{\text{eq}}(\mathbf{\Gamma}^{\text{eq}}(A) \cup \mathbf{Lin}_{\mathcal{G}(A)}^{\text{eq}}(A))$.

Proof. We apply Variant 5.2.2 with $R = \tilde{R} = \mathbb{Z}$. Since $\mathbf{\Gamma}$ is ordered is it a torsion free \mathbb{Z} -module, so (1) - (3) hold and (4) holds by hypothesis. \square

These result also apply with an automorphism:

Proposition 5.3.2. *Let $M \models \text{VFA}_{0,0}^{\text{mult}}$, $A \subseteq \mathcal{G}(M)$, $e \in (\mathbf{RV} \cup \mathbf{Lin}_A)^{\text{eq}}(M)$ and $E = \text{acl}^{\text{eq}}(e)$. Then $e \in \text{dcl}^{\text{eq}}(\mathbf{A}\mathbf{\Gamma}(E)\mathbf{RV}_{\mathbf{\Gamma}(E)}(E)\mathbf{Lin}_A(E))$.*

In particular, for $A = \text{acl}^{\text{eq}}(A) \subseteq M^{\text{eq}}$, $(\mathbf{RV} \cup \mathbf{Lin}_{\mathcal{G}(A)})^{\text{eq}}(A) \subseteq \text{dcl}^{\text{eq}}(\mathcal{G}(A))$.

Proof. We apply Variant 5.2.1 with $R = \mathbb{Z}[\sigma]$ and $I := \{P \in \mathbb{Z}[\sigma] : P(\mathbf{\Gamma}) = 0\}$. Hypothesis (1) holds by assumption. Hypothesis (2) and (3) hold by multiplicativity: if $c \in \mathbf{\Gamma}_{>0}$ and $P \in \mathbb{Z}[\sigma]$ are such that $P(c) = 0$, then for all $c \in \mathbf{\Gamma}$, $P(c) = 0$ and $P \in I$. Finally hypothesis (4) holds by divisibility.

So $e \in \text{dcl}^{\text{eq}}(\mathbf{\Gamma}^{\text{eq}}(E) \cup (\mathbf{Lin}_A \cup \mathbf{RV}_{\mathbf{\Gamma}(E)})^{\text{eq}}(E))$. But $\mathbf{\Gamma}$ is an ordered vector field over (the field of fraction of) $\mathbb{Z}[\sigma]/I$, so it eliminates imaginaries. Also, by Proposition 2.5.19, $\mathbf{Lin}_A \cup \mathbf{RV}_{\mathbf{\Gamma}(E)}$ weakly eliminate imaginaries. So $\mathbf{\Gamma}^{\text{eq}}(E) \subseteq \text{dcl}^{\text{eq}}(\mathbf{\Gamma}(E))$ and

$$(\mathbf{Lin}_A \cup \mathbf{RV}_{\mathbf{\Gamma}(E)})^{\text{eq}}(E) \subseteq \text{dcl}^{\text{eq}}(\mathbf{Lin}_A(E)\mathbf{RV}_{\mathbf{\Gamma}(E)}(E)).$$

The result follows. \square

6. IMAGINARIES IN VALUED FIELDS

6.1. The henselian case. Let Hen_0^{ac} denote the \mathbf{RV} -enrichment of Hen_0 with a compatible system of angular components $\text{ac}_n : \mathbf{K} \rightarrow \mathbf{R}_n^\times$ — we denote this language by \mathcal{L}_{ac} . We fix T a $\mathbf{\Gamma}$ - \mathbf{k} -enrichment of either Hen_0 or Hen_0^{ac} . Recall that the two sorted language \mathcal{L}_{mod} is given by a sort \mathbf{A} endowed with the language of rings, a sort \mathbf{V} endowed with the language of abelian groups and a function symbol $\mu : \mathbf{A} \times \mathbf{V} \rightarrow \mathbf{V}$ for scalar multiplication. Let t_ℓ denote the \emptyset -induced theory on \mathbf{R}_ℓ . For every $X \subseteq \mathbf{V}^2$ definable in the \mathbf{A} -enriched theory of free rank $n \in \mathbb{Z}_{>0}$ modules over models of t_ℓ ,

we define the equivalence relation E_X on \mathbf{V} by vE_Xw if $X_v = X_w$ and $\mathbf{T}_{n,\ell,X} := \bigsqcup_{s \in \mathbf{S}_n} (\mathbf{V}/E_X)^{(\mathbf{R}_{\ell,s/\ell\mathbf{m}s})}$. Let \mathbf{R}^{leq} denote $\bigsqcup_{n,\ell,X} \mathbf{T}_{n,\ell,X}$, the \mathbf{R} -linear imaginaries. We can now prove our imaginary Ax-Kochen-Ershov principle:

Theorem 6.1.1 (Theorem A). *Let T be a Γ - \mathbf{k} -enrichment of either Hen_0 or Hen_0^{ac} , such that:*

- (\mathbf{C}_Γ) *T has definably complete value group;*
- (\mathbf{FR}) *For every $\ell \in \mathbb{Z}_{>0}$, the interval $[0, v(\ell)]$ is finite and \mathbf{k} is perfect;*
- ($\mathbf{I}_\mathbf{k}$) *The residue field \mathbf{k} is infinite;*
- ($\mathbf{E}_\mathbf{k}^\infty$) *The induced theory on \mathbf{k} eliminates \exists^∞ .*

Then T weakly eliminates imaginaries in $\mathbf{K} \cup \Gamma^{\text{eq}} \cup \mathbf{R}^{\text{leq}}$.

Proof. We will use Proposition 4.4.8. By Theorem 3.1.3, hypothesis (\mathbf{D}) holds. Hypothesis ($\mathbf{Q}_\mathbf{K}$) holds trivially for $\mathcal{L}_1 = \mathcal{L}$ — and $f = \text{id}$. Also, \mathbf{RV} and \mathbf{R} are stably embedded in characteristic zero henselian fields. Let $M \models T$, $e \in M^{\text{eq}}$ and $A = \text{acl}^{\text{eq}}(e)$. By Proposition 4.4.8, $e \in \text{dcl}^{\text{eq}}(\mathbf{K}(A) \cup (\mathbf{RV} \cup \mathbf{Lin}_{\mathcal{G}(A)})^{\text{eq}}(A))$.

Claim 6.1.2. $(\mathbf{RV} \cup \mathbf{Lin}_{\mathcal{G}(A)})^{\text{eq}}(A) \subseteq \text{dcl}^{\text{eq}}(\Gamma^{\text{eq}}(A) \cup \mathbf{Lin}_{\mathcal{G}(A)}^{\text{eq}}(A))$

Proof. If $T \supseteq \text{Hen}_0^{\text{ac}}$, then \mathbf{RV}_n is \mathcal{L}_{ac} isomorphic to $\mathbf{R}_n^\times \times \Gamma$ and the isomorphisms are compatible as n varies. It follows that $(\mathbf{RV} \cup \mathbf{Lin}_{\mathcal{G}(A)})^{\text{eq}} \subseteq (\Gamma \cup \mathbf{Lin}_{\mathcal{G}(A)})^{\text{eq}}$, and since Γ and $\mathbf{Lin}_{\mathcal{G}(A)}$ are orthogonal, the claim follows. If T is a Γ - \mathbf{k} -enrichment of Hen_0 , the claim follows from Proposition 5.3.1. Note that by (\mathbf{C}_Γ), $\Gamma \cong \mathbb{Q}$ or $\Gamma \cong \mathbb{Z}$ and hence $\Gamma/n\Gamma$ is finite and every coset is represented in $\Gamma(\text{dcl}^{\text{eq}}(\emptyset))$. \square

Claim 6.1.3. $\mathbf{Lin}_{\mathcal{G}(A)}^{\text{eq}}(A) \subseteq \text{dcl}^{\text{eq}}(\mathbf{R}^{\text{leq}}(A))$

Proof. Recall that $\mathbf{Lin}_{\mathcal{G}(A)}$ is stably embedded. It follows that, for every $e \in \mathbf{Lin}_{\mathcal{G}(A)}^{\text{eq}}(A)$, taking tensor products of lattices, we may assume that there exists $n, \ell \in \mathbb{Z}_{>0}$ and $s \in \mathbf{S}_n(A)$ such that e codes some subset X_a of $s/\ell\mathbf{m}s$ and a a single parameter in $s/\ell\mathbf{m}s$. Since $s/\ell\mathbf{m}s$ is definably isomorphic to \mathbf{R}_ℓ^n once we name a basis, it follows that X is definable with parameters in the \mathcal{L}_{mod} -structure $(\mathbf{R}_{\ell,s/\ell\mathbf{m}s})$ — so $e \in \text{dcl}^{\text{eq}}(\mathbf{T}_{n,\ell,X}(A))$. \square

It follows that $e \in \text{dcl}^{\text{eq}}(\mathbf{K}(A) \cup \Gamma^{\text{eq}}(A) \cup \mathbf{R}^{\text{leq}}(A))$, which concludes the proof. \square

Let $\mathbf{k}^{\text{leq}} := \bigsqcup_{s,n,X} (\mathbf{V}/E_X)^{(\mathbf{k},s/\mathbf{m}s)}$.

Corollary 6.1.4. *Let F be a characteristic zero field that eliminates \exists^∞ . Then any valued field elementarily equivalent to $F((t))$ or $F((t^\mathbb{Q}))$ (with or without angular components) weakly eliminate imaginaries in $\mathbf{K} \cup \mathbf{k}^{\text{leq}}$. \square*

In certain cases, the elimination of imaginaries in $\text{Th}(\mathbf{k})$ -linear structures allows to further reduce this result to the geometric sorts. We can then also code finite sets, by adapting an argument of Johnson [Joh20, Section 5.3]:

Proposition 6.1.5. *Let M be a henselian valued field such that:*

- (**I_k**) *The residue field \mathbf{k} is infinite;*
- (**Cb_K**) *For any $A = \text{dcl}^{\text{eq}}(A) \subseteq M^{\text{eq}}$, any $\mathcal{L}(A)$ -definable type $p \in S_{\mathbf{K}^n}^0(M)$ finitely satisfiable in M is $\mathcal{L}_0(\mathcal{G}(A))$ -definable.*

Then every finite set in \mathcal{G} is coded in \mathcal{G} .

Proof. The proposition follows from:

Claim 6.1.6. *Let $C \subseteq \mathcal{G}$ be a finite set. There exists an $\mathcal{L}(\ulcorner C \urcorner)$ -definable type $p_C \in S_{\mathbf{K}^n}^0(M)$ finitely satisfiable in M such that C is $\mathcal{L}(a)$ -definable for any $a \models p_C$.*

Indeed taking p_C as in the claim, by (**Cb_K**), p_C is $\mathcal{G}(\text{dcl}^{\text{eq}}(\ulcorner C \urcorner))$ -definable and hence, so is C .

We now prove the claim. We start by considering $C = \{s\} \subseteq \mathbf{S}_n$. By (**I_k**), the generic type q_n of $\text{GL}_n(\mathcal{O})$ (that is, the quantifier free type that reduces to the generic of $\text{GL}_n(\mathbf{k})$ modulo \mathfrak{m}) is finitely satisfiable in M . Note also that it is \mathcal{L}_0 -definable and symmetric (cf., [Joh20, Section 3.3]): for every definable quantifier free type r , $q_n \otimes r = r \otimes q_n$. Let B be the matrix associated to some basis of s in M and let $p_s = B \cdot q_n$. This type does not depend on B , is $\mathcal{L}_0(s)$ -definable, finitely satisfiable in M and symmetric.

If $C \subseteq \mathbf{S}_n$, let $a \models \bigotimes_{s \in C} p_s$ and A be the set of the a_s , for $s \in S$. Since finite subsets of \mathbf{K} are coded in \mathbf{K} , $\ulcorner A \urcorner$ can be identified with a tuple in \mathbf{K} . Then $p_C = \text{tp}_0(\ulcorner A \urcorner/M)$ does not depend on a choice of enumeration of C and thus it is $\mathcal{L}(\ulcorner C \urcorner)$ -definable (and finitely satisfiable in M). Note that if $\mathbf{\Gamma}(M)$ has a smallest positive element, we are done since, for any lattice s , $\mathfrak{m}s$ is (\mathcal{L} -definably isomorphic to) a lattice and hence \mathbf{T}_n \mathcal{L} -definably embeds in \mathbf{S}_{n+1} by the usual identification of translates of linear spaces with higher dimensional linear spaces.

Let us now assume that $\mathbf{\Gamma}(M)$ does not admit a smallest positive element. We first consider the case where $C \subseteq \mathbf{K}^i \times \mathbf{k}^j$. Let $E \subseteq \mathbf{k}$ be the set of elements of \mathbf{k} appearing as coordinates of elements in C , let b be the tuple of coefficients of the polynomial $\prod_{e \in E} (x - e)$ and let $p_E \in S^0(M)$ be the type of generic lifts of $b \in \mathbf{k}$ to \mathcal{O} . This type is $\mathcal{L}(\ulcorner C \urcorner)$ -definable and finitely satisfiable in M since the value group does not admit a smallest positive element. Then for any $a \models p_E$, res induces a bijection between the $\mathcal{L}_0(a)$ -definable set of roots of the polynomial $\sum_{\ell} a_{\ell} x^{\ell}$ and E . It follows that C is in $\mathcal{L}_0(a)$ -definable bijection with a subset D of \mathbf{K}^{i+j} , which is coded in some cartesian power of \mathbf{K} . Then $p_C = \text{tp}_0(a \ulcorner D \urcorner/M)$ has the required properties. Let us finally consider $e \in \mathbf{T}_n(M)$ and let $s = \tau_n(e)$ (see page 9). If $a \models p_s$, then e is $\mathcal{L}_0(a)$ -definably isomorphic to a tuple $b \in \mathbf{k}$. The type $p_e = \text{tp}_0(ab/M)$ is $\mathcal{L}_0(e)$ -definable, finitely satisfiable in M and symmetric — since \mathbf{k} is quantifier free stable. If $C \subseteq \mathbf{T}_n$, let $a \models \bigotimes_{e \in C} p_e$ and $A \subseteq \mathbf{K}^i \times \mathbf{k}^j$ be the set of the a_e , for $e \in C$. Then, applying the previous paragraph to A , we find p_C as required. \square

The authors would like to thank Ehud Hrushovski for his insights on the correct statement of the following corollary.

Corollary 6.1.7. *Let F be a characteristic zero field that is either algebraically closed, pseudofinite or real closed. Then any valued field M elementarily equivalent to $F((t))$ or $F((t^{\mathbb{Q}}))$ (with or without angular components), eliminates imaginaries in \mathcal{G} provided we add the following (imaginary) constants:*

- if F is real closed and the value group is a \mathbb{Z} -group with minimal positive element γ_0 , a constant for a half line of \mathbf{RV}_{1,γ_0} ;
- if F is pseudofinite, constants for a generator of Galois of F — see [Hru12, Section 5.9];
- if F is pseudofinite and the value group is a \mathbb{Z} -group with minimal positive element γ_0 , a constant for a $(\mathbf{k}^*)^n$ -orbit in \mathbf{RV}_{1,γ_0} , for every $n \geq 1$.

Proof. Let $A = \text{dcl}^{\text{eq}}(A) \subseteq M^{\text{eq}}$. By 2.5.13, $\mathbf{Lin}_A(M)$ is a $\text{Th}(\mathbf{k}(M))$ -linear structure with flags. If $\mathbf{k}(M)$ is algebraically closed, by [Hru12, Lemma 5.6], it eliminates imaginaries. If $\mathbf{k}(M)$ is real closed, \mathbf{Lin}_A also eliminates imaginaries by Propositions 2.5.21 and 2.5.24 and Remark 2.5.25. If $\mathbf{k}(M)$ is pseudofinite and $\Gamma(M)$ is divisible, $\mathbf{Lin}_A(M)$ has roots and we conclude with [Hru12, Theorem 5.10]. If $\Gamma(M)$ is a \mathbb{Z} -group, \mathbf{Lin}_A does not have roots, but by [Hru12, Remark 5.8] it suffices to insure the existence of an \emptyset -definable $(\mathbf{k}^*)^n$ -orbit inside each one dimensional vector space of \mathbf{Lin}_A , i.e., each $\mathbf{RV}_{1,\gamma}$ with $\gamma \in \Gamma(A)$. For every n , write γ as $i\gamma_0 + n\delta$. By assumption, there exists \emptyset -definable $(\mathbf{k}^*)^n \cdot \xi \subseteq \mathbf{RV}_{1,\gamma_0}$. Then $(\mathbf{k}^*)^n \cdot \xi^i \zeta^n \subseteq \mathbf{RV}_{1,\gamma}$ is independent of the choice of $\zeta \in \mathbf{RV}_{1,\delta}$ and thus \emptyset -definable.

It now follows from Corollary 6.1.4 that M weakly eliminates imaginaries in \mathcal{G} and we conclude with Proposition 6.1.5. \square

Not all of these results are new, although all of the statements with angular components are. The case of $\mathbb{C}((t^{\mathbb{Q}}))$ just amounts to Haskel-Hrushovski-Macpherson's result [HHM06] for ACVF. The case of $\mathbb{R}((t^{\mathbb{Q}}))$ is Melior's result [Mel06] for RCVF and the case of $F((t))$, with F pseudofinite, is Hrushovski-Martin-Rideau's result [HMR18] for pseudolocal fields — slightly improved since we only require algebraic constants in $\mathbf{RV}_1^{\text{eq}}$ and not in \mathbf{K} .

Corollary 6.1.8. *Let F be a positive characteristic perfect field that eliminates \exists^∞ . Then $W(F)$ (with or without compatible angular components) weakly eliminate imaginaries in $\mathbf{K} \cup \mathbf{R}^{\text{leq}}$. \square*

It seems plausible that, if $F \models \text{ACF}$, the \mathbf{R} -linear imaginaries can also be eliminated, yielding elimination of imaginaries in \mathcal{G} for $W(\mathbb{F}_p^{\text{a}})$. However, this remains an open problem.

6.2. The σ -henselian case. Let us conclude with the description of the imaginaries in $\text{VFA}_{0,0}^{\text{mult}}$:

Theorem 6.2.1 (Theorem B). *The theory $\text{VFA}_{0,0}^{\text{mult}}$ (with or without equivariant angular components) eliminates imaginaries in \mathcal{G} .*

Proof. Any model of $\text{VFA}_{0,0}^{\text{mult}}$ is elementarily equivalent to a maximally complete one and hence (\mathbf{C}_B) holds. By Fact 2.4.7, the structure induced on Γ is σ -minimal. So (\mathbf{C}_Γ) and $(\mathbf{E}_\Gamma^\infty)$ hold. Finally, (\mathbf{E}_k^∞) holds since ACFA eliminates \exists^∞ . By Theorem 3.1.1, (\mathbf{D}) holds. Hypothesis (\mathbf{Q}_K) follows from Fact 2.4.4, with $\mathcal{L}_1 := \mathcal{L}_{\mathbf{RV}} \cup \{\sigma_{\mathbf{RV}}\}$ and $f(x) := (\sigma^n(c))_{n \in \mathbb{Z}_{\geq 0}}$. So, by Proposition 4.4.8, for every $M \models \text{VFA}_{0,0}^{\text{mult}}$, $e \in M^{\text{eq}}$ and $A = \text{acl}^{\text{eq}}(e)$, we have $e \in \text{dcl}^{\text{eq}}(\mathbf{K}(A) \cup (\mathbf{RV} \cup \text{Lin}_{\mathcal{G}(A)})^{\text{eq}}(A))$. By Proposition 5.3.2 — or using the angular components — we have $(\mathbf{RV} \cup \text{Lin}_{\mathcal{G}(A)})^{\text{eq}}(A) \subseteq \text{dcl}^{\text{eq}}(\mathcal{G}(A))$. \square

Similar results hold in differential valued fields.

Corollary 6.2.2. *The following two families of difference valued fields, indexed by integer primes p :*

- (1) $K_p := (\mathbb{F}_p(t)^a, v_t, \phi_p)$, where ϕ_p is the Frobenius automorphism;
- (2) $K_p := (\mathbb{C}_p, v_p, \sigma_p)$, where σ_p is an isometric lift of the Frobenius automorphism on $\mathbf{k}(\mathbb{C}_p) = \mathbb{F}_p^a$.

uniformly eliminate imaginaries in \mathcal{G} for large p : for every $\mathcal{L}_{\mathbf{RV}}^\sigma$ -definable sets $X \subseteq Y \times Z$, there exists an $\mathcal{L}_{\mathbf{RV}}^\sigma$ -definable map $f: Z \rightarrow W$, where W is a product of sorts in \mathcal{G} and some $N \in \mathbb{Z}_{\geq 0}$ such that, for every prime $p > N$ and $z_1, z_2 \in Z(K_p)$, $f(z_1) = f(z_2)$ iff and only if $X_{z_1}(K_p) = X_{z_2}(K_p)$. \square

As noted earlier, in case (1), since K_p is a definable expansion of a model of ACVF which also eliminates imaginaries in the geometric sorts, the result is even uniform in all p .

REFERENCES

- [AK65] J. Ax and S. Kochen. Diophantine problems over local fields I. *Amer. J. Math.* 87.3 (1965), pp. 605–630.
- [Azg10] S. Azgin (now known as Durhan). Valued fields with contractive automorphism and Kaplansky fields. *J. Algebra* 324.10 (2010), pp. 2757–2785.
- [AvdD10] S. Azgin (now known as Durhan) and L. van den Dries. Elementary theory of valued fields with a valuation-preserving automorphism. *J. Inst. Math. Jussieu* 9.4 (2010), pp. 1–35.
- [Bas91] Ş. A. Basarab. Relative elimination of quantifiers for Henselian valued fields. *Ann. Pure Appl. Logic* 53.1 (1991), pp. 51–74.
- [Bau82] W. Baur. “Die Theorie der Paare reell abgeschlossener Körper”. In: *Logic and Algorithmic: An International Symposium Held in Honour of Ernst Specker*. Monogr. Enseign. Math. 30. Kundig, 1982, pp. 25–34.
- [BMS07] L. Bélair, A. Macintyre, and T. Scanlon. Model theory of the Frobenius on the Witt vectors. *Amer. J. Math.* 129.3 (2007), pp. 665–721.
- [BPW01] S. Buechler, A. Pillay, and F. Wagner. Supersimple theories. *J. Amer. Math. Soc.* 14.1 (2001), pp. 109–124. ISSN: 0894-0347. DOI: 10.1090/S0894-0347-00-00350-7.
- [CH99] Z. Chatzidakis and E. Hrushovski. Model theory of difference fields. *Transactions of the American Mathematical Society* 351.8 (1999), pp. 2997–3071.

- [CP98] Z. Chatzidakis and A. Pillay. Generic structures and simple theories. *Ann. Pure Appl. Logic* 95.1-3 (1998), pp. 71–92. ISSN: 0168-0072. DOI: 10.1016/S0168-0072(98)00021-9.
- [CH14] A. Chernikov and M. Hils. Valued difference fields and NTP_2 . *Israel J. Math.* 204.1 (2014), pp. 299–327. ISSN: 0021-2172. DOI: 10.1007/s11856-014-1094-z.
- [CHR19] R. Cluckers, I. Halupczok, and S. Rideau-Kikuchi. Hensel minimality. arXiv:1909.13792. To appear in Forum of Mathematics, Pi. 2019.
- [CL08] R. Cluckers and F. Loeser. Constructible motivic functions and motivic integration. *Invent. Math.* 173.1 (2008), pp. 23–121.
- [Con+22] G. Conant et al. Enriching a predicate and tame expansions of the integers. arXiv:2203.07226. 2022.
- [Del81] F. Delon. Types sur $\mathbb{C}((x))$. In: *Groupe d'étude sur les théories stables, année 2, 1978/79*. exp. 5. Secrétariat Math., 1981, pp. 1–29.
- [Del82] F. Delon. *Quelques propriétés des corps valués en théorie des modèles*. Thèse de Doctorat d'état. Université Paris VII, 1982.
- [DO15] S. Durhan and G. Onay. Quantifier elimination for valued fields equipped with an automorphism. *Selecta Math. (N.S.)* 21.4 (2015), pp. 1177–1201.
- [EP05] A. J. Engler and A. Prestel. *Valued Fields*. Springer Monographs in Mathematics. Springer-Verlag, 2005.
- [Ers65] J. L. Ershov. On the elementary theory of maximal normed fields. *Dokl. Akad. Nauk SSSR* 165 (1965), pp. 21–23.
- [Fle11] J. Flenner. Relative decidability and definability in Henselian valued fields. *J. Symbolic Logic* 76.4 (2011), pp. 1240–1260. ISSN: 0022-4812. DOI: 10.2178/jsl/1318338847.
- [HHM06] D. Haskell, E. Hrushovski, and D. Macpherson. Definable sets in algebraically closed valued fields: elimination of imaginaries. *J. Reine Angew. Math.* 597 (2006), pp. 175–236.
- [HHM08] D. Haskell, E. Hrushovski, and D. Macpherson. *Stable Domination and Independence in Algebraically Closed Valued Fields*. Lect. Notes Log. 30. Assoc. Symbol. Logic, Chicago, IL; Cambridge Univ. Press, Cambridge, 2008.
- [HKR18] M. Hils, M. Kamensky, and S. Rideau. Imaginaries in separably closed valued fields. *Proc. Lond. Math. Soc. (3)* 116.6 (2018), pp. 1457–1488. ISSN: 0024-6115. DOI: 10.1112/plms.12116.
- [Hru02] E. Hrushovski. Valued fields with automorphisms. manuscript. 2002.
- [Hru04] E. Hrushovski. The elementary theory of the Frobenius automorphism. arXiv: math/0406514. 2004.
- [Hru12] E. Hrushovski. Groupoids, imaginaries and internal covers. *Turkish J. Math.* 36.2 (2012), pp. 173–198.
- [Hru14] E. Hrushovski. Imaginaries and definable types in algebraically closed valued fields. In: *Valuation Theory in Interaction*. Ed. by A. Campillo Lopez, F.-V. Kuhlmann, and B. Teissier. EMS Ser. Congr. Rep. 10. Eur. Math. Soc., Zürich, 2014, pp. 297–319.
- [HK06] E. Hrushovski and D. Kazhdan. Integration in valued fields. In: *Algebraic Geometry and Number Theory*. Progr. Math. 253. Birkhäuser Boston, 2006, pp. 261–405.
- [HL16] E. Hrushovski and F. Loeser. *Non-archimedean tame topology and stably dominated types*. Vol. 192. Ann. of Math. Stud. Princeton University Press, Princeton, NJ, 2016, pp. vii+216.
- [HMR18] E. Hrushovski, B. Martin, and S. Rideau. Definable equivalence relations and zeta functions of groups. *J. Eur. Math. Soc. (JEMS)* 20.10 (2018). With an appendix by Raf Cluckers, pp. 2467–2537.

- [Joh20] W. Johnson. On the proof of elimination of imaginaries in algebraically closed valued fields. *Notre Dame J. Form. Log.* 61.3 (2020), pp. 363–381.
- [Joh21] W. Johnson. Dp-finite fields I(A): The infinitesimals. *Ann. Pure Appl. Logic* 172.6 (2021), Paper No. 102947, 39.
- [Kei64] H. J. Keisler. Complete theories of algebraically closed fields with distinguished subfields. *Michigan Math. J.* 11 (1964), pp. 71–81. ISSN: 0026-2285.
- [Kuh94] F.-V. Kuhlmann. Quantifier elimination for Henselian fields relative to additive and multiplicative congruences. *Israel J. Math.* 85 (1994), pp. 277–306.
- [Lel90] G. Leloup. Théories complètes de paires de corps valués henséliens. *J. Symbolic Logic* 55.1 (1990), pp. 323–339. ISSN: 0022-4812. DOI: 10.2307/2274971.
- [Mel06] T. Mellor. Imaginaries in real closed valued fields. *Ann. Pure Appl. Logic* 139 (2006), pp. 230–279.
- [MR21] S. Montenegro and S. Rideau-Kikuchi. Imaginaries, invariant types and pseudo p -adically closed fields. *Trans. Amer. Math. Soc.* 374.2 (2021), pp. 803–828.
- [Pal12] K. Pal. Multiplicative valued difference fields. *J. Symb. Log.* 77.2 (2012), pp. 545–579.
- [Poi83] B. Poizat. Une théorie de Galois imaginaire. *J. Symb. Log.* 48.4 (1983), pp. 1151–1170.
- [Rid17] S. Rideau. Some properties of analytic difference valued fields. *J. Inst. Math. Jussieu* 16.3 (2017), pp. 447–499.
- [Rid19] S. Rideau. Imaginaries and invariant types in existentially closed valued differential fields. *J. Reine Angew. Math.* 750 (2019), pp. 157–196.
- [Rio17] R. Rioux. *Théorie des modèles d'expansion de corps valués: phénomènes de séparation*. PhD thesis. Université Claude Bernard Lyon 1, 2017.
- [Rob77] A. Robinson. *Complete Theories*. Second. Stud. Logic Found. Math. 46. North-Holland Publishing Co., Amsterdam, 1977, pp. x+129.
- [Sca03] T. Scanlon. “Quantifier elimination for the relative Frobenius”. In: *Valuation Theory and Its Applications, Conference proceedings of the International Conference on Valuation Theory (Saskatoon, 1999)*. Vol. II. Fields Inst. Commun. 33. Amer. Math. Soc., Providence, RI, 2003, pp. 323–352.
- [Ser68] J.-P. Serre. *Corps locaux*. Hermann, 1968.
- [TZ12] K. Tent and M. Ziegler. *A Course in Model Theory*. Assoc. Symbol. Logic, Chicago, IL. Lect. Notes Log. 40. Cambridge Univ. Press, Cambridge, 2012.
- [Vic21] M. Vicaria. Elimination of imaginaries in multivalued fields. arXiv:2109.08140. 2021.
- [Wag00] F. O. Wagner. *Simple theories*. Vol. 503. Mathematics and its Applications. Kluwer Academic Publishers, Dordrecht, 2000, pp. xii+260.
- [Wei99] V. Weispfenning. “Mixed real-integer linear quantifier elimination”. In: *Proceedings of the 1999 International Symposium on Symbolic and Algebraic Computation (Vancouver, BC)*. ACM, New York, 1999, pp. 129–136. DOI: 10.1145/309831.309888.

INSTITUT FÜR MATHEMATISCHE LOGIK UND GRUNDLAGENFORSCHUNG, WESTFÄLISCHE WILHELMS-UNIVERSITÄT MÜNSTER, EINSTEINSTR. 62, D-48149 MÜNSTER, GERMANY
Email address: hils@uni-muenster.de

UNIVERSITÉ DE PARIS CITÉ AND SORBONNE UNIVERSITÉ, CNRS, IMJ-PRG, F-75006 PARIS, FRANCE.
Email address: silvain.rideau@imj-prg.fr