

Article

Closed similarity manifolds.

Fried, David

in: Commentarii mathematici Helvetici | Commentarii Mathematici Helvetici - 55

Terms and Conditions

The Göttingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library. Each copy of any part of this document must contain there Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept there Terms and Conditions. Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact:

Niedersächsische Staats- und Universitätsbibliothek
Digitalisierungszentrum
37070 Goettingen
Germany
Email: gdz@sub.uni-goettingen.de

Purchase a CD-ROM

The Goettingen State and University Library offers CD-ROMs containing whole volumes / monographs in PDF for Adobe Acrobat. The PDF-version contains the table of contents as bookmarks, which allows easy navigation in the document. For availability and pricing, please contact:

Niedersaechsische Staats- und Universitaetsbibliothek Goettingen - Digitalisierungszentrum
37070 Goettingen, Germany, Email: gdz@sub.uni-goettingen.de

Closed similarity manifolds

DAVID FRIED*

We classify similarity structures on closed manifolds. A *similarity structure* on a manifold M is defined by an atlas of charts with gluing functions that are the restrictions of similarity transformations (those affine isomorphisms that change distance by a constant factor) to open subsets of Euclidean space. Thus M has a preferred conformal structure. The universal cover \tilde{M} is locally Euclidean since one may choose a preferred inner product at a basepoint and parallel translate it over \tilde{M} without ambiguity. Similarity manifolds are the affine manifolds (i.e. manifolds with a given connection of zero curvature and zero tension) with a “parallel protractor” in the sense that Euclidean manifolds are affine manifolds with a “parallel ruler.”

The classification we obtain in Theorem 2 below was given (with one omission) by Kuiper but under the implicit hypothesis that the development map D cover its image $[K]$. Here D is an affine immersion from \tilde{M} to Euclidean space E^d (these properties determine $D: \tilde{M} \rightarrow E^d$ up to composition by a similarity of E^d). For general affine structures, the development map doesn't cover its image [S–Th] but it will follow from our classification that D does cover its image for similarity structures. To show this covering property directly would require essentially all of our demonstration, so it is natural to proceed directly to the classification without using [K].

In addition to extending the classification of [K] to all similarity structures, our methods show that similarity manifolds which aren't Euclidean admit a natural metric μ in the preferred conformal class, unlike Euclidean manifolds where the preferred metric can be renormalized by any scale factor. This situation is somewhat reminiscent of the case of Riemannian manifolds of constant curvature K , where the metric can be normalized so that $|K| = 1$ unless the metric is flat ($K = 0$). The non-Euclidean similarity manifolds possess a natural volume which we compute in Theorem 2. Our classification of non-Euclidean closed similarity manifolds of M of dimension $d \geq 3$ shows they correspond 1-1 to the following

* Partially supported by National Science Foundation Grant #MCS-8003622.

arbitrary data

- (1) an isometry of a $d-1$ dimensional Clifford–Klein space
- (2) a positive real value of the natural volume.

Since similarity manifolds arise in the behavior of certain hyperbolic manifolds at infinity [Th] we see that they are closely connected with Riemannian manifolds of constant curvature K for K positive, negative or zero.

We thank Dennis Sullivan and Nico Kuiper for their helpful comments. We gratefully acknowledge the support of I.H.E.S. during the preparation of this paper.

1. Incomplete implies radiant

The uniqueness of D determines similarities $\phi(g) \in \text{Sim}(E^d)$, $g \in \pi_1 M$, with the property that $\phi(g) \circ D = D \circ g: \tilde{M} \rightarrow E^d$. This *holonomy* homomorphism $\phi: \pi_1 M \rightarrow \text{Sim}(E^d)$ is crucial to the study of similarity structures. We call the similarity structure *reducible* if $\phi(\pi_1 M)$ fixes some affine subspace of E^d and *radiant* when this subspace is a single point [FGH]. Choosing the fixed point as origin, radiant similarity structures are based on the group of linear similarities $\text{Sim}_0(E^d) = \text{Sim}(E^d) \cap \text{Gl}(d, \mathbf{R})$. Since we later show that (geodesically) complete similarity manifolds are Euclidean, the following theorem shows the structure group of a similarity manifold may always be reduced to either $\text{Sim}_0(E^d)$ or the Euclidean group $\text{Euc}(E^d)$.

THEOREM 1. *A connected closed similarity manifold which is incomplete must be radiant.*

Proof. Fix a development map $D: \tilde{M} \rightarrow E^d$. This defines a Euclidean metric on \tilde{M} and determines scale factors $\alpha(g) \in \mathbf{R}^+$ for $g \in \pi_1 M$ by the rule $\|gv\| = \alpha(g)\|v\|$, $v \in T\tilde{M}$. Here $\alpha: \pi_1 M \rightarrow \mathbf{R}^+$ is a homomorphism related to the holonomy ϕ by the equation $\alpha(g)^d = |\det \phi(g)|$.

For each $\tilde{m} \in \tilde{M}$ let $D_{\tilde{m}}$ be the largest open disc in $T_{\tilde{m}}\tilde{M}$ on which \exp is defined. Let $r(\tilde{m}) \in (0, \infty]$ be the radius of $D_{\tilde{m}}$. We see that $r(\tilde{m}) \geq r(\tilde{n}) - \text{dist}(\tilde{m}, \tilde{n})$ (Fig. 1). Thus r satisfies $|r(x) - r(y)| \leq \text{dist}(x, y)$.

Since M is incomplete, r is finite at some point and hence finite on all of \tilde{M} . Clearly r is continuous and $r(g\tilde{m}) = \alpha(g)r(\tilde{m})$ for all $g \in \pi_1 M$. Thus

$$\frac{\|Tg(y)\|}{r(g\tilde{m})}: T\tilde{M} \rightarrow [0, \infty)$$

is independent of g and defines a continuous metric μ on M such that

- (1) μ is in the preferred conformal class
- (2) for each $m \in M$, the unit disc in $T_m M$ is the largest disc (relative to any preferred inner product at m) on which \exp is defined.

For any fixed $m \in M$ there is a $v_0 \in \partial D_m$ such that $\gamma(t) = \exp(tv_0)$ is defined for $0 \leq t < 1$ but not for $t = 1$. The μ -speed of the affine ray $\gamma(t)$ is $(\mu(\gamma'(t), \gamma'(t)))^{1/2} = 1/(1-t)$ that is inversely proportional to the distance from $\gamma(t)$ to $\partial(\exp D_m)$ (Fig. 2). Since $\gamma(t)$ has infinite μ -length it passes by some point $p \in M$ infinitely often and arbitrarily closely. Fixing some orthonormal frame F_m at m there is an orthonormal frame F_p at p and times $t_i \nearrow 1$ such that F_p is the limit of $(1-t_i) \cdot (\gamma|_{[0, t_i]} F_m)$ which denotes the orthonormal frame obtained by parallel translating F_m along $\gamma(t)$ for time t_i and scaling down by $(1-t_i)$. Let g_{ij} denote the element of $\pi_1(M, p)$ approximated by $\gamma|_{[t_i, t_j]}$.




Fig. 1.




Fig. 2.

Now lift everything to \tilde{M} and fix $\tilde{w} \in T_{\tilde{m}} \tilde{M}$ with inner product $\tilde{\mu}(\tilde{w}, \tilde{v}_0) < 1$. For $i \gg 0$ and $j \gg i$, g_{ij} carries $\tilde{\gamma}(t_i)$ close to $\tilde{\gamma}(t_j)$ in the $\tilde{\mu}$ metric. The holonomy $\phi(g_{ij})$ is a very sharp contraction ($\alpha(g_{ij}) \approx (1-t_j)/(1-t_i)$) with almost no rotation (Fig. 3). Using the local coordinate given by D , we see that (for $j \gg i \gg 0$) $\exp(t \cdot g_{ij} \tilde{w})$ lies within $\exp D_{\tilde{m}}$ for all $t \in [0, 1]$. Thus, applying g_{ij}^{-1} , we see that $\exp \tilde{w}$ is defined.

This shows that the vector v_0 considered above is unique, since all $w \in \tilde{D}_m - v_0$ satisfy $\mu(w, v_0) < 1$. So we have a vector field X on M whose value at $m \in M$ is




Fig. 3. (Note that $\partial(\exp D_m)$ is perpendicular to the radius γ)

Considering the compact set $\{\tilde{w} \in J_{\tilde{m}} \mid \mu(\tilde{w} - \tilde{X}(\tilde{m}), \tilde{w} - \tilde{X}(\tilde{m})) = 1 \text{ and } \mu(\tilde{w} - \tilde{X}, I_{\tilde{m}}) = 0\} = C_{\tilde{m}}$ and the open union $\bigcup_{\tilde{w} \in C_{\tilde{m}}} H_{\exp \tilde{w}} \subset \tilde{M}$ we see that $E(I_{\tilde{m}})$ is locally constant. Thus $I = E(I_{\tilde{m}})$ is constant and always outside $\text{im } D$. Since I is intrinsically described, it is clearly invariant under $\phi(\pi_1 M)$.




Fig. 6.

We now show I is a point. The vectorfield X on M is seen to correspond to the vector field Y on E^m which assigns to each $x \in E^m$ the vector $Y(x)$ from x to I which is perpendicular to I . It is now easily checked that $\text{div}_\mu X = \dim(I)$. Since $\text{Vol}_\mu(M)$ is finite we must have $\dim I = 0$. Q.E.D.

The preceding argument raises the hope that incompleteness may be a useful property in the study of affine manifolds.

2. The classification

Using Theorem 1, we may classify similarity manifolds in terms of the well-understood Riemannian manifolds of constant nonnegative curvature [W].

THEOREM 2. *A closed connected similarity manifold M is either Euclidean (if it is complete) or radiant (if it isn't complete). All radiant similarity manifolds M are constructed in the following ways.*

(1) $d \leq 2$: \tilde{M} is \mathbf{R}^+ or $\tilde{\mathbf{C}}^*$. Via e^z , the action of $\pi_1 M$ on \tilde{M} by similarities corresponds to a uniform discrete group of rigid motions of \mathbf{R} or \mathbf{C} (translations or glide reflections), in the coordinate z in \mathbf{R} or \mathbf{C} . This change of coordinates associates a Euclidean manifold to M in a natural way.

(2) $d \geq 3$: M is the quotient of $E^d - 0$ by a group $K \rtimes \mathbf{Z}$, where $K \subset O(d)$ is finite and \mathbf{Z} is generated by an expansion $g \in \text{Sim}_0(E^d)$.

Topologically, radiant similarity manifolds are precisely the mapping tori of isometries of Clifford-Klein spaces. For $d \geq 3$, this isometry and the volume of M (relative to the natural metric μ introduced in Theorem 1) are natural invariants that classify M , and the volume can be freely chosen in \mathbf{R}^+ . The value of $\text{Vol}_\mu(M)$ for $d \geq 3$ is

$$\frac{(\log |\det g|)(\text{vol } S^{d-1})}{d(\text{card } K)}.$$

All similarity manifolds are finitely covered by T^d or $S^{d-1} \times S^1$.

Proof. When M is Euclidean the Hopf–Rinow Theorem shows M is complete. If M is complete then $M = E^d / \pi_1 M$ so $\pi_1 M$ acts without fixed points, the holonomy contains only isometries and M is Euclidean [K]. Theorem 1 shows that incomplete M are radiant. Conversely when M is radiant, the fixed point $e \in E^d$ lies outside the developing image $\text{im}(D)$ (see [FGH, Thm. 3.3] for a correct proof) and so M isn't complete.

When M is radiant, we choose $e \in E^d$ as origin and find that the holonomy preserves the complete metric $ds/\|x\|$ on $E^d - 0$. This induces a metric on M which is clearly just the metric μ constructed in Theorem 1.

It's known [e.g. Th, Thm. 3.6] that when G is a group of analytic isometries of a complete simply connected Riemannian manifold X and M is a closed (G, X) manifold the development map $D: \tilde{M} \rightarrow X$ is a homeomorphism. Let G be $\widetilde{\text{Sim}}_0(E^d)$. Choosing X appropriately, we find that \tilde{M} is \mathbf{R}^+ , $\tilde{\mathbf{C}}^*$ or $E^d - 0$ ($d \geq 3$).

It remains to analyze the action of $\pi_1 M$ on \tilde{M} . The case $d \leq 2$ is easy, since the only fixed point free rigid motions of \mathbf{R} and \mathbf{R}^2 are translations and glide reflections. For $d \geq 3$, the subgroup $\alpha(\pi_1 M) \subset \mathbf{R}^+$ must be discrete for $\pi_1 M$ to act discontinuously (since $O(d) = \widetilde{O}(d)$ is compact, $\alpha(g)$ near 1 implies $\phi(g^n)$ near 1 for some $n > 0$). As $\pi_1 M$ acts uniformly, $\alpha(\pi_1 M)$ is infinite cyclic. The kernel $K = \phi(\pi_1 M) \cap O(d)$ acts properly discontinuously on S^{d-1} and hence is finite. The map $\log \|x\|: E^d - 0 \rightarrow \mathbf{R}$ induces a fibration $f: M \rightarrow S^1$ with fiber the Clifford–Klein space S^{d-1}/K . Transverse to f is the radiant vectorfield \tilde{X} considered in Theorem 1, given in coordinates by $\tilde{X}(\tilde{m}) = -\tilde{m}$. The flow ϕ generated by X permutes the fibers of f and has return map corresponding to the isometry $g/(\alpha(g))$ of S^{d-1}/K .

In dimensions $d \leq 2$, the only mapping tori of Clifford–Klein spaces are S^1 , T^2 and the Klein bottle. These admit Euclidean metrics and e^z gives rise to radiant similarity structures on these manifolds. For $d \geq 3$, an isometry $h: S^{d-1}/K \leftarrow$ of a Clifford–Klein space gives rise to a one-parameter family of radiant similarity structures on the mapping torus M_h as follows. Lift h to $h' \in O(d)$ and let G be the subgroup of $\text{Sim}_0(E^d)$ generated by K and $g = \alpha h'$, $\alpha > 1$. Then G acts properly discontinuously on $E^d - 0$ and quotient is the desired similarity manifold.

The volumes are easily computed by passing to a finite cover to reduce to $K = \{1\}$ and considering the fundamental domain $\{1 \leq \|x\| \leq \alpha(g)\}$. It follows that the parameter $\alpha > 1$ may be replaced by $\text{vol}_\mu(M) > 0$.

Since Bieberbach proved that flat manifolds M are covered by T^d [W], we have shown all portions of Theorem 2. Q.E.D.

We note that the existence of a finite cover with abelian fundamental group allows one to apply [S] and [FGH] to affine structures on M . For instance, when $d \geq 3$ and M is an incomplete closed connected similarity manifold, every affine structure on M is radiant. Also, perturbations of similarity structures within the class of all affine structures are well-understood.

REFERENCES

- [FGH] FRIED, DAVID, GOLDMAN, WILLIAM and HIRSCH, MORRIS. *Affine Manifolds with Nilpotent Holonomy*, to appear.
- [K] KUIPER, N. *Compact Spaces with a Local Structure Determined by the Group of Similarity Transformations in E^m* , Indag. Math. Vol. XII, Fasc. 4, 1950.
- [S] SMILLIE, JOHN. *Affinely Flat Manifolds*. Ph.D. Thesis, Univ. of Chicago, 1977.
- [S-Th] SULLIVAN, DENNIS and THURSTON, W. *Manifolds with Canonical Coordinate Charts: Some Examples*. I.H.E.S. preprint 1979.
- [Th] THURSTON, W. Princeton University notes, 1977.
- [W] WOLF, J. *Spaces of Constant Curvature*, McGraw-Hill, 1967.

University of California at Santa Cruz

Received January 10/July 1 1980